

**WILBERT S. ARONOW, MD**

Adjunct Professor of Geriatrics and Adult Development, Mount Sinai School of Medicine; Corporate Medical Director, Hebrew Hospital Home, Bronx and Westchester County, New York

# Hypertension in elderly patients: Treatment reduces mortality, but is underused

## ■ ABSTRACT

Recently published clinical trials provide compelling evidence of the effectiveness of antihypertensive therapy to reduce cardiovascular mortality. The Sixth Joint National Committee on Detection, Evaluation, and Treatment of Hypertension emphasizes that hypertension is undertreated in this country: only approximately one fourth of hypertensive persons have their blood pressure controlled to the current recommended level of less than 140/90 mm Hg. The point is especially pertinent to elderly patients, most of whom have hypertension. Moreover, most elderly hypertensive persons have systolic hypertension, which has an even stronger association with risk than does diastolic pressure. Current recommendations are reviewed.

## ■ KEY POINTS

Although increased systolic blood pressure is a stronger risk factor for cardiovascular mortality than increased diastolic pressure, increased pulse pressure may be an even better marker of risk.

Office readings may overestimate true average blood pressure owing to pseudohypertension and "white-coat" hypertension.

The goal of treatment in older persons is to lower the blood pressure to < 140/90 mm Hg if possible.

Initial doses of antihypertensive drugs should be lower in elderly than in younger patients, owing to the risk of orthostatic hypotension.

**T**HE OLDER YOU ARE, the more likely you are to have hypertension—and the less likely you are to have it under control.

Government data<sup>1,2</sup> show that by age 60 more than half of the population has hypertension, and more than half of hypertensive persons in this age group have isolated systolic hypertension, shown by epidemiologic data to be a stronger cardiovascular risk factor than diastolic hypertension. Yet, only 26% of hypertensive persons over age 60 have their blood pressure controlled to the currently recommended level of 140/90 mm Hg or less, and at age 75 and older the numbers drop to 21% for women and 17% for men.<sup>2</sup>

Recently published clinical trials provide compelling evidence of the effectiveness of antihypertensive therapy to reduce cardiovascular mortality. This paper summarizes the treatment of hypertension in elderly patients, outlining recent recommendations by the Sixth Joint National Committee on Detection, Evaluation, and Treatment of Hypertension.<sup>1,2</sup>

## ■ MOST OLDER PERSONS HAVE HYPERTENSION

More elderly persons have hypertension than do not. The third National Health and Nutrition Examination Survey<sup>3</sup> found that the prevalence of hypertension (systolic blood pressure  $\geq$  140 mm Hg or diastolic blood pressure  $\geq$  90 mm Hg or under a physician's care for hypertension) in persons aged 60 years and older was:

- 60% in non-Hispanic whites

TABLE 1

### Increased risk in elderly patients with hypertension

ENDPOINT	NO. OF PATIENTS	RELATIVE RISK
<b>New coronary events<sup>4</sup></b>		
Men	664	2.0
Women	1,488	1.6
<b>New atherothrombotic brain infarction<sup>16</sup></b>		
Men	664	2.2
Women	1,488	2.4
<b>Peripheral arterial disease<sup>14</sup></b>		
Men	244	1.7
Women	625	1.5

- 71% in non-Hispanic African Americans
- 61% in Mexican Americans.

Other population studies<sup>4,5</sup> had roughly similar findings.

Isolated systolic hypertension (an elevated systolic pressure with a normal diastolic pressure) accounts for more than half of cases of hypertension in the elderly.<sup>4,5</sup> In contrast, most middle-aged persons with hypertension have diastolic or mixed systolic-diastolic hypertension.

#### ■ HYPERTENSION INCREASES CARDIOVASCULAR MORBIDITY AND MORTALITY IN THE ELDERLY

Isolated systolic hypertension and diastolic hypertension are both associated with increased cardiovascular morbidity and mortality in older persons.<sup>6-17</sup> The higher the systolic or diastolic blood pressure, the greater the morbidity and mortality from cardiovascular disease. Hypertension is the principal risk factor for stroke,<sup>9,16</sup> and it is also associated with coronary events,<sup>4,8</sup> peripheral arterial disease,<sup>10-14</sup> and heart failure (TABLE 1).<sup>9,15</sup>

Of note: Increased systolic blood pressure is a stronger risk factor for cardiovascular morbidity and mortality than is increased diastolic blood pressure.<sup>2,6</sup> The Cardiovascular Health Study, a 5-year study of 5,202 older men and women, found that a brachial systolic blood pressure higher than 169 mm Hg increased the mortality rate 2.4 times.<sup>7</sup>

An even better marker of risk than either systolic or diastolic blood pressure may be increased pulse pressure. Madhavan et al<sup>18</sup> found that a pulse pressure wider than 63 mm Hg was associated with an increase in cardiovascular events. An increased pulse pressure, found in older persons with isolated systolic hypertension, indicates decreased vascular compliance in the large arteries.

#### ■ ANTIHYPERTENSIVE TREATMENT REDUCES STROKES AND CORONARY EVENTS

A number of clinical trials have demonstrated the benefit of treating both diastolic and isolated systolic hypertension in the elderly.<sup>19-25</sup> In general, these studies showed a greater percentage decrease in strokes than in coronary events. MacMahon and Rodgers,<sup>26</sup> in a meta-analysis of five studies in the elderly, calculated that antihypertensive treatment decreased the incidence of strokes by 34% and the incidence of coronary heart disease events by 19%. However, cardiac events are much more common than strokes, and antihypertensive treatment will prevent a greater number of coronary events in terms of absolute numbers.

#### ■ HOW TO DIAGNOSE HYPERTENSION

The report of the Sixth Joint National Committee on Detection, Evaluation, and Treatment of Hypertension (JNC VI)<sup>1</sup> defines hypertension as a systolic blood pressure of 140 mm Hg or higher, or a diastolic blood pressure of 90 mm Hg or higher, or both. The number must be the average of at least two blood pressure readings taken at each of at least two visits after an initial screening visit, using a standardized technique.

#### Stages of hypertension

To gauge the severity of hypertension, JNC VI<sup>1</sup> devised a system of "stages."

**Stage 1.** 140–159 mm Hg systolic, or 90–99 mm Hg diastolic, or both.

**Stage 2.** 160–179 mm Hg systolic, or 100–109 mm Hg diastolic, or both.

**Stage 3.** ≥ 180 mm Hg systolic, or ≥ 110 mm Hg diastolic, or both.<sup>1</sup>

If the systolic and diastolic blood pressures fall into different stages, the higher stage is

**Take at least 2 readings during at least 3 visits, and find the average**



used. Isolated systolic hypertension—a systolic blood pressure of  $\geq 140$  mm Hg with a diastolic blood pressure of  $< 90$  mm Hg—is staged according to the systolic pressure.

### Problems in measuring blood pressure

In some elderly persons, office readings may overestimate the true, average blood pressure, owing to two problems: pseudohypertension and “white-coat” hypertension.

**Pseudohypertension** is due to excessive vascular stiffness—the calcified brachial artery resists compression by the sphygmomanometer, leading to spuriously high readings.<sup>27</sup>

**White-coat hypertension**, ie, an excessive increase in blood pressure in response to the stress of being in a physician’s office, is especially common in older women and particularly affects the systolic blood pressure.<sup>28</sup>

If office readings are high but the patient does not have target organ damage (ie, heart disease, stroke, transient ischemic attacks, nephropathy, peripheral arterial disease, or retinopathy), consider the possibility that the patient has pseudohypertension or white-coat hypertension. In this situation, blood pressure readings should be obtained outside of the office by self-measurement or by ambulatory monitoring.<sup>1</sup>

### Stratifying risk

Hypertension is a cardiovascular risk factor, but not the *only* risk factor. Accordingly, JNC VI also recommends a simple risk-factor analysis, as outlined below.

**Does the patient have other major cardiovascular risk factors?** These are:

- Smoking
- Dyslipidemia
- Diabetes mellitus
- Age older than 60 years
- Male gender
- Postmenopausal status in women
- Family history of cardiovascular disease.

**Does the patient have target organ damage (ie, damage due to high blood pressure) or clinical cardiovascular disease?** These include:

- Left ventricular hypertrophy
- Angina pectoris

- Previous myocardial infarctions
- Previous coronary revascularizations
- Heart failure
- Stroke or transient cerebral ischemic attack
- Nephropathy
- Peripheral arterial disease
- Retinopathy.

Using this information, you can divide patients into three risk groups:

**Risk group A.** No other risk factors and no target organ damage or clinical cardiovascular disease

**Risk group B.** One or more risk factors (not including diabetes mellitus) and no target organ damage or clinical cardiovascular disease

**Risk group C.** Diabetes (with or without other risk factors) or target organ damage or both.

The urgency of beginning antihypertensive drug therapy depends on both the stage of hypertension and the patient’s risk group.

### ■ TREATMENT

The goal of treatment of hypertension in older persons is the same as in younger persons—to decrease the blood pressure to  $< 140/90$  mm Hg if possible.<sup>1</sup> Further reductions in diastolic pressure seem to be safe,<sup>29</sup> although the benefit is not proven.

Although the benefits of antihypertensive treatment in older persons with stage 1 isolated systolic hypertension have not yet been demonstrated, these persons are at increased risk for cardiovascular morbidity and mortality,<sup>30</sup> and I believe they should be treated with antihypertensive therapy, especially if they are in risk groups B or C.

### Lifestyle modification

Like younger patients, older persons with hypertension should undertake lifestyle modifications. Specifically, they should be instructed to:

- Restrict dietary sodium to 2.4 grams daily
- Lose weight if obese
- Discontinue drugs that increase blood pressure
- Avoid tobacco

**A simple risk-factor analysis helps determine whether to start treatment**

- Limit alcohol intake to no more than 1 ounce daily in men and no more than one-half ounce daily in women and lighter-weight persons
- Increase aerobic physical activity
- Reduce dietary saturated fat and cholesterol
- Maintain adequate dietary potassium, calcium, and magnesium intake.

### When to start drug therapy

Lifestyle modifications alone may suffice for some patients. JNC VI<sup>1</sup> recommends trying lifestyle modification for up to 6 months in persons with stage 1 hypertension in risk group B, and starting antihypertensive drug therapy if this does not reduce the blood pressure to below 140/90 mm Hg.

However, older persons should start antihypertensive drug therapy immediately in addition to lifestyle modifications if they have:

- Stage 2 or 3 hypertension
- Stage 1 hypertension and target organ damage or evidence of clinical cardiovascular disease or diabetes mellitus.

### What drug to use?

JNC VI recommends diuretics or beta-blockers as initial drugs,<sup>1</sup> because these drugs have been found to decrease cardiovascular morbidity and mortality in controlled clinical trials,<sup>19–22</sup> whereas few trials have been performed with other classes of agents. I agree with this recommendation.

Older persons with hypertension should receive another drug as initial therapy if they also have any of the conditions listed below:

- Previous myocardial infarction, angina pectoris, myocardial ischemia, or complex ventricular arrhythmias: a beta-blocker is indicated.<sup>31</sup>

- Heart failure: both a diuretic and an angiotensin-converting enzyme (ACE) inhibitor are indicated.<sup>32,33</sup>

- Diabetes mellitus: an ACE inhibitor is indicated. If the patient cannot tolerate an ACE inhibitor because of cough, rash, or altered taste sensation, an angiotensin II type 1 receptor antagonist should be given. Alternatively, a long-acting dihydropyridine can be effective.<sup>34</sup>

The JNC VI report includes tables show-

ing the range of doses of different antihypertensive drugs.<sup>1</sup> The lower doses listed in these tables should be used in treating older persons (TABLE 2).

### ■ ISOLATED SYSTOLIC HYPERTENSION

Persons with isolated systolic hypertension are the largest subgroup of older persons with hypertension. This group also represents the largest population group inadequately treated at the present time. In an academic geriatrics practice, 30% of persons with hypertension did not have their blood pressure lowered to below 140/90 mm Hg by antihypertensive therapy, and the percentage was the same in patients with isolated systolic hypertension.<sup>5</sup>

The benefit of treating isolated systolic hypertension is well documented. In the Systolic Hypertension in the Elderly Program, isolated systolic hypertension was defined as a systolic blood pressure  $\geq 160$  mm Hg and the diastolic blood pressure  $\leq 90$  mm Hg.<sup>22</sup> In this randomized, double-blind, placebo-controlled study, chlorthalidone 12.5 mg to 25 mg daily was used as step 1 drug therapy and atenolol 25 mg to 50 mg daily as step 2. At 4.5-years of follow-up, compared with placebo, antihypertensive drug therapy caused a 36% significant reduction in stroke, a 27% significant decrease in nonfatal myocardial infarction plus coronary death, a 33% significant reduction in major cardiovascular events, a 49% significant decrease in heart failure,<sup>25</sup> and a 13% insignificant decrease in total mortality.

The Systolic Hypertension in Europe (Syst-Eur) Trial<sup>23</sup> used a slightly different definition of isolated systolic hypertension: a sitting systolic blood pressure  $\geq 160$  mm Hg with a sitting diastolic blood pressure  $< 95$  mm Hg, and a standing systolic blood pressure  $\geq 140$  mm Hg. In this randomized, double-blind, placebo-controlled study in older patients, active treatment was started with nitrendipine 10 mg to 40 mg daily and, if necessary, combined with or replaced by enalapril 5 mg to 20 mg daily, hydrochlorothiazide 12.5 mg to 25 mg daily, or both. At 2 years of follow-up, compared with placebo, antihypertensive drug therapy caused a 42% significant reduction in stroke and a 31% significant reduction in all fatal and nonfatal cardiovascular endpoints. In

**Start drug therapy immediately in stage 2 or 3 hypertension**

**TABLE 2****Initial dosages of antihypertensive agents in elderly patients****Thiazide diuretics**

Bendroflumethiazide 2.5 mg daily  
Benzthiazide 12.5 mg daily  
Chlorothiazide 75 mg twice a day  
Chlorthalidone 12.5 mg daily  
Hydrochlorothiazide 12.5 mg daily  
Hydroflumethiazide 12.5 mg daily  
Indapamide 2.5 mg daily  
Methyclothiazide 2.5 mg daily  
Metolazone 0.5 mg daily  
Polythiazide 1 mg daily  
Quinethazone 25 mg daily  
Trichlormethiazide 1 mg daily

**Loop diuretics**

Bumetanide 0.25 mg twice a day  
Ethacrynic acid 12.5 mg twice a day  
Furosemide 20 mg twice a day  
Torsemide 5 mg daily

**Potassium-sparing diuretics**

Amiloride 5 mg daily  
Spironolactone 25 mg daily  
Triamterene 25 mg daily

**Alpha 1 receptor blockers**

Doxazosin 1 mg daily  
Prazosin 1 mg twice a day  
Terazosin 1 mg daily

**Beta-blockers**

Acebutolol 200 mg daily  
Atenolol 25 mg daily  
Betaxolol 5 mg daily  
Bisoprolol 2.5 mg daily  
Carteolol 2.5 mg daily  
Metoprolol 50 mg daily  
Nadolol 40 mg daily  
Penbutolol 10 mg daily  
Pindolol 5 mg twice a day  
Propranolol 20 mg twice a day  
Propranolol, long-acting 40 mg daily  
Timolol 10 mg twice a day

**Alpha-beta blockers**

Carvedilol 6.25 mg twice a day  
Labetalol 100 mg twice a day

**Other adrenergic inhibitors**

Clonidine 0.1 mg twice a day  
Guanabenz 4 mg twice a day  
Guanadrel 5 mg twice a day  
Guanethidine 10 mg daily  
Guanfacine 1 mg daily  
Methyldopa 250 mg twice a day  
Reserpine 0.05 mg daily

**ACE inhibitors**

Benazepril 5 mg daily  
Captopril 12.5 mg twice a day  
Enalapril 5 mg daily  
Fosinopril 10 mg daily  
Lisinopril 5 mg daily  
Moexipril 3.75 mg twice a day  
Quinapril 5 mg daily  
Ramipril 1.25 mg daily  
Trandolapril 1 mg daily

**Angiotensin II receptor blockers**

Irbesartan 125 mg daily  
Losartan 25 mg daily  
Valsartan 80 mg daily

**Calcium antagonists**

Amlodipine 2.5 mg daily  
Diltiazem, sustained release 60 mg twice a day  
Diltiazem, extended release 120 mg daily  
Felodipine 5 mg daily  
Isradipine 1.25 mg twice a day  
Nicardipine, sustained release 30 mg twice a day  
Nifedipine, long-acting 30 mg daily  
Nisoldipine 20 mg daily  
Verapamil 40 mg twice a day  
Verapamil, long-acting 120 mg daily

SOURCE: ADAPTED FROM THE JNC VI REPORT, REFERENCE 1

addition, there were 14% fewer deaths, 20% fewer myocardial infarctions, 29% fewer cases of heart failure, and 27% fewer deaths due to cardiovascular diseases, but these trends did not achieve statistical significance.

In these studies, no J-curve was clearly defined for either systolic or diastolic blood pressure, ie, no point beyond which further blood pressure reduction was associated with an increase in adverse events.

The JNC VI report recommends thiazide diuretics or beta-blockers in combination with diuretics for treating older persons with isolated systolic hypertension, based on the results from the SHEP study.<sup>22</sup> In addition, because of data from the Syst-Eur trial,<sup>23</sup> the JNC VI report also recommends long-acting dihydropyridine calcium antagonists for this indication.<sup>1</sup>

**■ LEFT VENTRICULAR HYPERTROPHY**

Left ventricular hypertrophy (LVH) is an independent risk factor for coronary events and stroke. In two studies in elderly hypertensive patients in our clinic,<sup>35,36</sup> we found that LVH increased the risk of coronary events approximately threefold, and the risk of atherothrombotic brain infarction between threefold and fourfold.

A meta-analysis of 109 treatment studies showed that ACE inhibitors were more effective than other antihypertensive drugs in reducing left ventricular mass.<sup>37</sup> However, other types of antihypertensive drugs also reduce left ventricular mass. In an echocardiographic substudy of SHEP, at 3-year follow-up, the left ventricular mass index had decreased by 13% in the group receiving

TABLE 3

### Reasons to look for secondary hypertension in elderly patients

Onset of diastolic blood pressure > 100 mm Hg after age 60 years  
 Exacerbation of hypertension or resistance to a previously effective drug regimen  
 Accelerated hypertension  
 Diastolic blood pressure > 100 mm Hg despite triple drug therapy  
 Increasing renal dysfunction  
 Symptoms suggesting pheochromocytoma or other curable types of hypertension  
 Spontaneous hypokalemia

active drug treatment (a diuretic-based regimen) and had increased by 6% in the placebo group.<sup>38</sup>

Prospective studies using different types of antihypertensive drugs are necessary to determine whether regression of left ventricular mass leads to a decrease in cardiovascular morbidity and mortality in older men and women with hypertension.

### Renovascular disease is the most common cause of secondary hypertension in the elderly


#### ■ ORTHOSTATIC AND POSTPRANDIAL HYPOTENSION

Antihypertensive drug therapy may cause orthostatic hypotension<sup>39</sup> or postprandial hypotension<sup>40</sup> in older persons, especially in those who are frail or institutionalized. Management of orthostatic and postprandial hypotension in older persons is discussed in detail elsewhere.<sup>41</sup> The dose of antihypertensive drug may need to be decreased or another antihypertensive drug used.

#### ■ SECONDARY HYPERTENSION

Most cases of hypertension in the elderly are primary (ie, idiopathic), but a sizable minority may be secondary (ie, due to an identifiable cause). TABLE 3 lists the reasons for looking for secondary hypertension in older persons. The most common form of secondary hypertension in older persons is atherosclerotic renovascular disease. Using renal duplex ultrasonography, we found proximal renal artery stenosis ( $\geq 60\%$  narrowing of the luminal diameter of

the renal artery) in 18 (22%) of 81 older hypertensive patients (mean age 82 years).<sup>42</sup> Bilateral renal artery stenosis was present in 5 (6%) of the 81 patients.

Medical therapy should be used for treating renovascular hypertension in older persons if blood pressure is controlled and renal function remains stable. Surgical revascularization or percutaneous transluminal renal angioplasty and stent placement should be considered for patients with renovascular hypertension that is refractory to medical therapy. Surgical revascularization may also preserve renal function. 

#### ■ REFERENCES

- 1997 Joint National Committee. The Sixth Report of the Joint National Committee on the Detection, Evaluation, and Treatment of High Blood Pressure (JNC VI). *Arch Intern Med* 1997; 157:2413–2444.
- National High Blood Pressure Education Program Working Group. National High Blood Pressure Education Program working group report on hypertension in the elderly. *Hypertension* 1994; 23:275–285.
- Burt VL, Whelton P, Roccella EJ, et al. Prevalence of hypertension in the US adult population: results from the third National Health and Nutrition Examination Survey, 1988–1991. *Hypertension* 1995; 25:305–313.
- Aronow WS, Ahn C. Risk factors for new coronary events in a large cohort of very elderly patients with and without coronary artery disease. *Am J Cardiol* 1996; 77:864–866.
- Mendelson G, Ness J, Aronow WS. Drug treatment of hypertension in older persons in an academic hospital-based geriatrics practice. *J Am Geriatr Soc* 1999; 47:597–599.
- Applegate WB, Rutan GH. Advances in management of hypertension in older persons. *J Am Geriatr Soc* 1992; 40:1164–1174.
- Fried LP, Kronmal RA, Newman AB, et al. Risk factors for 5-year mortality in older adults. The Cardiovascular Health Study. *JAMA* 1998; 279:585–592.
- Vokonas PS, Kannel WB. Epidemiology of coronary heart disease in the elderly. In: Tresch DD, Aronow WS, editors. *Cardiovascular Disease in the Elderly Patient*. New York: Marcel Dekker, Inc 1994:91–123.
- Wolf PA. Cerebrovascular disease in the elderly. In: Tresch DD, Aronow WS, editors. *Cardiovascular Disease in the Elderly Patient*. New York: Marcel Dekker, Inc, 1994:125–147.
- Kannel WB. Hypertension. In: Aronow WS, Stemmer EA, Wilson SE, editors. *Vascular disease in the elderly*. Armonk: Futura Publishing Co, Inc, 1997:177–198.
- Juergens JL, Barker NW, Hines EA Jr. Arteriosclerosis obliterans: review of 520 cases with special reference to pathogenic and prognostic factors. *Circulation* 1960; 21:188–195.
- Hughson WG, Mann JI, Garrod A. Intermittent claudication: prevalence and risk factors. *Br Med J* 1978; 1:1379–1381.
- Schroll M, Munck O. Estimation of peripheral arteriosclerotic disease by ankle blood pressure measurements in a population study of 60-year-old men and women. *J Chron Dis* 1981; 34:261–269.



14. Aronow WS, Sales FF, Etienne F, Lee NH. Prevalence of peripheral arterial disease and its correlation with risk factors for peripheral arterial disease in elderly patients in a long-term health care facility. *Am J Cardiol* 1988; 62:644–646.
15. Levy D, Larson MG, Vasan RS, et al. The progression from hypertension to congestive heart failure. *JAMA* 1996; 275:1557–1562.
16. Aronow WS, Ahn C, Gutstein H. Risk factors for new atherothrombotic brain infarction in 664 older men and 1,488 older women. *Am J Cardiol* 1996; 77:1381–1383.
17. Garland C, Barrett-Connor E, Suarez L, Criqui MH. Isolated systolic hypertension and mortality after age 60 years. *Am J Epidemiol* 1983; 118:365–376.
18. Madhavan S, Ooi WL, Cohen H, Alderman MH. Relation of pulse pressure and blood pressure reduction to the incidence of myocardial infarction. *Hypertension* 1994; 23:395–401.
19. Amery A, Birkenhager W, Brixko P, et al. Mortality and morbidity results from the European Working Party on Hypertension in Elderly Trial. *Lancet* 1985; 1:1349–1354.
20. Dahlöf B, Lindholm LH, Hansson L, et al. Morbidity and mortality in the Swedish Trial in Old Patients With Hypertension (STOP Hypertension). *Lancet* 1991; 338:1281–1285.
21. MRC Working Party. Medical Research Council trial of treatment of hypertension in older adults: principal results. *Br Med J* 1992; 304:405–412.
22. SHEP Cooperative Research Group. Prevention of stroke by antihypertensive drug treatment in older persons with isolated systolic hypertension: final results of the Systolic Hypertension in the Elderly Program (SHEP). *JAMA* 1991; 265:3255–3264.
23. Staessen JA, Fagard R, Thijs L, et al. Randomised double-blind comparison of placebo and active treatment for older patients with isolated systolic hypertension. *Lancet* 1997; 350:757–764.
24. Coope J, Warrender TS. Randomised trial of treatment of hypertension in the elderly in primary care. *Br Med J* 1986; 293:1145–1151.
25. Kostis JB, Davis BR, Cutler J, et al. Prevention of heart failure by antihypertensive drug treatment in older persons with isolated systolic hypertension. *JAMA* 1997; 278:212–216.
26. MacMahon S, Rodgers A. The effects of blood pressure reduction in older patients: an overview of five randomized controlled trials in elderly hypertensives. *Clin Exp Hypertens* 1993; 15:967–978.
27. Messerli FH. Osler's maneuver, pseudohypertension, and true hypertension in the elderly. *Am J Med* 1986; 80:906–910.
28. Wiinberg N, Hoegholm A, Christensen HR, et al. 24-h ambulatory blood pressure in 352 normal Danish subjects, related to age and gender. *Am J Hypertens* 1995; 8:978–986.
29. Hansson L, Zanchetti A, Carruthers SG, et al. Effects of intensive blood-pressure lowering and low-dose aspirin in patients with hypertension: principal result of the Hypertension Optimal Treatment (HOT) randomised trial. *Lancet* 1998; 351:1755–1762.
30. Sagie A, Larson MG, Levy D. The natural history of borderline isolated systolic hypertension. *N Engl J Med* 1993; 329:1912–1917.
31. Aronow WS, Ahn C, Mercado AD, et al. Effect of propranolol versus no antiarrhythmic drug on sudden cardiac death, total cardiac death, and total death in patients  $\geq 62$  years of age with heart disease, complex ventricular arrhythmias, and left ventricular ejection fraction  $\geq 40\%$ . *Am J Cardiol* 1994; 74:267–270.
32. Garg R, Yusuf S, for the Collaborative Group on ACE Inhibitor Trials. Overview of randomized trials of angiotensin-converting enzyme inhibitors on mortality and morbidity in patients with heart failure. *JAMA* 1995; 273:1450–1456.
33. Aronow WS, Kronzon I. Effect of enalapril on congestive heart failure treated with diuretics in elderly patients with prior myocardial infarction and normal left ventricular ejection fraction. *Am J Cardiol* 1993; 71:602–604.
34. Tuomilehto J, Rastenyte D, Birkenhager W, et al. Effects of calcium-channel blockade in older patients with diabetes and hypertension. *N Engl J Med* 1999; 340:677–684.
35. Aronow WS, Ahn C, Kronzon I, Koenigsberg M. Congestive heart failure, coronary events and atherothrombotic brain infarction in elderly blacks and whites with systemic hypertension and with and without echocardiographic and electrocardiographic evidence of left ventricular hypertrophy. *Am J Cardiol* 1991; 67:295–299.
36. Aronow WS, Ahn C, Kronzon I, Gutstein H. Association of plasma renin activity and echocardiographic left ventricular hypertrophy with frequency of new coronary events and new atherothrombotic brain infarction in older persons with systemic hypertension. *Am J Cardiol* 1997; 79:1543–1545.
37. Dahlöf B, Pennert K, Hansson L. Reversal of left ventricular hypertrophy in hypertensive patients: a metaanalysis of 109 treatment studies. *Am J Hypertens* 1992; 5:95–110.
38. Ofili EO, Cohen JD, St. Vrain JA, et al. Effect of treatment of isolated systolic hypertension on left ventricular mass. *JAMA* 1998; 279:778–780.
39. Aronow WS, Lee NH, Sales FF, Etienne F. Prevalence of postural hypotension in elderly patients in a long-term health care facility. *Am J Cardiol* 1988; 62:336.
40. Aronow WS, Ahn C. Postprandial hypotension in 499 elderly persons in a long-term health care facility. *J Am Geriatr Soc* 1994; 42:930–932.
41. Aronow WS. Dizziness and syncope. In: Hazzard WR, Blass JP, Ettinger WH Jr, Halter JB, Ouslander JG, editors. *Principles of Geriatric Medicine and Gerontology* (4th ed). New York City: McGraw-Hill, Inc, 1998:1519–1534.
42. Schoenfeld MR, Aronow WS, Kutcher R, et al. The incidence of atheromatous renovascular disease in the elderly diagnosed by duplex ultrasonography. *J Cardiovasc Tech* 1989; 8:283–286.

ADDRESS: Wilbert S. Aronow, MD, Hebrew Hospital Home, 801 Co-op City Boulevard, Bronx, NY 10475.

## We Welcome Your Letters

WE ENCOURAGE YOU TO WRITE, either to respond to an article published in the *Journal* or to address a clinical issue of importance to you. You may submit letters by mail, fax, or e-mail.

### MAILING ADDRESS

Letters to the Editor  
*Cleveland Clinic Journal of Medicine*  
9500 Euclid Ave., NA32  
Cleveland, OH 44195

### FAX

216.444.9385

E-MAIL [ccjm@ccf.org](mailto:ccjm@ccf.org)

Please be sure to include your full address, phone number, fax number, and e-mail address. Please write concisely, as space is limited. Letters may be edited for style and length. We cannot return materials sent. Submission of a letter constitutes permission for the *Cleveland Clinic Journal of Medicine* to publish it in various editions and forms.