

To the Editor: I read with interest the review article by Yahr et al¹ in the February issue, about resistant hypertension. Under the heading “Does the patient have lifestyle factors that raise blood pressure?” there is no mention of stress (chronic or acute crisis) or of psychiatric conditions such as generalized anxiety disorder. In treating the whole patient, these factors should be considered.

An example: a thin 77-year-old White man with no chronic illnesses other than moderate hypertension, well-controlled until 6 months previous to pre-senting with 180/95 mm Hg pressures (despite adher-

ing to diet and medications), and with a normal lipid profile and normal renal and hepatic function. On questioning, the patient says his spouse was diagnosed with dementia and a movement disorder 8 months before this visit. The patient is the sole caregiver, doing all cooking, cleaning, care related to activities of daily living, and supervision for a “nocturnal wanderer” spouse in her late 70s. He reports 4 to 5 hours of uninterrupted sleep daily.

This is why I think that changes in relationships and environment should be questioned.

Otherwise, an excellent article.

Leslie E. F. Page, DO, MPH
Wichita, KS

■ REFERENCES

1. Yahr J, Thomas G, Calle J, Taliencio J. Resistant hypertension: a step-wise approach. *Cleve Clin J Med* 2023; 90(2):115–125.
doi:10.3949/ccjm.90a.22046

doi:10.3949/ccjm.90c.05002