

## Why 25-dehydroxyvitamin D is a negative acute-phase reactant

**To the Editor:** In previous publications, we presented evidence that 25-dehydroxyvitamin D (25[OH]D) in serum behaves as a negative acute-phase reactant, ie, that its concentration decreases in the presence of inflammatory states.<sup>1,2</sup> Low levels may thus reflect low vitamin D stores or inflammation, complicating the clinical interpretation of test results.

We have recently become aware of the mechanism underlying this phenomenon: less than 1% of circulating 25(OH)D exists in unbound form, and the majority is tightly bound to vitamin D binding protein, while 10% to 15% is bound to albumin. Both are negative acute-phase proteins.<sup>3–5</sup> As the serum concentrations of these proteins decrease, so does that of 25(OH)D. Similarly, the positive acute-phase behavior of copper is explained by the fact that it is bound to ceruloplasmin, a positive acute-phase protein.<sup>6</sup>

Maria J. Antonelli, MD  
Assistant Professor of Medicine,  
Department of Medicine,  
Division of Rheumatology,  
Case Western Reserve University,  
MetroHealth Medical Center, Cleveland, OH

Irving Kushner, MD  
Emeritus Professor of Medicine,  
Department of Medicine,  
Division of Rheumatology,  
Case Western Reserve University,  
MetroHealth Medical Center, Cleveland, OH

Murray Epstein, MD, FASN, FACP  
Emeritus Professor of Medicine,  
Division of Nephrology and Hypertension,  
University of Miami Miller School of Medicine,  
Miami, FL

### REFERENCES

1. Antonelli M, Kushner I. Low serum levels of 25-hydroxyvitamin D accompany severe COVID-19 because it is a negative acute phase reactant. *Am J Med Sci* 2021; 362(3):333–335. doi:10.1016/j.amjms.2021.06.005
2. Antonelli MJ, Kushner I, Epstein M. The constellation of vitamin D, the acute-phase response, and inflammation. *Cleve Clin J Med* 2023; 90(2):85–89. doi:10.3949/ccjm.90a.22048
3. Yousefzadeh P, Shapses SA, Wang X. Vitamin D binding protein impact on 25-hydroxyvitamin D levels under different physiologic and pathologic conditions. *Int J Endocrinol* 2014; 2014:981581. doi:10.1155/2014/981581
4. Dahl B, Schiødt FV, Gehrchen PM, Ramlau J, Kiaer T, Ott P. Gc-globulin is an acute phase reactant and an indicator of muscle injury after spinal surgery. *Inflamm Res* 2001; 50(1):39–43. doi:10.1007/s000110050722
5. Liberman U, Bikle DD. Disorders in the action of vitamin D. In: Feingold KR, Anawalt B, Blackman MR, et al, eds. *Endotext*. South Dartmouth, MA: MDText.com, Inc.; 2023.
6. Sattar N, Scott HR, McMillan DC, Talwar D, O'Reilly DS, Fell GS. Acute-phase reactants and plasma trace element concentrations in non-small cell lung cancer patients and controls. *Nutr Cancer* 1997; 28(3):308–312. doi:10.1080/01635589709514592

doi:10.3949/ccjm.90c.09001