# Disordered immune regulation in autoimmune rheumatic diseases

Autologous mixed lymphocyte reactivity in polymyositis/dermatomyositis and systemic lupus erythematosus<sup>1</sup>

Meher M. Dustoor, Ph.D. Richard Ransohoff, M.D. Allen M. Segal, D.O.

<sup>1</sup> Departments of Immunology and Cancer Research (M.M.D.), Neurology (R.R.), and Rheumatic and Immunologic Diseases (A.M.S.), The Cleveland Clinic Foundation. Submitted for publication Jan 1986; accepted April 1986.

#### 0009-8787/86/03/0241/08/\$3.00/0

Copyright © 1986, The Cleveland Clinic Foundation

Polymyositis/dermatomyositis (PM/DM) and systemic lupus erythematosus (SLE) are autoimmune disorders of unknown etiology. In order to study whether immunoregulatory abnormalities might be involved in these autoimmune states, the autologous mixed lymphocyte reaction (AMLR) was investigated in patients with either adult PM/DM, childhood DM (CDM), or SLE. The AMLR was found to be significantly depressed in adult PM/DM regardless of disease activity. By contrast, in CDM the AMLR was normal. In SLE, the AMLR was depressed in patients with both active and inactive disease and the responding T cell population appeared to be defective, as shown in a study on a pair of identical twins. These studies thus underscore the existence of a primary immunoregulatory dysfunction in these two autoimmune diseases.

**Index terms:** Dermatomyositis • Lupus erythematosus, systemic

**Cleve Clin Q 53:241–248**, Fall 1986

The autologous mixed lymphocyte reaction (AMLR) is the *in vitro* proliferative response of T cells to autologous non-T or activated T cells.<sup>1,2</sup> The response is directed against Class II histocompatibility antigens present on the stimulating cells.<sup>3</sup> The AMLR is capable of generating various immunoregulatory activities including both T cell helper and suppressor function, broadly reactive cytotoxic cells, and lymphokines such as interleukin-2 and interferon- $\gamma$ .<sup>4-10</sup> The AMLR thus represents an *in vitro* model for dissecting cellular and molecular interactions leading to specific immunological functions and can be used to analyze interactions occurring in an aberrant fashion, for example, in autoimmune disorders and malignancy. Support for this rationale has in fact been obtained in studies of the AMLR in various diseases where it has been shown to be defective.<sup>11-15</sup>

Both polymyositis/dermatomyositis (PM/DM) and systemic lupus erythematosus (SLE) are rheumatic diseases in which autoimmunity plays a direct pathologic role. Muscle destruction in PM/ DM is the result of a cell-mediated immune reaction to autologous skeletal muscle<sup>16</sup> whereas in SLE, humoral mechanisms, in the form of antigen-antibody complexes deposited in various tissues, are primarily involved in the pathogenesis.<sup>17</sup> In autoimmune diseases, the presence of an underlying disorder of immunoregulation contributing to the pathogenesis is an important consideration. In this article, we report the results of our study of AMLR reactivity in adult and childhood PM/DM and in SLE.

# Methods

# Patients

Adult polymyositis/dermatomyositis: Diagnosis was based on the criteria of Bohan et al<sup>18</sup>; all patients included in the study had "probable" or "definite" disease by these criteria. Patients with malignancy, overlap syndromes (defined as the presence of a second established rheumatologic diagnosis), or childhood DM were excluded. All patients whose disease was described as "active" were studied after diagnosis was established and before treatment was begun. These patients had elevated muscle enzyme levels and weakness at the time of study. All, however, were ambulatory and without intercurrent illness. Patients whose disease was described as "inactive" were studied upon return to their physicians for routine checkup. These latter patients were in clinical and biochemical remission at the time of study. Patients with inactive disease and receiving doses of corticosteroid greater than 7.5 mg/day prednisone equivalent were excluded; patients (Patient 6 and Patient 7) receiving smaller doses of corticosteroid (ie, 7.5 mg/day or less) were studied at least 24 hours after a dose of medication, since it has been demonstrated that the AMLR returns to normal levels within 24 hours of administration of an oral dose of prednisone.<sup>19</sup> Patients receiving cytotoxic immunosuppressive medications were excluded. No patients studied had light- or electron-microscopic criteria for the diagnosis of inclusion body myositis.

Childhood dermatomyositis: All patients with childhood dermatomyositis (CDM) fulfilled the criteria of Bohan et al.<sup>18</sup> Other conditions for inclusion in the study were the same as described for patients with adult PM/DM. Patient 1 was being treated with a 2 mg/day course of prednisone.

Systemic lupus erythematosus: All SLE patients met the American Rheumatism Association 1982 revised criteria for the classification of SLE.<sup>20</sup> None of the patients was receiving cytotoxic drugs or corticosteroids >7.5 mg/day at the time of the study. As with PM/DM patients, those receiving corticosteroids were studied at least 24 hr after a dose of medication. Of the 16 patients studied, six were receiving an alternate-day course of prednisone (7.5 mg or less). Active or inactive disease was defined by using a modification of a quantitative scoring system of flare severity previously published by Lokshin et al.<sup>21</sup>

#### Laboratory methods

Preparation of lymphocytes and purification of T cells and non-T cells: Peripheral blood mononuclear cells were isolated by centrifugation on a Ficoll-Hypaque (Pharmacia Fine Chemicals, Piscataway, NJ) gradient. T and non-T cells were separated by rosette technique with sheep erythrocytes (SRBC) using an overnight procedure. T cells recovered from the rosette pellet by lysis with 0.14 M NH<sub>4</sub>Cl were used as responder cells and were 95–97% E-rosette positive. Non-T fractions contained 55–65% SIg-positive cells, 25– 35% esterase-positive cells, and less than 5% Erosette-positive cells.

Fractionation of T cells into T4 and T8 subpopulations: This was accomplished by use of a panning previously described procedure.<sup>4</sup> Briefly,  $30-40 \times 10^6$  T cells were treated with optimal dilutions of either OKT4 or OKT8 monoclonal antibodies (Ortho Diagnostics, Raritan, NJ) for 45 min at 4° C. The treated cells were allowed to adhere for 1 hour at 4° C on plates coated with goat anti-mouse IgG (Zymed Labs, San Francisco, CA). Nonadherent cells, after OKT8 or OKT4 treatment, were generally >90% T4 or >90% T8, respectively. Contamination with T8 or T4 cells, respectively, was <2%.

Autologous mixed lymphocyte reaction: A total of  $1 \times 10^5$  T cells was cultured with  $1 \times 10^5$ mitomycin C-treated (50 µg mitomycin C/2 ×  $10^6$  cells/mL) non-T cells in RPMI-1640 medium

			<sup>8</sup> H-thymidine incorpo cont	pration (cpm) in culture aining		
Subject*	Age (yr) Age (yr)	Activity of disease†	Responders alone	Responders + stimulators	Δ СРМ‡	
Patient 1	21	I	$7,965 \pm 3,823$	$17,098 \pm 2,827$	9,133	
Control 1			$750 \pm 568$	$10,012 \pm 6,738$	9,262	
Patient 2	23	А	$3,060 \pm 591$	$4,028 \pm 2,865$	968	
Control 2			$4,528 \pm 888$	$20,202 \pm 3,115$	15,674	
Patient 3	45	I	$363 \pm 228$	$1,145 \pm 261$	782	
Control 3			$62 \pm 74$	$1,072 \pm 596$	1,010	
Patient 4	47	Α	$3,185 \pm 233$	$11,646 \pm 1,135$	8,461	
Control 4			$3,924 \pm 314$	$14,546 \pm 1,668$	10,622	
Patient 5	50	Α	$800 \pm 80$	$1,038 \pm 345$	238	
Patient 6	55	Ι	$698 \pm 260$	$329 \pm 193$	-369	
Control 5			$1,148 \pm 299$	$3,893 \pm 36$	2,745	• •
Patient 7	57	Α	$1,669 \pm 265$	$425 \pm 263$	-1,244	.*
Control 6			$2,954 \pm 188$	$6,934 \pm 226$	3,980	
Patient 8	62	I	$406 \pm 468$	$211 \pm 13$	-195	
Control 6			$2,954 \pm 188$	$6,934 \pm 226$	3,980	
Patient 9	67	I	$730 \pm 670$	$1,088 \pm 722$	358	
Control 7			$594 \pm 389$	$2,253 \pm 658$	1,659	
Patient 10	70	I	$472 \pm 161$	$1,704 \pm 273$	1,232	
Control 10			$407 \pm 150$	$9,676 \pm 615$	9,269	
Patient 11	58	А	$52 \pm 13$	$49 \pm 9$	3	

# Table 1. Autologous mixed lymphocyte reaction in patients with polymyositis/dermatomyositis and matched controls

\* Normal controls were age-, sex-, and race-matched with each patient and were tested concurrently.

 $\dagger$  Assessed as described under Materials and Methods. A = active disease; I = inactive disease.

 $\ddagger$  Significantly depressed in patients (p < 0.01, Wilcoxon signed rank test for paired data).

supplemented with 10% heat-inactivated pooled human AB serum, 1 mM l-glutamine, 25 mM Hepes buffer, 100 U/mL penicillin, and 100  $\mu$ g/ mL streptomycin. Triplicate cultures were carried out in round-bottom microtiter plates (Linbro Chemical Co., Hamden, CT) in 5% CO<sub>2</sub> and 95% air at 37° C. On day 5, 0.5  $\mu$ Ci (1.85 × 10<sup>4</sup> Bq) <sup>3</sup>H-thymidine (3 Ci/mMol [1.11 × 10<sup>5</sup> MBq/ mMol], New England Nuclear, Boston, MA) was added per well and the cultures were harvested 16 hours later using a MASH Harvester. Uptake of <sup>3</sup>H-thymidine was measured by liquid scintillation counting. The results are expressed as  $\Delta$ counts per minute ( $\Delta$  cpm) where

 $\Delta$  cpm = (cpm of AMLR cultures)

- (cpm of responders cultured alone)

### Statistical analysis

In our study, patients were paired with ageand sex-matched normal subjects. Because our study population was not a random sample from the population at large, there was no presumption that our results would follow a normal distribution. All data were therefore analyzed without assumptions about the distribution; the Wilcoxon signed rank test for paired data<sup>22</sup> was used to test for significant differences between patients and controls, and between patients with active and inactive disease. Results were analyzed for normality by means of the Wilk-Shapiro test, and where results were found to follow a normal distribution, the t test for paired data was also performed; there were no differences between results of the Wilcoxon signed rank test and the

**Table 2.** Allogeneic mixed lymphocyte reaction in patients with polymyositis/dermatomyositis

<b>.</b>			
Subject	Responders alone (cpm)	Responders + autologous stimulators* (cpm)	Responders + allogeneic stimulators† (cpm)
Patient 3	$363 \pm 228$	$1,145 \pm 261$	$19,473 \pm 618$
Control 3	$62 \pm 74$	$1,072 \pm 596$	$9,032 \pm 1,370$
Patient 9	$730 \pm 670$	$1,088 \pm 722$	$26,883 \pm 2,266$
Control 8	$594 \pm 389$	$2,253\pm 658$	$41,082 \pm 5,654$

\*  $1 \times 10^5$  responding T cells were cultured with  $1 \times 10^5$  mitomycin-C treated autologous non-T cells for 6 days.

† 1 × 10<sup>5</sup> responding T cells were cultured with 1 × 10<sup>5</sup> mitomycin-C treated allogeneic non-T cells for 6 days.

t test for paired data. All data are presented as analyzed with the Wilcoxon signed rank test.

#### Results

#### Polymyositis/dermatomyositis

The AMLR proliferative response was studied in eleven PM/DM patients and their age-, race-, and sex-matched normal controls. As shown in Table 1, this AMLR proliferative response was significantly depressed in PM/DM patients (p <.01). A time-course study of the AMLR in normal individuals and in PM/DM patients revealed a continuous low reactivity by PM/DM lymphocytes throughout the culture period (data not shown) indicating that altered kinetics does not account for the poor reactivity of PM/DM lymphocytes. Furthermore, addition of increasing numbers of stimulator cells did not reconstitute the response (data not shown). It is important to note that a nonspecific depression of all T lymphocyte functions was not present in these patients, as they were shown to possess reactivity in allogeneic MLR (Table 2).

A comparison of five patients with active disease and six with inactive disease showed a similar impairment of AMLR proliferative responses in both groups.

#### Childhood dermatomyositis

Twelve patients with CDM were evaluated for AMLR reactivity. As shown in *Table 3*, the patients as a group responded well in the AMLR and had no statistically significant differences when compared with normal controls, although three of the 12 patients did demonstrate impaired AMLR responses (Patients 1, 2, and 8). The impaired response did not appear to be related to disease activity since Patients 1 and 8 had clinically inactive disease whereas Patient 2 had active disease.

#### Systemic lupus erythematosus

The *Figure* shows the autologous reactivity of T cells as well as plate-separated T4 and T8 cells in SLE patients and normal controls. T cells from 16 SLE patients showed significantly decreased AMLR reactivity when compared with matched normal T cells (p < 0.01). When T4 (OKT8<sup>-</sup>) and T8 (OKT4<sup>-</sup>) subpopulations were compared, the T4 cells from SLE patients showed poor proliferation compared with normal T4 cells (p < .05). It has been shown that T8 cells will proliferate in an AMLR only in the presence of T4 cells or exogenously supplied IL-2.23 Thus, in these experiments, neither T8 cells from normal controls nor those from SLE patients showed proliferation when cultured alone in an AMLR system.

When the patient group was divided into groups with active and inactive disease, all nine patients with active disease had decreased responses compared with their matched normal counterparts as well as the normal group as a whole (p < 0.01). Of the seven patients with inactive disease, two demonstrated normal AMLR reactivity whereas five patients responded with much lower proliferation. On a statistical basis, therefore, the group with inactive disease did not differ significantly (p > 0.05) from the normals, although the majority of these patients (5/7) did not respond well.

We had the opportunity to study a set of identical twins discordant for the expression of SLE. Cross-stimulation studies were performed on responder and stimulator cells from each member of the pair (*Table 4*). There was a major defect in the patient's T cells, which were unable to respond to her own or the twin's non-T cells. A similar lack of response was seen with the patient's T4 cell population. The SLE non-T cells appeared capable of stimulating the normal twin's T cells. The defect in the T cells also extended to the allogeneic MLR where the patient's cells responded poorly to allogeneic thirdparty non-T cells.

# Discussion

The AMLR, which denotes proliferation of T cells in response to autologous non-T cells, appears to be a complex and important example

			<sup>3</sup> H-thymidine incorpo conta	ration (cpm) in culture		
Subject*	Age (yr)	Activity of disease†	Responders alone	Responders + stimulators	Δ СРМ‡	
Patient 1	35	I	$161 \pm 33$	$241 \pm 160$	80	
Control 1			$1,048 \pm 490$	$6,846 \pm 1,574$	5,798	
Patient 2	15	Α	$438 \pm 60$	$119 \pm 33$	-319	
Control 2			$5,296 \pm 1,048$	$188 \pm 68$	-5,108	
Patient 3	20	. <b>I</b>	$799 \pm 271$	$3,458 \pm 699$	2,659	
Control 3			$5,421 \pm 1,411$	$8,734 \pm 1,309$	3,313	
Patient 4	27	I	$5,020 \pm 1,601$	$12,222 \pm 675$	7,202	
Control 4			$6,552 \pm 2,019$	$15,288 \pm 2,804$	8,736	
Patient 5	18	Α	$1,634 \pm 1,397$	$2,849 \pm 968$	1,215	
Control 5			$2,191 \pm 1,660$	$13,319 \pm 1,600$	11,128	
Patient 6	27	ľ	$4,582 \pm 2,949$	$21,682 \pm 5,008$	17,100	
Control 6			$944 \pm 823$	$3,135 \pm 683$	2,191	
Patient 7	33	I	$358 \pm 86$	$2,516 \pm 622$	2,158	
Patient 8	20	I	$23,400 \pm 6,621$	$22,878 \pm 2,511$	-522	
Control 8			$3,534 \pm 732$	$1,777 \pm 625$	-1,757	•
Patient 9	16	I	$340 \pm 114$	$2,890 \pm 683$	2,550	
Control 9			$58 \pm 1$	$3,678 \pm 974$	3,620	
Patient 10	12	I	$739 \pm 527$	$6,448 \pm 3,510$	5,709	
Control 10			$2,714 \pm 1,469$	$7,050 \pm 1,366$	4,336	
Patient 11	31	I	$224 \pm 47$	$2,694 \pm 1,071$	2,470	
Control 11			$2,467 \pm 627$	$16,344 \pm 3,836$	13,877	
Patient 12	11	I	$837 \pm 71$	$7,152 \pm 2,789$	6,315	
Control 12			$509 \pm 245$	$5,756 \pm 1,408$	5,247	

Table 3.	Autologous mixed lymphocyte reaction in patients with childhood dermatomyositis
	and matched controls

\* Normal controls were age-, sex-, and race-matched with each patient and were tested concurrently.

 $\dagger$  Assessed as described under Materials and Methods. A = active disease; I = inactive disease.

<sup>‡</sup> No significant difference between patients and controls using the Wilcoxon signed rank test for paired data.

of autoreactivity or autorecognition. T cell recognition of cell surface structures encoded by Ia/ DR genes of the major histocompatibility complex is the basis of the AMLR, and such recognition has been shown essential for the afferent limb of normal immune responses to foreign antigens. It has further been shown that immunoregulatory activities of T cells are activated in AMLR cultures. It has therefore been hypothesized that T cell recognition of self, as reflected in appropriate AMLR reactivity, is essential for the maintenance of normal immune homeostasis.<sup>24</sup> AMLR defects have been reported in other autoimmune disorders as well as in neoplastic diseases.<sup>11-15</sup> Autoimmunity-prone strains of mice also demonstrate reduced syngeneic MLR responses.<sup>25</sup> Thus, decreased AMLR may reflect an immunoregulatory imbalance that, together with other factors, may lead to autoimmune disease.<sup>24</sup>

It is likely that cell-mediated autoimmunity to muscle is important in the pathophysiology of PM/DM. The following observations support this hypothesis: (a) lymphocytes sensitized to skeletal muscle are present in patients<sup>26</sup>; (b) patient lymphocytes can mediate injury to muscle cells in



**Figure.** Autologous mixed lymphocyte reactivity (AMLR) activity of T, T4 (OKT8<sup>-</sup>), or T8 (OKT4<sup>-</sup>) cells from normal subjects and individual patients with systemic lupus erythematosus (SLE). Horizontal bars represent the mean.

tissue culture<sup>27</sup>; and (c) lymphocytes from PM/ DM patients can elaborate myotoxic lymphokines when incubated with autologous muscle.<sup>28</sup> Analysis of an autoimmune state such as that seen in PM/DM requires consideration of a diversity of etiologic possibilities including genetic predisposition, viral induction, and immunoregulatory abnormalities.<sup>29</sup>

The impaired AMLR response in our group of patients with PM/DM thus suggests the presence of abnormal immunoregulation in this expression of autoimmunity. It is currently unclear whether this defect precedes disease expression or results from it, through the depletion of AMLR-reactive T cell subsets by their commitment as part of the disease process. We do not

**Table 4.** Autologous mixed lymphocyte reaction in patient with SLE and healthy identical twin

	Proliferative response (Δcpm)		
Stimulator cells	SLE patient	Healthy twin	
T responder cells			
SLE non-T	330	5,392	
Twin non-T	433	3,173	
Third-party non-T	12,887	48,512	
T4 responder cells			
SLE non-T	700	5,906	
Twin non-T	1,210	5,082	

SLE = systemic lupus erythematosus.

think that the decrease in AMLR reactivity could be due to nonspecific illness, as those patients with inactive disease who were clinically well also showed the AMLR defect. Further, in another study, the AMLR has been shown to be unhindered in at least one group of normal individuals who were nonspecifically ill.<sup>30</sup> The cellular basis for this defect is at present unclear. Studies in our laboratory have shown decreased IL-2 production in PM/DM AMLR cultures (unpublished observations). Furthermore, AMLR responses were partially restored following removal of adherent cells (unpublished observations).

In contrast to adult PM/DM, the CDM patients as a group showed no significantly different AMLR responses from normals. CDM differs from adult PM/DM in other respects also. Clinically, there is an increased incidence of calcification of muscle and skin in gastrointestinal vasculitis.<sup>31,32</sup> Conversely, ophthalmologic, pulmonary, and malignant occurrences are less common in CDM than adult PM/DM.<sup>31,33,34</sup> Histologically, most authors agree on the higher incidence of vasculitis affecting small arteries, capillaries, and venules in CDM.<sup>30,35,36</sup> Reticulotubular inclusions within endothelial cells and a reduced capillary/ muscle-fibre ratio have both been demonstrated in CDM,<sup>36</sup> whereas they have not been consistently identified in adult PM/DM.<sup>36</sup> Immunologic differences between CDM and adult PM/DM have also been suggested. Unlike adult PM/DM, lymphocytes from CDM patients do not demonstrate significant cytotoxicity.<sup>37,38</sup> Our finding of an immunoregulatory defect in adult PM/DM but not in CDM thus lends further support to the argument that the pathophysiologic basis for CDM differs from that for adult PM/DM.

SLE is an autoimmune disorder in which multiple immune abnormalities are associated with recurrent or continuous inflammatory processes in a variety of organ systems. It is characterized by the presence of autoantibodies to nuclear and other tissue antigens.<sup>39</sup> Disordered immune system control in SLE has been documented by several investigators.<sup>40</sup> The studies reported in this paper and by others<sup>14,41</sup> indicate that the AMLR is impaired in SLE patients. Furthermore, in a serial study of SLE patients, the AMLR was found to correlate with disease activity; ie, it returned to normal when the disease was quiescent.<sup>42</sup> However, other studies, including this one, have found the AMLR to be uniformly depressed regardless of disease activity.<sup>43</sup> Although as a group our patients with inactive disease were not statistically different from the controls, the finding of a significantly impaired AMLR response in 5/7 of these patients indicates that in most cases the AMLR response does not normalize with a decline in disease activity. The T4 subpopulation is defective since these cells are unable to respond. Whether the T8 cells also have a defect in responding to autologous signals is not known at present since these cells proliferate in an AMLR only in the presence of a source of IL-2.<sup>23</sup> These questions are currently being studied.

Our results with the set of twins discordant for SLE suggest that the AMLR defect resides in the responding T cells and not in the ability of the non-T cells to act as stimulators. Sakane et al<sup>44</sup> came to the same conclusion in a similar study of two sets of identical twins. However, using HLA, A and B, and MLR-matched normal siblings of patients with inactive SLE, Kuntz et al<sup>45</sup> found the defect to be in the non-T stimulators. In the latter study, it is difficult to know whether the patients were matched at the HLA-D locus because the MLR was used both as the criterion of identity and measure of abnormality. Our interpretation of the AMLR defect in SLE is based on a study of a single twin pair. It is entirely possible, given the heterogeneous nature of SLE, that in some patients a stimulatory defect is also present, as suggested by a recent study.<sup>46</sup>

The AMLR T cell defect in SLE appears to represent an intrinsic abnormality in the T cells since it was present regardless of disease activity. SLE is a heterogeneous disease in both its clinical and laboratory manifestations. Thus far, no correlations have been reported in the literature between the degree of AMLR reactivity and clinical and/or laboratory parameters of autoimmunity.

In summary, we have shown, utilizing the AMLR, that defects in immunoregulation exist in adult PM/DM and SLE. Characterization of these defects would aid in the understanding of the pathogenesis of these and other autoimmune disorders.

# Acknowledgments

The authors would like to thank Geri Locker for assistance with statistical analysis and Martina Steele and Gwendolyn Nolan-Jones for excellent secretarial support.

#### References

- Weksler ME, Moody CE, Kozak, RW. The autologous mixed lymphocyte reaction. Adv Immunol 1981; 31:271-312.
- Battisto JR, Ponzio NM. Autologous and syngeneic mixed lymphocyte reactions and their immunological significance. Prog Allergy 1981; 28:160-192.
- Palacios R, Möller G. HLA-DR antigens render resting T cells sensitive to interleukin-2 and induce production of the growth factor in the autologous mixed lymphocyte reaction. Cell Immunol 1981; 63:143-153.
- Gatenby PA, Kotzin BL, Engleman EG. Induction of immunoglobulin secreting cells in the human autologous mixed leukocyte reaction: regulation by helper and suppressor lymphocyte subsets defined with monoclonal antibodies. J Immunol 1981;127:2130-2135.
- Smith JB, Knowlton RP. Activation of suppressor T cells in human autologous mixed lymphocyte culture. J Immunol 1979; 123:419–422.
- Wolos JA, Smith JB. Helper cells in the autologous mixed lymphocyte reaction. I. Generation of cytotoxic T cells recognizing modified self H-2. J Immunol 1981; 127:711-715.
- Goto M, Zvaifler NJ. Characterization of the killer cell generated in the autologous mixed leukocyte reaction. J Exp Med 1983; 157:1309-1323.
- Lattime EC, Gillis S, David C, Stutman O. Interleukin 2, production in the syngeneic mixed lymphocyte reaction. Eur J Immunol 1981; 11:67–69.
- Yu DTY, Chiorazzi N, Kunkel HG. Helper factors derived from autologous mixed lymphocyte cultures. Cell Immunol 1980; 50:305-313.
- Argov S, Cantell K, Klein E, Klein G. Immune interferon (IFN-γ) production in autologous mixed cultures. Cell Immunol 1983; 76:196-199.
- Pope RM, McChesney L, Talal N, Fischbach M. Characterization of the defective autologous mixed lymphocyte response in rheumatoid arthritis. Arth Rheum 1984; 27:1234-1244.
- Engleman EG, Benike CJ, Hoppe RT, Kaplan HS, Berbenich FR. Autologous mixed lymphocyte reaction in patients with Hodgkin's disease. J Clin Invest 1980; 66:149–158.
- James SP, Elson CO, Waggoner JG, Jones EA, Strober W. Deficiency of the autologous mixed lymphocyte reaction in patients with primary biliary cirrhosis. J Clin Invest 1980; 66:1305-1310.
- 14. Sakane T, Steinberg AD, Green I. Failure of autologous mixed lymphocyte reactions between T and non-T cells in patients with systemic lupus erythematosus. Proc Natl Acad Sci (USA) 1978; 75:3464-3468.
- Miyasaka N, Sauvezie B, Pierce D, Daniels TE, Talal N. Decreased autologous mixed lymphocyte reaction in Sjögren's syndrome. J Clin Invest 1980; 66:928-933.
- 16. Mastaglia FL, Ojeda VJ. Inflammatory myopathies: Part 2. Ann Neurol 1985; **17**:317–323.
- 17. Koffler D, Agnello V, Kunkel HG. Polynucleotide immune complexes in serum and glomeruli of patients with systemic lupus erythematosus. Am J Pathol 1974; 74:109-122.
- Bohan A, Peter JB, Bowman RL, Pearson CM. A computerassisted analysis of 153 patients with polymyositis and dermatomyositis. Medicine 1977; 56:255-286.
- Hahn BH, MacDermott RP, Jacobs SB, Pletscher LS, Beale MG. Immunosuppressive effects of low doses of glucocorti-

coids: effects on autologous and allogenic mixed leukocyte reactions. J Immunol 1980; **124**; 2812-2817.

- 20. Tan EM, Cohen AS, Fries JF, et al. The 1982 revised criteria for the classification of systemic lupus erythematosus. Arth Rheum 1982; 25:1271-1277.
- Lokshin MD, Reinitz E, Druzin ML, Murrman M, Estes D. Lupus pregnancy. Case-control prospective study demonstrating absence of lupus exacerbation during or after pregnancy. Am J Med 1984; 77:893-898.
- 22. Hollander M, Wolfe DA. Nonparametric Statistical Methods. New York, Wiley and Sons, 1973.
- Smolen JS, Luger TA, Chused TM, Steinberg AD. Responder cells in the human autologous mixed lymphocyte reaction. J Clin Invest 1981; 68:1601-1604.
- Smith JB, Talal N. Significance of self-recognition and interleukin-2 for immunoregulation, autoimmunity and cancer. Scand J Immunol 1982; 16:269–278.
- Glimcher LH, Steinberg AD, House SB, Green I. The autologous mixed lymphocyte reaction in strains of mice with autoimmune diseases. J Immunol 1980; 125:1832-1838.
- Esiri MM, MacLennan CM, Hazleman BL. Lymphocyte sensitivity to skeletal muscle in patients with polymyositis and other disorders. Clin Exp Immunol 1973; 14:25–35.
- 27. Haas DC, Arnason BGW. Cell-mediated immunity in polymyositis: creatine phosphokinase release from muscle cultures. Arch Neurol 1974; **31**:192–196.
- Johnson RL, Fink CW, Ziff M. Lymphotoxin formation by lymphocytes and muscle in polymyositis. J Clin Invest 1972; 51:2435-2449.
- Allison A. Autoimmune disease: concepts of pathogenesis and control. [In] Talal N, ed. Autoimmunity: Genetic, Immunologic, Virologic and Clinical Aspects. New York, Academic Press, 1977.
- Moody CE, Casazza BA, Christenson WN, Weksler ME. Lymphocyte transformation induced by autologous cells. VIII. Impaired autologous mixed lymphocyte reaction in patients with infectious mononucleosis. J Exp Med 1979; 150:1448-1455.
- 31. Banker BQ, Victor M. Dermatomyositis (systemic angiopathy) of childhood. Medicine 1966; **45:**261-289.
- 32. Crowe WE, Bove KE, Levinson JE, Hilton PK. Clinical and pathogenetic implications of histopathology in childhood polydermatomyositis. Arth Rheum 1982; **25**:126–139.
- Fruman LS, Sullivan DB, Petty RE. Retinopathy in juvenile dermatomyositis. J Pediatr 1976; 88:267-269.
- Park S, Nyhan WL. Fatal pulmonary involvement in dermatomyositis. Am J Dis Child 1975; 129:723-726.
- Banker BQ. Dermatomyositis of childhood: ultrastructural alterations of muscle and intramuscular blood vessels. J Neuropathol Exp Neurol 1975; 34:46–75.

- Carpenter S, Karpati G, Rothman S, Watters G. The childhood type of dermatomyositis. Neurology 1976; 26:952-962.
- Iannaccone ST, Bowen DE, Samaha FJ. Cell-mediated cytotoxicity and childhood dermatomyositis. Arch Neurol 1982; 39:400-402.
- Whitaker JN. Inflammatory myopathy: a review of etiologic and pathogenetic factors. Muscle Nerve 1982; 5:573–592.
- Hughes GRV. Anti-nucleic acid antibodies in SLE—clinical and pathological significance. Clin Rheum Dis 1975; 1:545– 559.
- 40. Steinberg AD, Taurog JD, Cowdery JS, et al. Cellular basis of SLE. [In] Krakauer R, Cathcart M, eds. Immunoregulation and Autoimmunity. New York, Elsevier, 1980.
- 41. Smith JB, DeHoratius RJ. Deficient autologous mixed lymphocyte reactions correlate with disease activity in systemic lupus erythematosus and rheumatoid arthritis. Clin Exp Immunol 1982; 48:155-162.
- 42. Sakane T, Steinberg AD, Green I. Studies of immune functions of patients with systemic lupus erythematosus. V. T cell suppressor function and autologous mixed lymphocyte reaction during active and inactive phases of the disease. Arth Rheum 1980; 23:225-231.
- 43. Sakane T, Kotani H, Takada S, Murakawa Y, Ueda Y. A defect in the suppressor circuits among OKT4+ cell populations in patients with systemic lupus erythematosus occurs independently of a defect in the OKT8+ suppressor T cell function. J Immunol 1983; 131:753-761.
- 44. Sakane T, Steinberg AD, Arnett FC, Reinertsen JL, Green I. Studies of immune functions of patients with systemic lupus erythematosus. III. Characterization of lymphocyte subpopulations responsible for defective autologous mixed lymphocyte reactions. Arth Rheum 1979; 22:770-776.
- Kuntz MK, Innes JB, Weksler ME. The cellular basis of the impaired autologous mixed lymphocyte reaction in patients with systemic lupus erythematosus. J Clin Invest 1979; 63:151-153.
- Crow MK. Enhancement of the impaired autologous mixed leukocyte reaction in patients with systemic lupus erythematosus. J Clin Invest 1985; 76:807-815.

Meher M. Dustoor, Ph.D. Department of Immunology and Cancer Research The Cleveland Clinic Foundation 9500 Euclid Avenue Cleveland, OH 44106