



WILLIAM R. HART, MD, SECTION EDITOR

In vitro comparison of activity of cefixime with activities of other orally administered antimicrobial agents

CYNTHIA C. KNAPP, MS AND JOHN A. WASHINGTON, MD

■ Cefixime, cefuroxime, cephalexin, cefaclor, penicillin, ampicillin, amoxicillin alone and combined with clavulanate, erythromycin, and trimethoprim/sulfamethoxazole were tested in vitro against nearly 1,100 clinical isolates of bacteria. Minimal inhibitory concentrations (MICs) of $> 32 \mu\text{g/mL}$ were observed for all cephalosporins against $> 90\%$ (MIC_{90}) of isolates of *Citrobacter freundii*, *Enterobacter aerogenes*, *Enterobacter cloacae*, *Morganella morganii*, *Acinetobacter anitratus*, and *Pseudomonas* sp. An MIC_{90} of $\leq 0.25 \mu\text{g}$ of cefixime per mL was observed with nonenterococcal streptococci, *Klebsiella* sp, *Proteus mirabilis*, *Proteus vulgaris*, *Providencia stuartii*, *Salmonella* sp, and *Shigella* sp. The MIC_{90} values of cefuroxime and amoxicillin/clavulanate were, respectively, > 32 and $16 \mu\text{g/mL}$ for *Klebsiella oxytoca*, 4 and $32 \mu\text{g/mL}$ for *Klebsiella pneumoniae*, 2 and $1 \mu\text{g/mL}$ for *P mirabilis*, and $> 32 \mu\text{g/mL}$ for *P stuartii*. The MIC_{90} values of cefixime, cefuroxime, amoxicillin/clavulanate, and trimethoprim/sulfamethoxazole were, respectively, 0.12, 2, 0.12, and $\leq 0.5 \mu\text{g/mL}$ for *Branhamella catarrhalis* and 0.25, 4, 8, and $4 \mu\text{g/mL}$ for *Haemophilus influenzae*. Overall, trimethoprim/sulfamethoxazole was the most active compound, followed by cefixime.

□ INDEX TERM: ANTIBIOTICS, LACTAM □ CLEVE CLIN J MED 1988; 55:477-482

CEFIXIME [(6R,7R)-7-[(Z)-2-(2-amino-4-thiazolyl)-2-(carboxymethoxyimino)-acetamido]-8-vinyl-5-thia-1-azabicyclo-[4,2,0]-oct-2-ene-2-carboxylic acid] is a new orally administered cephalosporin that has β -lactamase stability and an in vitro spectrum of activity similar to that of third-generation cephalosporins.¹⁻³

The objective of this study was to compare the activity of cefixime with that of other orally administered antimicrobials including ampicillin, amoxicillin, amoxicillin/clavulanic acid, penicillin, cefaclor, cephalexin, ce-

furoxime (the orally administered axetil ester of which is currently in clinical trials), erythromycin, and trimethoprim/sulfamethoxazole (TMP-SMX).

MATERIALS AND METHODS

Fresh and stock clinical bacterial isolates from cultures of blood, respiratory secretions, urine specimens, and wounds of in- and out-patients seen at The Cleveland Clinic Foundation were selected for testing. Fresh clinical isolates were tested on a consecutive basis between March and July 1987, and stock clinical isolates were used to supplement species represented by few fresh isolates.

Minimal inhibitory concentrations (MICs) were de-

Department of Microbiology, The Cleveland Clinic Foundation.
Submitted for publication Nov 1987; accepted Jan 1988.

This project was supported by a grant from Lederle Laboratories.

TABLE 1
COMPARISON OF IN VITRO ACTIVITY OF CEFIXIME WITH OTHER ANTIMICROBIAL AGENTS*

Organisms	Number tested	Cefixime	Cefuroxime	Cephalexin	Cefaclor	Penicillin	Ampicillin	Amoxicillin	Amoxicillin/Clavulanate†	Erythromycin	Sulfamethoxazole/Trimethoprim‡
<i>Citrobacter diversus</i>	18										
MIC range		≤0.06–0.25	0.5–>32	4–32	0.5–>32	—	2–>32	>32	0.5–>32	—	≤0.5
MIC ₅₀		≤0.06	4	4	4	—	>32	>32	2	—	≤0.5
MIC ₉₀		0.25	16	32	32	—	>32	>32	32	—	≤0.5
<i>Citrobacter freundii</i>	58										
MIC range		≤0.6–>32	0.5–>32	4–32	0.5–>32	—	0.5–>32	0.5–>32	0.5–>32	—	≤0.5–>16
MIC ₅₀		>32	>32	>32	>32	—	>32	>32	>32	—	≤0.5
MIC ₉₀		>32	>32	>32	>32	—	>32	>32	>32	—	>16
<i>Enterobacter aerogenes</i>	65										
MIC range		≤0.06–>32	0.5–>32	16–>32	8–>32	—	8–>32	32–>32	32–>32	—	≤0.5
MIC ₅₀		2	8	>32	>32	—	>32	>32	>32	—	≤0.5
MIC ₉₀		>32	>32	>32	>32	—	>32	>32	>32	—	≤0.5
<i>Enterobacter cloacae</i>	103										
MIC range		≤0.06–>32	1–>32	4–>32	16–>32	—	4–32	8–>32	8–>32	—	≤0.5–>16
MIC ₅₀		4	16	>32	>32	—	>32	>32	>32	—	≤0.5
MIC ₉₀		>32	>32	>32	>32	—	>32	>32	>32	—	≤0.5
<i>Escherichia coli</i>	136										
MIC range		≤0.06–>32	0.25–>32	2–>32	0.25–>32	—	0.25–>32	0.12–>32	≤0.06–>32	—	≤0.5–>16
MIC ₅₀		0.25	2	8	2	—	4	4	4	—	≤0.5
MIC ₉₀		32	16	>32	>32	—	>32	>32	>32	—	≤0.5
<i>Klebsiella oxytoca</i>	30										
MIC range		≤0.06–8	0.25–>32	2–>32	0.25–>32	—	16–>32	16–>32	0.5–>32	—	≤0.5
MIC ₅₀		≤0.06	1	4	0.5	—	>32	>32	2	—	≤0.5
MIC ₉₀		≤0.06	8	32	16	—	>32	>32	32	—	≤0.5
<i>Klebsiella pneumoniae</i>	88										
MIC range		≤0.06–1	0.5–32	0.5–>32	0.25–32	—	8–>32	2–>32	0.5–>32	—	≤0.5–>16
MIC ₅₀		≤0.06	2	4	0.5	—	32	>32	2	—	≤0.5
MIC ₉₀		≤0.6	4	8	2	—	>32	>32	32	—	≤0.5
<i>Morganella morganii</i>	41										
MIC range		≤0.06–>32	4–>32	32–>32	>32–>32	—	16–>32	32–>32	32–>32	—	≤0.5–>16
MIC ₅₀		2	>32	>32	>32	—	>32	>32	>32	—	≤0.5
MIC ₉₀		>32	>32	>32	>32	—	>32	>32	>32	—	≤0.5
<i>Proteus mirabilis</i>	62										
MIC range		≤0.06–8	0.25–16	4–>32	0.5–>32	—	0.5–>32	0.25–>32	0.25–>32	—	≤0.5–>16
MIC ₅₀		≤0.06	1	8	1	—	1	1	0.5	—	≤0.5
MIC ₉₀		≤0.06	2	16	2	—	2	1	1	—	≤0.5
<i>Proteus vulgaris</i>	21										
MIC range		≤0.06–>32	8–>32	16–>32	16–>32	—	32–>32	>32	1–>32	—	≤0.5
MIC ₅₀		≤0.06	>32	>32	>32	—	>32	>32	8	—	≤0.5
MIC ₉₀		0.12	>32	>32	>32	—	>32	>32	>32	—	≤0.5

TABLE 1 – continued
COMPARISON OF IN VITRO ACTIVITY OF CEFIXIME WITH OTHER ANTIMICROBIAL AGENTS*

Organisms	Number tested	Cefixime	Cefuroxime	Cephalexin	Cefaclor	Penicillin	Ampicillin	Amoxicillin	Amoxicillin/Clavulanate†	Erythromycin	Sulfamethoxazole/Trimethoprim‡
<i>Providencia rettgeri</i>	17										
MIC range		≤0.06–2	≤0.06–>32	2–>32	0.25–>32	—	0.5–>32	2–>32	4–>32	—	≤0.5–>16
MIC ₅₀		≤0.06	0.25	>32	>32	—	32	>32	32	—	1
MIC ₉₀		0.5	>32	>32	>32	—	>32	>32	>32	—	>16
<i>Providencia stuartii</i>	14										
MIC range		≤0.06–0.5	≤0.06–16	4–>32	0.5–>32	—	0.5–>32	4–>32	1–32	—	≤0.5–>16
MIC ₅₀		≤0.06	0.5	>32	>32	—	>32	>32	32	—	>16
MIC ₉₀		0.12	4	>32	>32	—	>32	>32	>32	—	>16
<i>Serratia marcescens</i>	58										
MIC range		≤0.06–>32	8–>32	32–>32	16–>32	—	2–>32	2–>32	4–>32	—	≤0.5–>16
MIC ₅₀		0.5	>32	>32	>32	—	>32	>32	>32	—	≤0.5
MIC ₉₀		16	>32	>32	>32	—	>32	>32	>32	—	1
<i>Streptococcus</i>	50										
Groups A,B,C,F,G											
MIC range		≤0.06–0.25	≤0.06	<0.06–4	≤0.06–0.5	≤0.06	≤0.06	≤0.06	≤0.06	≤0.06–8	≤0.5
MIC ₅₀		≤0.06	≤0.06	0.25	≤0.06	≤0.06	≤0.06	≤0.06	≤0.06	≤0.06	≤0.5
MIC ₉₀		0.25	≤0.06	2	0.5	≤0.06	≤0.06	≤0.06	≤0.06	≤0.06	≤0.5
<i>Streptococcus pneumoniae</i>	12										
MIC range		≤0.06–0.5	≤0.06–0.25	≤0.6–8	≤0.06–0.5	≤0.06	≤0.06	≤0.06	≤0.06	≤0.06–0.25	≤0.5–2
MIC ₅₀		≤0.06	≤0.06	≤0.06	≤0.06	≤0.06	≤0.06	≤0.06	≤0.06	≤0.06	≤0.5
MIC ₉₀		2	≤0.06	2	0.5	≤0.06	≤0.06	≤0.06	≤0.06	≤0.06	≤0.5
<i>Branhamella catarrhalis</i>	17										
MIC range		≤0.06–0.25	0.12–2	0.12–4	0.12–4	≤0.06–16	≤0.06–4	≤0.06–4	≤0.06–0.12	0.12–5	≤0.5
MIC ₅₀		≤0.06	1	2	1	4	1	1	≤0.06	0.12	≤0.5
MIC ₉₀		0.25	2	4	2	16	4	4	0.12	0.5	≤0.5
<i>Haemophilus influenzae</i>	23										
MIC range		≤0.06–4	≤0.06–16	2–>32	0.5–>32	—	0.06–>32	0.12–>32	0.12–8	0.5–8	≤0.5–4
MIC ₅₀		≤0.06	0.5	8	2	—	8	8	0.5	4	≤0.5
MIC ₉₀		0.25	4	>32	16	—	32	32	8	8	4
<i>Staphylococcus aureus</i>	10										
MIC range		8–32	1–2	2–8	2–32	1–>16	2–>32	1–>32	0.5–4	≤0.06–0.12	≤0.5
MIC ₅₀		16	2	4	4	4	4	4	1	0.12	≤0.5
MIC ₉₀		16	2	4	8	>16	32	16	2	0.12	≤0.5
<i>Staphylococcus</i>	27										
coagulase-neg.											
MIC range		2–>32	0.12–>32	1–>32	1–>32	≤0.06–>16	≤0.06–>32	0.06–>32	0.06–32	≤0.06–>8	0.5–>16
MIC ₅₀		16	0.5	16	8	4	8	8	1	>8	2
MIC ₉₀		>32	>32	>32	>32	>16	>100	>32	>32	>8	>16

* MIC = Minimal inhibitory concentration in µg/mL; MIC₅₀ = MIC for 50% of isolates; MIC₉₀ = MIC for 90% of isolates.

† MIC listed expresses amoxicillin activity (combined on a 2:1 basis with clavulanate).

‡ MIC listed expresses trimethoprim activity (combined on a 1:20 basis with sulfamethoxazole).

TABLE 2
ACTIVITY OF CEFIXIME AGAINST SPECIES REPRESENTED BY
<10 ISOLATES

Organism (no.)	MIC (μg/mL) for individual isolates
<i>Acinetobacter lwoffii</i> (5)	1, 2, 4(2), 32
<i>Alcaligenes odorans</i> (2)	2, 4
<i>Alcaligenes xylosoxidans</i> (1)	>32
<i>Cedecea</i> sp (1)	0.5
<i>Cedecea neteri</i> (4)	0.125, 0.25, 0.5 (2)
<i>Citrobacter amalonaticus</i> (1)	≤0.06
<i>Enterobacter agglomerans</i> (3)	0.25, 1
<i>Haemophilus parainfluenzae</i> (1)	≤0.06
<i>Hafnia alvei</i> (1)	>32
<i>Providencia alcalifaciens</i> (1)	≤0.06
<i>Salmonella</i> sp (9)	≤0.06(6), 0.125(3)
<i>Serratia liquefaciens</i> (8)	≤0.06(2), 0.125(3), 0.25(2), >32
<i>Serratia odorifera</i> (1)	≤0.06
<i>Serratia rubidaea</i> (1)	0.125
<i>Shigella dysenteriae</i> (1)	0.125

terminated according to the microdilution methods and controls recommended by the National Committee for Clinical Laboratory Standards in which approximately 5×10^5 colony forming units (CFU)/mL were inoculated into a \log_2 progression of concentrations of each antimicrobial that was incorporated in cation-supplemented Mueller-Hinton broth.⁴ Studies with 10^7 CFU of seven isolates of β -lactamase-positive isolates of *H influenzae* and 10 β -lactamase-positive isolates of *Staphylococcus aureus* per mL were performed to determine inoculum effects on MICs of cefixime and cefuroxime. The ranges of concentrations of antimicrobials tested were as follows: cephalosporins, 0.06–32 μg/mL; ampicillin and amoxicillin alone and when combined in a 2:1 ratio with clavulanate, 0.06–32 μg/mL; penicillin, 0.06–16 μg/mL; erythromycin, 0.06–8 μg/mL; and trimethoprim, 0.5–16 μg/mL combined in a 1:20 ratio with sulfamethoxazole.

RESULTS

A total of 1,096 freshly collected and stocked clinical isolates was tested. Listed in Table 1 are the ranges of MICs and the MICs for 50 and 90% of isolates for species represented by at least 10 isolates. Listed in Table 2 are the MICs of cefixime for species represented by fewer than 10 isolates. Among cephalosporins tested, cefixime, followed by cefuroxime, was the most active against the Enterobacteriaceae, although the MIC₉₀ values of both cephalosporins against isolates of *Citrobacter freundii*, *Enterobacter* sp, and *Morganella morganii* were > 32 μg/mL. Cefixime was generally inactive against *Acineto-*

bacter anitratus, *Pseudomonas aeruginosa*, and *Pseudomonas maltophilia* (data not shown). The MIC ranges and MIC₉₀ values of amoxicillin/clavulanate closely paralleled those of cefixime and cefuroxime against the Enterobacteriaceae, *B catarrhalis*, and *H influenzae*; however, the MIC₅₀ values of cefixime were substantially lower (generally 3 to 5 \log_2 dilutions) than those of cefuroxime and amoxicillin/clavulanate against the Enterobacteriaceae. The activities of amoxicillin/clavulanate and cefixime against *B catarrhalis* and *H influenzae* were virtually identical. The most active compound overall against the Enterobacteriaceae was the combination of TMP-SMX. TMP-SMX was also active against *B catarrhalis*.

Inoculum effects on the activities of cefixime and cefuroxime were tested with inocula of 5×10^5 CFU/mL and 5×10^7 CFU/mL of 7- β -lactamase-producing *H influenzae*. MICs of each antimicrobial agent remained unchanged in five instances and increased ≥ 5 -fold in two instances. No inoculum effects on the activities of cefixime and cefuroxime were noted in studies of 5×10^5 CFU/mL and 5×10^7 CFU/mL of 10- β -lactamase-producing *S aureus*.

All antimicrobials tested were active against nonenterococcal streptococci. Most staphylococci tested were penicillin-resistant (MIC > 0.12 g/mL) and, therefore, resistant to ampicillin and amoxicillin. Cefuroxime, cephalixin, amoxicillin/clavulanate, erythromycin, and TMP-SMX were the most active antimicrobials against *S aureus*. None of the antimicrobials tested were consistently active against coagulase-negative staphylococci. Cefixime was generally inactive against staphylococci.

DISCUSSION

The α -methoxyimino aminothiazole side chain at the 7-position of the cephem nucleus is shared by cefixime, cefotaxime, ceftizoxime, and cefmenoxime; it provides enhanced stability to plasmid and chromosomal β -lactamases and enhanced affinity for target enzymes (penicillin-binding proteins) relative to first-generation cephalosporins.¹⁻³ Thus, cefixime has a substantially broader spectrum of activity against gram-negative bacteria, including *H influenzae* and *B catarrhalis*, than do currently available orally absorbed cephalosporins, such as cephalixin and cefaclor. Although cefixime is generally more active than cefuroxime and ampicillin/clavulanate on a weight for weight basis, the spectrum of activity of these three antimicrobial agents is very similar. This similarity exists because neither cefixime nor cefuroxime

is hydrolyzed by plasmid-mediated β -lactamases and because clavulanic acid inactivates plasmid-mediated β -lactamases.^{2,3}

Conversely, since clavulanic acid does not inactivate the type I chromosomally mediated β -lactamase, which is most notably produced by *C freundii*, *Enterobacter* sp, *Serratia marcescens*, *M morganii*, and *P aeruginosa*, and since mutants of these species that are derepressed for the type I β -lactamase slowly hydrolyze second- and third-generation cephalosporins,⁵ it is not surprising that amoxicillin/clavulanate, cefuroxime, and cefixime exhibited poor activity against our isolates of *C freundii*, *Enterobacter* sp, *M morganii*, and *S marcescens*. The MIC₉₀ values for all three antimicrobials against these species were, with a single exception, $> 32 \mu\text{g/mL}$. The frequency of resistance of our isolates of *C freundii*, *Enterobacter* sp, *M morganii*, and *S marcescens*, as well as of Proteaceae other than *P mirabilis*, to ampicillin/clavulanate and cefuroxime is in accord with that reported by others,^{2,6,7} while that to cefixime is in accord with that reported by Neu et al² and Utsui et al⁷ but is higher than that reported by Fuchs et al,³ Mulligan and Kwok,⁸ and Tanaka et al.⁶ As reported by others,^{1,3,7} cefixime, cefuroxime, and amoxicillin/clavulanate were active against β -lactamase-producing isolates of *H influenzae* and *B catarrhalis*. With the exceptions of *Klebsiella pneumoniae* and *Proteus mirabilis*, other gram-negative bacilli were generally resistant to cephalexin and cefaclor, and gram-negative bacilli other than *P mirabilis* were largely resistant to amoxicillin and ampicillin. Overall, the most active antimicrobial agent tested against gram-negative bacteria was TMP-SMX, resistance to which was limited to *Providencia* sp and *H influenzae*.

Although active against nonenterococcal streptococci, cefixime had, as also reported by others,^{1-3,7} little activity against staphylococci. All strains of *S aureus* tested were penicillinase producers and were, therefore, resistant to the penicillins but susceptible to amoxicillin/clavulanate, cephalosporins other than cefixime, TMP-SMX, and erythromycin. None of the antimicrobials tested were particularly active against coagulase-negative staphylococci.

Peak serum levels of cefixime have been reported to be in the range of $3.85 \pm 0.23 \mu\text{g/mL}$ to $4.92 \pm 0.51 \mu\text{g/mL}$ following an oral dose of 400 mg in healthy subjects.^{9,10} Thus, it has been proposed that bacteria inhibited by $\leq 1 \mu\text{g/mL}$ of cefixime per mL be considered susceptible and that those with MICs of $\geq 4 \mu\text{g/mL}$ be considered resistant.³ The corresponding MIC equivalents of susceptibility and resistance for all cephalosporins other than cefoperazone are $\leq 8 \mu\text{g/mL}$ and $\geq 32 \mu\text{g/mL}$, respectively⁴;

however, since peak levels of cefuroxime average $8.6 \mu\text{g/mL}$ following a 500-mg oral dose of the axetil ester,¹¹ the MIC equivalents of susceptibility and resistance for this compound will likely be reduced somewhat.

Because of the considerable variation in peak cefuroxime levels ($4.5 \mu\text{g/mL}$ to $12.8 \mu\text{g/mL}$) following a 500-mg oral dose of the axetil ester¹¹ and the high relapse rate of women treated with cefuroxime axetil for recurrent bacteriuria,¹² it may be advisable to set the MIC equivalents for susceptibility and resistance of the axetil ester at the same concentrations as those proposed for cefixime. If this were to be the case, few gram-negative bacteria tested in our laboratory would be considered susceptible to cefuroxime axetil, and the oral antimicrobials of choice in descending order of activity would be TMP-SMX, cefixime, and amoxicillin-clavulanate. Ultimately, however, the role of cefixime in the treatment of urinary tract infections must be assessed in clinical trials similar to those carried out with cefuroxime axetil by Brumfitt et al¹² and must be compared with that of TMP-SMX, which exhibits broader activity in vitro and has a well-established record in the treatment of bacteriuria.

In conclusion, cefixime is a novel third-generation oral cephalosporin with activity against nonenterococcal streptococci and plasmid-mediated β -lactamase-producing gram-negative bacteria but with poor activity against staphylococci and chromosome-mediated β -lactamase-producing gram-negative bacteria. Cefixime is more active than other oral cephalosporins against β -lactamase-producing strains of *H influenzae* and *B catarrhalis*, the incidence of which has increased in acute otitis media and in acute exacerbations of chronic bronchitis. Whether, however, cefixime will have any advantages over amoxicillin/clavulanate in the treatment of these types of respiratory tract infections remains to be seen. Finally, cefixime would appear to have little application in skin and soft-tissue infections because of its poor antistaphylococcal activity.

ACKNOWLEDGMENTS

Isolates of *Branhamella catarrhalis*, *Haemophilus influenzae*, and *Streptococcus pneumoniae* included strains generously provided by Michael Jacobs, MD, PhD, University Hospitals of Cleveland (OH) and Clyde Thornsberry, PhD, Centers for Disease Control, Atlanta, GA.

Cefixime was supplied by Lederle Laboratories; cefuroxime by Glaxo Inc.; penicillin, cephalexin, cefaclor, and erythromycin by Lilly Research Laboratories; amox-

icillin and amoxicillin/clavulanate by Beecham Laboratories; TMP-SMX by Burroughs Wellcome Co.; and ampicillin by Bristol Laboratories.

The authors also gratefully acknowledge Lynne Atkinson for technical assistance in preparing microdilution trays and Estella Stovall for assistance in preparing the manuscript.

JOHN A. WASHINGTON, MD
Department of Microbiology
The Cleveland Clinic Foundation
One Clinic Center
9500 Euclid Avenue
Cleveland, Ohio 44195

REFERENCES

1. Kaminura T, Kojo H, Matsumoto Y, Mine Y, Goto S, Kuwahara S. In vitro and in vivo antibacterial properties of FK 027, a new orally active cephem antibiotic. *Antimicrob Agents Chemother* 1984; 25:98-104.
2. Neu HC, Labthavikul P. Comparative in vitro activity and β -lactamase stability of FR 17027, a new orally active cephalosporin. *Antimicrob Agents Chemother* 1984; 26:174-180.
3. Fuchs PC, Jones RN, Barry AL, et al. In vitro evaluation of cefixime (FK 027, FR 17027, CL 28465): Spectrum against recent clinical isolates, comparative antimicrobial activity, β -lactamase stability, and preliminary susceptibility testing criteria. *Diagn Microbiol Infect Dis* 1986; 5:151-162.
4. National Committee for Clinical Laboratory Standards. Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically; Approved Standard. NCCLS publication M7-A. Villanova, PA, NCCLS, 1985.
5. Nayler JHC. Resistance to β -lactams in gram-negative bacteria: relative contributions of β -lactamase and permeability limitations. *J Antimicrob Chemother* 1987; 19:713-732.
6. Tanaka SK, Schwind Summerill RA, Minassian BF, et al. In vitro evaluation of tigemonam, a novel oral monobactam. *Antimicrob Agents Chemother* 1987; 31:219-225.
7. Utsui Y, Matsuhisa I, Mitsuhashi S. In vitro and in vivo antibacterial activities of CS-807, a new oral cephalosporin. *Antimicrob Agents Chemother* 1987; 31:1085-1092.
8. Mulligan ME, Kwok YY. In vitro activity of cefixime and six other agents against nosocomial pathogens of the Enterobacteriaceae family. *Infect Control* 1987; 8:241-244.
9. Guay DRP, Meatherall RC, Harding GK, Brown GR. Pharmacokinetics of cefixime (CL 284,635;FK 027) in healthy subjects and patients with renal insufficiency. *Antimicrob Agents Chemother* 1986; 30:485-490.
10. Brittain DC, Scully BE, Hirose T, Neu HC. The pharmacokinetic and bactericidal characteristics of oral cefixime. *Clin Pharmacol Ther* 1985; 38:590-594.
11. Sommers DK, Van Wyk M, Williams PEO, Harding SM. Pharmacokinetics and tolerance of cefuroxime axetil in volunteers during repeated dosing. *Antimicrob Agents Chemother* 1984; 25:344-347.
12. Brumfitt W, Hamilton-Miller JMT, Smith GW. Comparative trial of cefuroxime axetil in recurrent urinary tract infections illustrating importance of 6-week follow-up. *Antimicrob Agents Chemother* 1987; 31:1442-1443.