#### CLINICAL DECISION-MAKING GARY S. HOFFMAN, MD, EDITOR



KENNETH E. SACK, MD Professor of Clinical Medicine, Director of Clinical Programs in Rheumatology, University of California San Francisco

# The difficulties of differentiating vasculitis from its mimics

# ABSTRACT

**Even biopsy** 

misleading

findings can be

The signs and symptoms of vasculitis are not specific, and tests for confirming the diagnosis can be misleading. Thus, when considering a diagnosis of vasculitis, physicians need to keep an open mind. With a case vignette, the author illustrates some of the difficulties in diagnosing "vasculitis."

NFLAMMATION IS ONLY one of the many causes of vascular injury. For this reason, and because the treatment of vasculitis often involves the use of potentially toxic agents, the clinician should keep an open mind when evaluating patients with organ ischemia.

## IS IT VASCULITIS— OR SOME OTHER VESSEL INJURY?

Numerous conditions can injure or occlude blood vessels and mimic the clinical picture of vasculitis.<sup>1</sup> In fact, the clinical manifestations of vascular injury depend more on the size and location of the affected vessel than on the cause of the injury.

For example, damage to small vessels can cause a variety of cutaneous changes, including palpable purpura, urticaria, livedo reticularis, papulovesicular lesions, and nodules. Likewise, such involvement of small vessels can lead to dysfunction of the heart, kidney, gastrointestinal tract, and nerves. When medium-sized arteries are affected, cutaneous nodules or ulcerations, peripheral gangrene, or organ infarction can evolve. Disease of large vessels can result in claudication or major organ dysfunction.

# Tests used to diagnose vasculitis lack specificity

Lack of specificity of angiographic, biopsy, and

laboratory findings may create further difficulty in diagnosing vasculitis.

Angiography. Irregularities of vessel walls, segmental arterial occlusions, and vascular dilatations—angiographic findings considered characteristic of vasculitis—may appear in a variety of nonvasculitic conditions (TABLE 1).

**Biopsy findings.** The histopathology of vascular tissues can also be misleading. For example, fibrinoid changes in a vessel wall may result from acute arterial hypertension rather than from vasculitis. Vascular hypertrophy is common in the lower leg of normal individuals, as is cellular infiltration of perivascular tissues.<sup>2</sup> And vascular inflammation can result not only from immunologic events but also from such processes as infection,<sup>3</sup> embolic phenomena,<sup>4</sup> and cold-induced injury.<sup>5</sup>

Laboratory tests may likewise lead the clinician astray. The appearance of an anti-DNA antibody or an antineutrophilic cytoplasmic antibody (ANCA), which are deemed fairly specific for systemic lupus erythematosus (SLE) and Wegener's granulomatosis, respectively, may actually reflect an infectious or neoplastic process.<sup>6,7</sup>

**Clinical findings.** Finally, cutaneous lesions of a nonvascular nature (eg, neutrophilic dermatoses,<sup>8</sup> pyoderma gangrenosum,<sup>9</sup> erythema nodosum,<sup>10</sup> insect bites<sup>11</sup>) may be mistaken for those caused by vasculitis.

The following vignette illustrates some of the difficulties in diagnosing "vasculitis."

#### CASE PRESENTATION

A 69-year-old woman presented with worsening hypertension, a cyanotic toe, and renal insufficiency. Two years previously, biopsy of a skin rash had shown "leukocytoclastic angiitis." At that time, laboratory studies disclosed a mild anemia, an erythrocyte sedimentation rate (ESR) of 85 mm/hour, a serum creatinine value of 2.2 mg/dL, a urine protein concentration of 30 mg/dL, and a negative test for antinuclear antibodies (ANA). The patient declined to undergo renal biopsy. She received prednisone in doses as high as 60 mg per day, but ischemic necrosis of the left hip necessitated stopping this treatment.

Four months before admission, the patient developed cyanosis of her left third toe. She also complained of fatigue, weakness, depression, and mild weight loss. The finding of severe stenoses of the carotid arteries led to bilateral carotid endarterectomies. Over the ensuing months, her hypertension and renal failure worsened.

At admission to the hospital, the patient's blood pressure was 200/126 mm Hg. She had several small purpuric lesions on her extremities and a faint erythematous eruption on her back. The left third toe was cyanotic, and pulses in the left leg were diminished. Cardiac examination was unremarkable, and there were no abdominal bruits. A radiograph showed aneurysmal dilatation of the thoracic aorta.

Laboratory studies were as follows:

- Hematocrit 24.6%
- White blood cell count  $12.8 \times 10^{9}/L$
- ESR 72 mm/hour
- Urine protein 1+ with a normal urine sediment
- Serum creatinine concentration 3 mg/dL
- Serum cholesterol concentration 228 mg/dL
- Serum triglyceride concentration 192 mg/dL
- Antinuclear antibodies positive at a 1:80 dilution (homogeneous pattern)
- Anti-DNA antibody present in a low titer
- Serum cryoglobulins were absent.

Initial diagnostic considerations included SLE and polyarteritis nodosa. However, the combination of progressive renal failure, ischemia of a digit, and asymmetric pulses in a patient with known atherosclerosis made atheromatous embolic disease a more likely diagnosis.

The patient was ultimately discharged

# TABLE 1

# Angiographic mimics of vasculitis

Amyloidosis

Atrial myxoma

Drug abuse (eg, ergot derivatives and sympathomimetic agents)

Ehlers-Danlos syndrome

Exposure to cold or radiation

Fibromuscular dysplasia

Infection (eg, bacterial, fungal, rickettsial, spirochetal, viral)

Injection of contrast material

Migraine

Moyamoya disease

Neoplasm (eg, pheochromocytoma, vascular lymphoma, vascular encasement by solid tumor)

Neurofibromatosis

Pseudoxanthoma elasticum

Systemic hypertension

Thrombotic thrombocytopenic purpura

#### Trauma

SOURCE: ADAPTED FROM SACK KE. MIMICKERS OF VASCULITIS, IN: KOOPMAN WJ, EDITOR. ARTHRITIS AND ALLIED CONDITIONS, 13TH EDITION. BALTIMORE: WILLIAMS AND WILKINS, 1997: 1525–1546.

and died at home. At autopsy, histologic examination of the kidneys showed intravascular clefts typical of cholesterol deposits in small renal arteries. There was no evidence of vasculitis.

## COMMENT: THE TREACHEROUS DIAGNOSTIC PATH

This case illustrates some of the ways atheromatous emboli can produce a picture of "vasculitis." Skin lesions accompanied by digital cyanosis, accelerated hypertension, and renal insufficiency certainly bring to mind polyarteritis nodosa. And positive tests for ANA and anti-DNA antibodies as well as elevated ESRs also suggest connective tissue disease. Indeed, many patients with atheroembolic disease have a low Anti-DNA and ANCA may reflect infection or neoplasms We thank those who reviewed manuscripts submitted to the *Cleveland Clinic Journal of Medicine* for the year ending September 30, 1998. Reviewing papers for scientific journals is an arduous task and involves considerable time and effort. We are grateful to these reviewers for contributing their expertise this past year.—John D. Clough, MD, Editor-in-Chief.

Achkar, Edgar Alfes, John Andrish, Jack T Bartholomew, John R Belinson, Jerome L Braun, William E Bronson, David Brouhard, Ben Burt, Randall W Cain, Robert A Carey, William D Castle, Lon Chilcote, William A Clough, Mary Cochran, Bertram H Crowe, Joseph Culver, James E Jr Cunningham, Robert D'Amico, Joseph A Daugherty, Silas Deal, Chad L Dinner, Dudley S Domen, Ronald E Donley, Brian G Durbeck, Donald C Farmer, Richard G Fessler, Barri J Foley, Kevin T Francis, Garv Frazee, Louise E Gidwani, Gita P Gifford, Ray W Jr Gonsalves-Ebrahim, Lilian V Good, Milton B Gorbien, Martin Gordon, Steven M Gorensek, Margaret Grant, R Peery Graves, Thomas M Gray, Lawrence Groene, Linda Gupta, Manjula K Hall, Geraldine S Hall, James O Hall, Phillip M Harris, Michael A Hart, William R Hartwell, Shattuck W Jr

Hayden, Stephen P Henry, Catherine Hobbs, Robert E Hoeltge, Gerald A Hoffman, Garv Holland, Jimmie CB Hoogwerf, Byron J James, Karen Kavuru, Mani Keys, Thomas F Khoury, Alan Kothari, Shakuntala Kratche, Richard P Kunkel, Robert Lakin, Milton M Lamb, James F Lang, Richard Lashner, Bret Lauer, Michael S Lederman, Richard J Levin, Kerry Licata, Angelo A Lichtin, Alan Lipton, Mark A Lowenthal, Gilbert Mandell, Brian Mayer, Mark E Mazanec, Daniel J McCollough, Arthur J McKibben, Jeanne Mehta, Neil Michota, Franklin Miller, Donna Moodie, Douglas S Morledge, Thomas J Murray, Richard Nelson, John Newman, Georgia Nickerson, Paul E Nissen, Steven Norr, Sigmund O'Toole, Elizabeth Ockner, Stephen Olin, Jeffrey Owens, Francis Sr Palmer, Robert M Potts, Jeanette Radwany, Steven

Ransohoff, David Ratliff, Norman B Reddy, Sethu Rehm, Susan J Ricketts, Gregory A Rollins, Michael B Rome, Ellen Rose, Leslie I Rosian, Rochelle Ross, Elisa K Roth, Mark Ruschhaupt, William F III Sahgal, Vinod Schuler, Michael A Schumacher, O Peter Shafer, William A Shane, Elizabeth Shapiro, Howard D Sheahan, Michael G Singsen, Bernhard H Smedira, Holly Smith, Martin L Soffer, Edy E Solomon, Glen D Somani, Peter Spiess, Patricia Staley, Ronna L Stoller, James K Sweeney, Patrick J Taylor, Harris C Tchou, Patrick J Thacker, Holly L Thomas, George Valera, Manuel Vann, Ana R Vidt, Donald G Wagner, William Walborn, Mary A Washington, John A II Weinhous, Martin S Weinstein, Cheryl E Wexner, Steven Whitcomb, Winthrop Wilke, William S Williams, Marc S Young, James B

SACK

serum complement level,<sup>12</sup> a finding typical of active SLE. This case also illustrates the nonspecificity of vascular inflammation in a biopsy specimen. Although this patient's rash (deemed leukocytoclastic vasculitis) could have reflected another disease process, similar rashes can occur in atheroembolic disease.<sup>13</sup>

This case illustrates how the pathway to the diagnosis of vasculitis can indeed be treacherous; and seldom in medicine is the admonition *primum non nocere*—first, do no harm—more applicable.

#### REFERENCES

- Sack KE. Mimickers of vasculitis, In: Koopman WJ, editor. Arthritis and Allied Conditions, 13th edition. Baltimore: Williams and Wilkins, 1997:1525–1546.
- Ryan TJ, Wilkinson DS. Cutaneous vasculitis; "Angiitis." In: Rook A, Wilkinson DS, Ebling FJG, Champion RH, Burton JL, editors. Textbook of Dermatology, 4th edition. Oxford: Blackwell Scientific, 1986:1121–1185.
- Somer T, Finegold SM. Vasculitis associated with infections, immunization, and antimicrobial drugs. Clin Infect Dis 1995; 20:1010–1036.
- 4. Retan JW, Miller RE. Microembolic complications of atherosclerosis. Literature review and report of a patient. Arch Intern Med 1966; 118:534–545.
- 5. Wall LM, Smith NP. Perniosis: A histopathological review. Clin Exper Dematol 1981; 6:263–271
- 6. Edgars JDM. The clinical utility of ANCA positivity. Ann Rheum Dis 1996; 55:494–496.
- Zuckerman KK, Leventhal L, Wynne C. Positive c-ANCA in a patient with lymphoma and without vasculitis. J Clin Rheumatol 1997; 3:279–281.
- Moreland LW, Brick JE, Kovack RE. Acute febrile neutrophilic dermatitis (Sweet syndrome): A review of the literature with emphasis on musculoskeletal manifestations. Semin Arthritis Rheum 1988; 17:143–155.
- 9. Callen JP. Pyoderma gangrenosum and related disorders. Med Clin North Am 1989; 73:1247–1261.
- Prestes C, Winklemann R, Su W. Septal granulomatous panniculitis: Comparison of the pathology of erythema nodosum migrans (migratory panniculitis) and chronic erythema nodosum. J Am Acad Dermatol 1990; 22:477–483.
- Rees RS, Fields JP, King LE. Do brown recluse spider bites induce pyoderma gangrenosum? South Med J 1985; 78:283–288.
- Cosio FG, Zager RA, Sharma HM. Atheroembolic renal disease causes hypocomplementemia. Lancet 1985; 1:118–121.
- Falanga V, Fine MJ, Kapoor WN. The cutaneous manifestations of cholesterol crystal embolization. Arch Dermatol 1986; 122:1194–1198.

ADDRESS: Kenneth E. Sack, MD, University of California San Francisco, 400 Parnassus Avenue, Room A-587, San Francisco, CA 94143-0326.