.....

Is God good for your health? The role of spirituality in medical care

DAVID B. LARSON, MD, MSPH

President, National Institute for Healthcare Research The National Institute for Healthcare Research, soon to be renamed the Institute for Spirituality and Health, is a private educational and research organization committed to exploring the relationship between spirituality and health in medicine and the social sciences.

HAROLD G. KOENIG, MD

Associate Professor of Psychiatry and Medicine, Duke University Medical Center

ABSTRACT

Many studies have found that religious belief and practice have a positive effect on physical and mental health, although the topic needs more research. As religious beliefs may affect both health and healthpromoting behavior, physicians should try to understand their patients' beliefs.

Studies say religious people are healthier

R ELIGION IS OFTEN considered antithetical to modern medicine, and many physicians still consider religion an inappropriate topic for a medical conversation. Yet a growing body of research suggests that religious commitment is associated with mental and physical health.

More than 40% of Americans attend worship services weekly, almost three-quarters say religious faith forms the basis for their approach to life, and 95% believe in God.¹ People are even more likely to consider their faith important when they have a severe illness. In one study, a majority of patients agreed or strongly agreed that they would like their physicians to ask whether they have spiritual or religious beliefs that would influence their medical decisions if they became gravely ill.²

At the very least, the gap between patients' and physicians' approach to spiritual issues makes many patients dissatisfied with the medical profession. In one survey, most hospital patients would have liked their physician to include spiritual issues when discussing medical care, but only about 10% of physicians did so.³

At worst, physicians' historical inattention to religious and spiritual issues may be causing them to overlook factors that play an important role in their patients' health.

RELIGION MAY PROMOTE HEALTH

Nearly every medical study that has considered religion has found it to be a positive factor. One review found that religious people had lower blood pressures and lower hypertension-related morbidity and mortality.⁴ The authors posit that members of these religious groups may follow health-promoting behaviors such as abstaining from alcohol and tobacco. But they also speculate that an additional health benefit may result from a stronger sense of peace and purpose.

In one prospective cohort study,⁵ attendance at a church or other religious institution at baseline predicted less disability during most of a 6- to 12-year follow-up.

Interestingly, the health benefits of religion are not confined to those who follow healthpromoting behaviors. One study found that smokers who rated themselves as not religious were more than seven times as likely to have abnormally high diastolic blood pressure than were religious smokers, and smokers who rarely attended church were four times as likely to have an abnormal diastolic blood pressure.⁶

RELIGION DECREASES DEPRESSION

Religious commitment seems to have a strong protective effect against depression and sui-

cide. A population study in Maryland found that people who did not attend church were four times as likely to commit suicide as frequent churchgoers.⁷ Stack⁸ found that the church attendance rate in a population was more closely associated with the suicide rate than any other factor studied, including the employment rate.

Religious commitment may also lower the risk of substance abuse. In a long-term study of male medical students, those who reported not having a religious affiliation when surveyed during medical school were much more likely to report an alcohol abuse problem 20 years later than were their religious colleagues.⁹ A large study in North Carolina found that a history of alcohol problems was less common among weekly churchgoers and people who called themselves "born-again."¹⁰

One study found that religious commitment also helps patients avoid depression triggered by serious illness.¹¹

RELIGION AND LONGEVITY

In light of these findings, it is not surprising to find evidence that religion may also be associated with longer life. A nationwide study showed that religious attendance added an average of 7 years of life, and for African-Americans, the figure was 14 years.¹² This follows up on previous research including a 28year follow-up of more than 5,000 residents of Alameda County, California, finding that those who attended religious services weekly or more were 23% less likely to die than infrequent attendees. In addition, once people started to attend services, they made healthier lifestyle choices, becoming more likely to stop smoking, increase exercising, expand their social network, and remain married.13

INTERPRETING THE FINDINGS

There are methodological problems with the studies of religion and health, and many are open to more alternative interpretations. For example, many of the studies measured only religious denomination or single items of religious attitudes, which are clearly inadequate measures of how people actually feel about, or practice, their faith. One could also argue that it is health that affects religious attendance, because healthier people are more likely to attend church regularly. Depression, substance addiction, physical illness, or disability is likely to increase social isolation and reduce levels of many activities, including church attendance.

Also, there are a number of ways that religion could affect one's health: by adding social or psychological support (or both) to people's lives, by giving a perspective on stress that reduces its negative impact, or by encouraging people to avoid risky behaviors such as drinking alcohol to excess.

Better-controlled studies are needed that better assess the mediating factors of religiousness, with measures of the depth or importance of the respondents' spirituality.

HOW PHYSICIANS SHOULD APPROACH RELIGION

As physicians, we should recognize that clergy, and chaplains in particular, could be valuable to our patients and to our own learning. In my medical training, we were not even informed about what hospital chaplains did. However, at many medical schools today, chaplains bring medical students on rounds and teach them about death and dying. Indeed, at more than half of US medical schools there are now courses addressing spiritual issues in medicine; many of these are required. These courses often include taking a spiritual history, a review of the research, and a discussion of the spiritual issues involved in death and dying.¹⁴

We should also consider asking tactful, sensitive questions about our patients' religion or spirituality to better respond to their preferences, especially in populations that have a high likelihood of relying on religion, such as the severely or chronically ill. Such questions could include, "Is religion helpful to you in handling your illness?" or a more open-ended question, "What gets you through the tough times in your life?" For patients who answer "yes," or "my faith, spirituality, or religion," an appropriate follow-up question might be, "What can I do to support or address spiritual or religious issues in your life?"

Some patients do not find religion important and do not want spirituality addressed in

Ask: "Is religion helpful to you in handling your illness?"

Dear Doctor:

As editors, we'd like you to look into every issue, every page of the Cleveland Clinic Journal of Medicine. We'd like to know...

1 How many issues do you look into? Here's our goal: Mall Most Half Few

2 How do you read the average issue? Here's our goal:

Cover-to-cover Most articles Selected articles

We put it in writing... please put it in writing for us. We want to hear from you.

CLEVELAND CLINIC JOURNAL OF MEDICINE The Cleveland Clinic Foundation 9500 Euclid Avenue, NA32 Cleveland, Ohio 44195

PHONE 216.444.2661 **FAX** 216.444.9385 **E-MAIL** ccjm@ccf.org MEDICAL GRAND ROUNDS

medical care.² However, for the more than 70% of the population for whom religion is central to life, treatment approaches that are not sensitive to spirituality may leave patients feeling disaffected. These people probably prefer medical care that is sympathetic or at least sensitive to their religious perspective.¹⁵ For those patients to whom religion is important, we should find ways to support their faith, such as referring them to chaplains or their clergy. We should learn to recognize the clinical relevance and importance of spirituality and religion in many of our patient's lives, for these factors may turn out to be good for one's health.

For more information on research concerning spirituality and health, visit NIHR's Web page at www.nihr.org or call 800-580-NIHR (6447).

REFERENCES

- Princeton Religious Research Center. Religion in America, 1993–94. Princeton, NJ: Princeton Religious Research Center; 1994.
- Ehman JW, Ott BB, Short TH. Do patients want physicians to inquire about their spiritual or religious beliefs if they become gravely ill? Arch Intern Med 1999; 159:1803–1806.
- King DE, Bushwick B. Beliefs and attitudes of hospital patients about faith healing and prayer. J Fam Pract 1994; 39:349–352.
- Levin JS, Vanderpool HY. Is religion therapeutically significant for hypertension? Soc Sci Med 1989; 29:69–78.
- Idler EL, Kasl SV. Religion, disability, depression, and the timing of death. Am J Sociol 1992; 97:1052–1079.
- Larson DB, Koenig HG, Kaplan BH, Greenberg RS, Logue E, Tyroler HA. The impact of religion on men's blood pressure. J Religion Health 1989; 28:265–278.
- 7. Comstock GW, Partridge KB. Church attendance and health. J Chronic Dis 1972; 25:665–672.
- Stack S. The effect of the decline in institutionalized religion on suicide: 1954–1978. J Sci Study Religion 1983; 22:239–252.
- Moore RD, Mead L, Pearson T. Youthful precursors of alcohol abuse in physicians. Am J Med 1990; 88:332–336.
- Koenig HG, George LK, Meador KG, Blazer DG, Ford SM. Religious practices and alcoholism in a Southern adult population. Hosp Community Psychiatry 1994; 45:225–231.
- Koenig HG, Cohen HJ, Blazer DG, et al. Religious coping and depression in elderly, hospitalized, medically ill men. Am J Psychiatry 1992; 149:1693–1700.
- Hummer RA, Rogers RG, Nam CB, et al. Religious involvement and US adult mortality. Demography 1999; 36:1–13.
- Strawbridge WJ, Cohen RD, Shema SJ, et al. Frequent attendance at religious services and mortality over 28 years. Am J Public Health 1997; 87:957–961.
- 14. Puchalski CM, Larson DB. Developing curricula in spirituality and medicine. Acad Med 1998 73:970–974.

CME ANSWERS Answers to the CREDIT TEST on page 143 of this issue 1 A 2 A 3 E 4 E 5 B 6 B 7 B 8 A 9 B

84 CLEVELAND CLINIC JOURNAL OF MEDICINE VOLUME 67 • NUMBER 2 FEBRUARY 2000 Downloaded from www.ccjm.org on May 3, 2025. For personal use only. All other uses require permission.