

**STEVEN R. LENTZ, MD, PhD**Department of Internal Medicine,
University of Iowa; Veterans Affairs Medical
Center, Iowa City**WILLIAM G. HAYNES, MD**General Clinical Research Center,
University of Iowa, Iowa City

Homocysteine: Is it a clinically important cardiovascular risk factor?

ABSTRACT

Elevated plasma homocysteine is associated with an increased risk of myocardial infarction, stroke, and venous thromboembolism. Folic acid and other B vitamins lower plasma homocysteine levels, but whether this therapy confers a clinical benefit has yet to be determined. Until we know the results of ongoing clinical trials of homocysteine-lowering therapy, testing for and treating elevated homocysteine is probably justified only in patients with known cardiovascular disease or who are at high risk.

KEY POINTS

High homocysteine levels are believed to directly increase the risk of cardiovascular disease, but they may actually result from cardiovascular events or merely be markers of cardiovascular disease.

Although homocysteine-lowering therapy has not been proven to lower cardiovascular risk, it is inexpensive and presumed safe.

Homocysteine can be lowered to normal levels in most patients with mildly or moderately elevated levels with either folic acid 0.4 to 5.0 mg/day, vitamin B₁₂ 0.5 to 1.0 mg/day, or both. The treatment may be less effective in patients with renal disease.

If treatment is undertaken, the goal of therapy should be a homocysteine level of less than 10 μ mol/L in patients at high cardiovascular risk.

Dr. Lentz's work is supported by the Office of Research and Development, United States Department of Veterans Affairs, and National Institutes of Health grants HL63943, NS24621, and HL62984. Dr. Haynes' work is supported by National Institutes of Health grants HL58972 and HL14388.

FOR ALMOST TWO DECADES we have known that a high plasma homocysteine level (hyperhomocysteinemia) is associated with increased cardiovascular risk. But whether elevated homocysteine causes cardiovascular disease or is a consequence of it remains unknown.

Adding folic acid and other B vitamins to the diet is effective in lowering levels, but whether it improves the clinical outlook is also still uncertain and is the focus of several ongoing clinical trials.

In this article, we summarize the causes and clinical manifestations of elevated homocysteine levels, proposed mechanisms linking homocysteine to vascular disease, and indications for testing for and treating hyperhomocysteinemia.

CAUSES OF HYPERHOMOCYSTEINEMIA

Homocysteine, a sulfur-containing amino acid, is a key intermediate in methionine metabolism (FIGURE 1). It is produced as a byproduct of methyl transfer reactions, which are important for the synthesis of nucleic acids, methylated proteins, neurotransmitters, and phospholipids.¹

Hyperhomocysteinemia can be caused by genetic defects, nutritional deficiencies, renal dysfunction, alcoholism, hypothyroidism, or certain medications (TABLE 1).

Severe hyperhomocysteinemia. People with rare inborn errors of methionine metabolism have very high levels of homocysteine in the plasma ($>100 \mu$ mol/L) and in the urine and a very high incidence of vascular disease.

How homocysteine is produced and metabolized




FIGURE 1

FROM BALLAL RS, JACOBSEN DW, ROBINSON K. HOMOCYSTEINE: UPDATE ON A NEW RISK FACTOR. CLEVE CLIN J MED 1997; 64:543-549.

This condition, recognized for more than 30 years, is called “hereditary homocystinuria.”²

Most cases of severe hyperhomocysteinemia are caused by autosomal-recessive defects in cystathionine beta-synthase, which also cause skeletal deformities, ocular abnormalities, mental retardation, hepatic steatosis, and premature death.³ Untreated patients have about a 50% chance of having a myocardial infarction, stroke, or venous thromboembolism before age 30.³

Moderate hyperhomocysteinemia (plasma total homocysteine 30–100 $\mu\text{mol/L}$) may be seen in patients with moderate vitamin B₁₂ deficiency, severe folate deficiency, or renal

failure. The cause of hyperhomocysteinemia in renal disease is not well understood, but may involve decreased clearance of homocysteine by the kidneys or extrarenal tissues.⁴

Mild hyperhomocysteinemia (plasma total homocysteine 10–30 $\mu\text{mol/L}$) is common in the general population, especially the elderly.^{5,6} It is often caused by renal insufficiency or mild to moderate vitamin B₁₂ deficiency. Folate deficiency is no longer a common cause of mild hyperhomocysteinemia in the United States and Canada since cereal grains have been fortified with folic acid.⁷

Mild hyperhomocysteinemia may also be caused by genetic factors, including heterozy-



gosity for mutations in the gene for cystathionine beta-synthase or homozygosity for a common polymorphism in the gene that encodes the enzyme 5,10-methylene tetrahydrofolate reductase (*MTHFR*), specifically a C-to-T substitution at base 677 of the gene (*MTHFR* 677 C→T).⁸ The clinical value of testing for these mutations has not been proven, and it is unlikely to provide more risk information than the homocysteine level. The *MTHFR* 677 C→T genotype appears to be associated with hyperhomocysteinemia and increased risk mainly in people with low folate intake.⁸

■ HOMOCYSTEINE AS A RISK FACTOR

A number of retrospective (case-control and observational) and prospective studies done over the past 15 years indicate that homocysteine is a graded, independent risk factor for myocardial infarction, stroke, and venous thromboembolism.^{9–11}

What is 'abnormal'?

The normal reference range for plasma total homocysteine is usually defined as the 2.5th to 97.5th percentile interval for presumably healthy people. The lower limit is typically 5 $\mu\text{mol/L}$, but the upper limit varies considerably among clinical laboratories. Furthermore, in different populations, the upper limit may vary between 10 and 20 $\mu\text{mol/L}$, depending on age (levels increase with age), sex (levels are higher in men than in women), ethnic group, and dietary intake of folate.¹² Adding to the confusion, mean levels of homocysteine have decreased in the United States since 1998, when enriched flour began to be fortified with folic acid to prevent neural tube defects in newborns.¹³

Rather than defining a level of homocysteine as either normal or abnormal, it may be more useful to consider homocysteine, like cholesterol and C-reactive protein, as a graded risk factor for cardiovascular disease.^{14,15} A meta-analysis¹⁶ found that for every 2.5- $\mu\text{mol/L}$ increase in plasma total homocysteine, the risk of myocardial infarction increases by about 10% and the risk of stroke increases by about 20%. The relationship between homocysteine and risk appears to hold for plasma concentrations of total homo-

TABLE 1

Causes of hyperhomocysteinemia

Severe hyperhomocysteinemia (> 100 $\mu\text{mol/L}$)

Hereditary homocystinuria (eg, homozygosity for defects in cystathionine beta-synthase, 5,10-methylene-tetrahydrofolate reductase, or other enzymes of methionine metabolism)
Hereditary disorders of vitamin B₁₂ utilization
Severe deficiency of vitamin B₁₂

Moderate hyperhomocysteinemia (30–100 $\mu\text{mol/L}$)

Renal failure
Moderate vitamin B₁₂ deficiency
Severe folate deficiency

Mild hyperhomocysteinemia (10–30 $\mu\text{mol/L}$)

Heterozygosity for defects in cystathionine beta-synthase
Homozygosity for the C677T polymorphism of 5,10-methylene-tetrahydrofolate reductase
Renal insufficiency
Renal transplantation
Mild folic acid deficiency
Mild vitamin B₁₂ deficiency
Vitamin B₆ deficiency
Hypothyroidism
Alcoholism
Medications (niacin, fibrates, methotrexate, isoniazid, levodopa, theophylline, phenytoin, nitrous oxide, trimethoprim)

cysteine between 10 and 30 $\mu\text{mol/L}$.

A more sensitive way to identify people at increased cardiovascular risk may be to measure plasma homocysteine after an oral methionine load. However, methods for administering methionine and collecting samples must be standardized before this can be generally recommended.^{12,17}

Is homocysteine a consequence, cause, or marker of disease?

It has not been established whether homocysteine is a consequence, a cause, or a marker of cardiovascular disease.

Consequence? Evidence that elevated homocysteine may be a consequence of vascular disease is that the relative risk associated with mild homocysteine elevation has tended to be higher in retrospective studies (in which homocysteine was measured after a vascular event) than in prospective studies (in which it was measured before an event).¹⁶

Cause? However, when analysis is limited

to prospective data and adjusted for other risk factors, hyperhomocysteinemia still emerges as a modest but significant risk factor for cardiovascular events.¹⁶ Elevated total homocysteine is also a strong predictor of mortality in patients with preexisting cardiovascular disease or other risk factors.^{12,18}

Other evidence that it is a cause is that vascular manifestations are common in patients with severe hyperhomocysteinemia caused either by defects in the cystathionine beta-synthase-dependent transsulfuration pathway that produce elevated levels of both homocysteine and methionine or by defects in the methionine synthase-dependent remethylation pathway that produce elevated levels of homocysteine but not methionine.

In addition, estimates of cardiovascular risk of hyperhomocysteinemia caused by genetic variation in 5,10-methylene tetrahydrofolate reductase, which are relatively free from confounding due to reverse causality,¹⁹ are similar to those estimated from prospective studies of homocysteine levels.^{8,15}

Experimental evidence also supports the theory that high homocysteine causes vascular problems: induced hyperhomocysteinemia in animals produces vascular abnormalities, including endothelial dysfunction and vascular hypertrophy, that are very similar to those seen with other risk factors such as hypertension, diabetes, and hypercholesterolemia.²⁰ Hyperhomocysteinemia also accelerates atherosclerosis progression in apolipoprotein E-deficient mice.^{21–23}

In human subjects, acute hyperhomocysteinemia induced by oral methionine loading causes endothelial dysfunction.²⁴

Possible mechanisms of hyperhomocysteinemia's vascular effects include impaired release of endothelium-derived nitric oxide, oxidative injury, accelerated atherosclerosis, altered hemostatic balance, endoplasmic reticulum stress, and activation of inflammatory pathways.^{25,26}

Marker? It is also plausible that the relationship between hyperhomocysteinemia and cardiovascular disease is indirect, and is confounded by other factors (eg, renal insufficiency or deficiencies of folate, vitamin B₁₂, or vitamin B₆) that influence both homocysteine levels and cardiovascular risk.

■ CLINICAL MEASUREMENT

Because of increasing interest in hyperhomocysteinemia as a cardiovascular risk factor, most clinical reference laboratories can now measure plasma homocysteine levels. Clinical assays measure total homocysteine, which includes the thiol homocysteine and its protein-bound and free disulfide derivatives.^{27,28}

Other than to help predict cardiovascular risk, reasons for measuring homocysteine are to screen for inborn errors of methionine metabolism and as an adjunctive test for vitamin B₁₂ deficiency.

Blood samples can be collected in either the fasting or nonfasting state. Plasma must be isolated quickly (within 1 hour at room temperature or 8 hours if the sample is kept on ice) to avoid spurious elevation of homocysteine due to leakage from erythrocytes and leukocytes. A reducing agent is added to the sample to convert disulfide forms to homocysteine, which is detected using either chromatographic or immunologic methods. The different methods give comparable results, but certified reference materials are lacking, and standardization is needed to minimize variation among laboratories.¹²

■ HOMOCYSTEINE-LOWERING THERAPY

Homocysteine-lowering therapy is life-saving for patients with severe hyperhomocysteinemia due to autosomal-recessive deficiency of cystathionine beta-synthase. Approximately 50% of patients respond to treatment with vitamin B₆ in pharmacological doses.³ Adjunctive therapy may include methionine restriction, cysteine supplementation, betaine, vitamin B₁₂, or folic acid.^{29,30} Interestingly, long-term therapy substantially decreases vascular risk despite only partially correcting elevated plasma homocysteine.³⁰

In people with mild or moderate hyperhomocysteinemia, B vitamins are highly effective in lowering homocysteine. A meta-analysis of 12 randomized trials performed before folic acid fortification was instituted³¹ found that treatment with folic acid 0.5 to 5 mg/day decreased homocysteine levels by 25%, and that the addition of vitamin B₁₂ 0.5 mg/day decreased homocysteine by another 7%.

The risk of MI increases about 10% for every 2.5 μ mol/L increase in homocysteine



Adding vitamin B₆ did not improve the homocysteine-lowering effect of folic acid and vitamin B₁₂.

Now that cereal grains are fortified with folic acid, additional folic acid supplementation may not work as well. Nevertheless, it is possible to achieve normal levels of plasma homocysteine in most people with mild or moderate hyperhomocysteinemia, although patients with renal failure may be resistant.³²

Benefit of therapy uncertain

Several randomized clinical trials are under way in people with coronary heart disease, stroke, or venous thromboembolism to test the effectiveness of homocysteine-lowering therapy with folic acid and vitamin B₁₂ (with or without vitamin B₆) in preventing secondary events.³³ Results from most of the trials will not be available until 2005 or 2006, but a few have been completed or reported in preliminary form.³⁴⁻⁴²

Results thus far are mixed. Two trials examined the effect of homocysteine-lowering therapy on the rate of restenosis following percutaneous coronary interventions. One found a lower rate of restenosis in the treated group,^{34,35} while the other found treatment to be detrimental.³⁷ Similar inconsistencies are emerging from other intervention trials.³⁸⁻⁴²

The conflicting results may be partly due to inadequate statistical power,³³ especially in the United States and Canada, where folic acid fortification has decreased homocysteine levels in the control population.⁴³ This appears to have been a factor in the recent Vitamin Intervention for Stroke Prevention (VISP) trial, which found no effect of homocysteine-lowering therapy on vascular outcomes in patients with prior stroke.³⁶

RECOMMENDATIONS

Given that the jury is still out on the benefit of homocysteine-lowering therapy, should physicians detect and treat hyperhomocysteinemia?

REFERENCES

1. Selhub J. Homocysteine metabolism. *Annu Rev Nutr* 1999; 19:217-246.
2. McCully KS. Vascular pathology of homocysteinemia: implications for the pathogenesis of arteriosclerosis. *Am J Pathol* 1969; 56:111-128.
3. Mudd SH, Skovby F, Levy HL, et al. The natural history of homocystinuria due to cystathionine beta-synthase deficiency. *Am J Hum Genet* 1985; 37:1-31.
4. Friedman AN, Bostom AG, Selhub J, Levey AS, Rosenberg IH. The kidney and homocysteine metabolism. *J Am Soc Nephrol* 2001; 12:2181-2189.
5. Selhub J, Jacques PF, Wilson PW, Rush D, Rosenberg IH. Vitamin status and intake as primary determinants of homocysteinemia in an elderly

From a strictly evidence-based perspective, routine screening for hyperhomocysteinemia and prescribing vitamins to lower homocysteine are unwarranted. The American Heart Association⁴⁴ and a European expert panel¹² advise against general screening for hyperhomocysteinemia, and the US Preventive Services Task Force has concluded that evidence is insufficient to recommend vitamin supplements to prevent cardiovascular disease.⁴⁵

On the other hand, there is good reason to believe that homocysteine is an independent cardiovascular risk factor and that homocysteine-lowering therapy ultimately may prove to have a modest clinical benefit. Treating with either folic acid 0.4-5.0 mg/day or vitamin B₁₂ 0.5-1.0 mg/day or both is inexpensive and presumably safe, and is likely to be cost-effective for preventing cardiovascular events if ongoing clinical trials confirm that homocysteine-lowering therapy is beneficial.³²

Even without definitive evidence of clinical benefit, a case can be made for detecting and treating hyperhomocysteinemia in selected patients (ie, those with a history of premature cardiovascular disease, stroke, or venous thromboembolism, or those thought to be at high risk because other risk factors are present). In such cases, treatment is unlikely to cause harm and may produce an important benefit.

When starting homocysteine-lowering therapy, the goal should be to maintain the plasma concentration of total homocysteine within the laboratory's low-normal reference range. An American Heart Association science advisory statement suggested that a level of less than 10 $\mu\text{mol/L}$ is a reasonable target for patients at high cardiovascular risk.⁴⁴ A European expert panel has proposed a slightly higher target level (13-15 $\mu\text{mol/L}$).¹² Because excess folate intake may exacerbate neurological complications in people with occult pernicious anemia, vitamin B₁₂ status should be determined before starting folic acid therapy.

Blood samples for homocysteine must be processed quickly



- population. *JAMA* 1993; 270:2693–2698.
6. Jacques PF, Rosenberg IH, Rogers G, et al. Serum total homocysteine concentrations in adolescent and adult Americans: results from the third National Health and Nutrition Examination Survey. *Am J Clin Nutr* 1999; 69:482–489.
 7. Liaugaudas G, Jacques PF, Selhub J, Rosenberg IH, Bostom AG. Renal insufficiency, vitamin B(12) status, and population attributable risk for mild hyperhomocysteinemia among coronary artery disease patients in the era of folic acid-fortified cereal grain flour. *Arterioscl Thromb Vasc Biol* 2001; 21:849–851.
 8. Klerk M, Verhoef P, Clarke R, Blom HJ, Kok FJ, Schouten EG; MTHFR Studies Collaboration Group. *MTHFR* 677C→T polymorphism and risk of coronary heart disease: a meta-analysis. *JAMA* 2002; 288:2023–2031.
 9. Boushey CJ, Beresford SA, Omenn GS, Motulsky AG. A quantitative assessment of plasma homocysteine as a risk factor for vascular disease. Probable benefits of increasing folic acid intakes. *JAMA* 1995; 274:1049–1057.
 10. Eikelboom JW, Lonn E, Genest J Jr, Hankey G, Yusuf S. Homocyst(e)ine and cardiovascular disease: a critical review of the epidemiologic evidence. *Ann Intern Med* 1999; 131:363–375.
 11. den Heijer M, Keijzer MB. Hyperhomocysteinemia as a risk factor for venous thrombosis. *Clin Chem Lab Med* 2001; 39:710–713.
 12. Refsum H, Smith AD, Ueland PM, et al. Facts and recommendations about total homocysteine determinations: an expert opinion. *Clin Chem* 2004; 50:3–32.
 13. Jacques PF, Selhub J, Bostom AG, Wilson PW, Rosenberg IH. The effect of folic acid fortification on plasma folate and total homocysteine concentrations. *N Engl J Med* 1999; 340:1449–1454.
 14. Graham IM, Daly LE, Refsum HM, et al. Plasma homocysteine as a risk factor for vascular disease. The European Concerted Action Project. *JAMA* 1997; 277:1775–1781.
 15. Wald DS, Law M, Morris JK. Homocysteine and cardiovascular disease: evidence on causality from a meta-analysis. *BMJ* 2002; 325:1202.
 16. Homocysteine Studies Collaboration. Homocysteine and risk of ischemic heart disease and stroke: a meta-analysis. *JAMA* 2002; 288:2015–2022.
 17. Silberberg J, Dudman N. Methionine loading. In: Carmel R, Jacobsen DW, editors. *Homocysteine in Health and Disease*. Cambridge, UK: Cambridge University Press; 2001:212–219.
 18. Nygård O, Nordrehaug JE, Refsum H, Ueland PM, Farstad M, Vollset SE. Plasma homocysteine levels and mortality in patients with coronary artery disease. *N Engl J Med* 1997; 337:230–236.
 19. Davey Smith G, Ebrahim S. 'Mendelian randomization': can genetic epidemiology contribute to understanding environmental determinants of disease? *Int J Epidemiol* 2003; 32:1–22.
 20. Lentz SR. Homocysteine and cardiovascular physiology. In: Carmel R, Jacobsen DW, editors. *Homocysteine in Health and Disease*. Cambridge, UK: Cambridge University Press; 2001:441–450.
 21. Hofmann MA, Lalla E, Lu Y, et al. Hyperhomocysteinemia enhances vascular inflammation and accelerates atherosclerosis in a murine model. *J Clin Invest* 2001; 107:675–683.
 22. Zhou J, Möller J, Danielsen CC, et al. Dietary supplementation with methionine and homocysteine promotes early atherosclerosis but not plaque rupture in ApoE-deficient mice. *Arterioscl Thromb Vasc Biol* 2001; 21:1470–1476.
 23. Wang H, Jiang X, Yang F, et al. Hyperhomocysteinemia accelerates atherosclerosis in cystathionine beta-synthase and apolipoprotein E double knock-out mice with and without dietary perturbation. *Blood* 2003; 101:3901–3907. Epub 2002 Dec 27.
 24. Kanani PM, Sinkey CA, Browning RL, Allaman M, Knapp HR, Haynes WG. Role of oxidant stress in endothelial dysfunction produced by experimental hyperhomocyst(e)inemia in humans. *Circulation* 1999; 100:1161–1168.
 25. Faraci FM. Hyperhomocysteinemia: a million ways to lose control. *Arterioscl Thromb Vasc Biol* 2003; 23:371–373.
 26. Austin RC, Lentz SR, Werstuck GH. Role of hyperhomocysteinemia in endothelial dysfunction and atherothrombotic disease. *Cell Death Differ* 2004; 11:556–564.
 27. Refsum H, Ueland PM, Nygård O, Vollset SE. Homocysteine and cardiovascular disease. *Annu Rev Med* 1998; 49:31–62.
 28. Mudd SH, Finkelstein JD, Refsum H, et al. Homocysteine and its disulfide derivatives: a suggested consensus terminology. *Arterioscl Thromb Vasc Biol* 2000; 20:1704–1706.
 29. Wilcken DE, Wilcken B. The natural history of vascular disease in homocystinuria and the effects of treatment. *J Inher Metab Dis* 1997; 20:295–300.
 30. Yap S, Boers GH, Wilcken B, et al. Vascular outcome in patients with homocystinuria due to cystathionine beta-synthase deficiency treated chronically: a multicenter observational study. *Arterioscler Thromb Vasc Biol* 2001; 21:2080–2085.
 31. Homocysteine Lowering Trialists' Collaboration. Lowering blood homocysteine with folic acid based supplements: meta-analysis of randomised trials. *BMJ* 1998; 316:894–898.
 32. Tice JA, Ross E, Coxson PG, et al. Cost-effectiveness of vitamin therapy to lower plasma homocysteine levels for the prevention of coronary heart disease: effect of grain fortification and beyond. *JAMA* 2001; 286:936–943.
 33. Clarke R. Design of clinical trials to test the homocysteine hypothesis of vascular disease. In: Carmel R, Jacobsen DW, editors. *Homocysteine in health and disease*. Cambridge, UK: Cambridge University Press; 2001:477–484.
 34. Schnyder G, Roffi M, Pin R, et al. Decreased rate of coronary restenosis after lowering of plasma homocysteine levels. *N Engl J Med* 2001; 345:1593–1600.
 35. Schnyder G, Roffi M, Flammer Y, Pin R, Hess OM. Effect of homocysteine-lowering therapy with folic acid, vitamin B₁₂, and vitamin B₆ on clinical outcome after percutaneous coronary intervention: the Swiss Heart study: a randomized controlled trial. *JAMA* 2002; 288:973–979.
 36. Toole JF, Malinow MR, Chambless LE, et al. Lowering homocysteine in patients with ischemic stroke to prevent recurrent stroke, myocardial infarction, and death: the Vitamin Intervention for Stroke Prevention (VISP) randomized controlled trial. *JAMA* 2004; 291:565–575.
 37. Lange H, Suryapranata H, De Luca G, et al. Folate therapy and in-stent restenosis after coronary stenting. *N Engl J Med* 2004; 350:2673–2681.
 38. Vermeulen EG, Stehouwer CD, Twisk JW, et al. Effect of homocysteine-lowering treatment with folic acid plus vitamin B₆ on progression of subclinical atherosclerosis: a randomised, placebo-controlled trial. *Lancet* 2000; 355:517–522.
 39. Liem A, Reynierse-Buitenwerf GH, Zwinderman AH, Jukema JW, van Veldhuisen DJ. Secondary prevention with folic acid: effects on clinical outcomes. *J Am Coll Cardiol* 2003; 41:2105–2113.
 40. Baker F, Blackwood S, Hunt J, et al. Blinded comparison of folic acid and placebo in patients with ischemic heart disease: an outcome trial [abstract]. *Circulation* 2003; 106(suppl 2):II-741.
 41. Marcucci R, Zanazzi M, Bertoni E, et al. Vitamin supplementation reduces the progression of atherosclerosis in hyperhomocysteinemic renal transplant recipients. *Transplantation* 2003; 75:1551–1559.
 42. den Heijer M, Willems HJ, Blom HJ, et al. Homocysteine lowering by B vitamins and the prevention of secondary deep-vein thrombosis and pulmonary embolism: a randomized, placebo-controlled, double-blind trial [abstract]. *J Thromb Haemost* 2003; 1(suppl 1):161.
 43. Bostom AG, Selhub J, Jacques PF, Rosenberg IH. Power shortage: clinical trials testing the "homocysteine hypothesis" against a background of folic acid-fortified cereal grain flour. *Ann Intern Med* 2001; 135:133–137.
 44. Malinow MR, Bostom AG, Krauss RM. Homocyst(e)ine, diet, and cardiovascular diseases: a statement for healthcare professionals from the Nutrition Committee, American Heart Association. *Circulation* 1999; 99:178–182.
 45. Morris CD, Carson S. Routine vitamin supplementation to prevent cardiovascular disease: a summary of the evidence for the U.S. Preventive Services Task Force. *Ann Intern Med* 2003; 139:56–70.

ADDRESS: Steven R. Lentz, MD, PhD, Department of Internal Medicine, C303 GH, The University of Iowa, Iowa City, IA 52242; e-mail steven-lentz@uiowa.edu.