



FROM THE OFFICE OF DR.

Nutrition: Know the facts

Most packaged foods have a Nutrition Facts label. For a healthier you, use this tool to make smart food choices quickly and easily. Try these tips:

Make your calories count. Look at the calories on the label and compare them with the nutrients listed to decide whether the food is worth eating. When a food item has more than 400 calories per serving, it is high in calories.

Know your fats. Look for foods low in saturated fats, *trans*-fats, and cholesterol to help reduce the risk of heart disease. Most of the fats you eat should be polyunsaturated and mono-unsaturated fats. Keep total fat intake between 20% and 35% of total calories.

To get enough fiber (25 grams per day), half of your intake of grains should be whole grains, such as in whole-grain bread.

Get enough of these vitamins and minerals.

Amount Per Serving		% Daily Value*	
Nutrition Facts			
Serving Size ½ cup (114g)			
Servings Per Container 4			
Calories 90 Calories from Fat 30			
Total Fat 3g		5%	
Saturated Fat 0g		0%	
Cholesterol 0mg		0%	
Sodium 300mg		13%	
Total Carbohydrate 13g		4%	
Dietary Fiber 3g		12%	
Sugars 3g			
Protein 3g			
Vitamin A 80% • Vitamin C 60%			
Calcium 4% • Iron 4%			
* Percent Daily Values are based on a 2,000 calorie diet. Your daily values may be higher or lower depending on your calorie needs:			
	Calories:	2,000	2,500
Total Fat	Less than	65g	80g
Sat Fat	Less than	20g	25g
Cholesterol	Less than	300mg	300mg
Sodium	Less than	2,400mg	2,400mg
Total Carbohydrate		300g	375g
Dietary Fiber		25g	30g
Calories per gram:			
Fat 9 • Carbohydrate 4 • Protein 4			

Check servings and calories. Look at the serving size and how many servings you are actually consuming. If you eat all four servings in this package you get not 90 but 360 calories. Servings may be smaller than you think: see <http://hin.nhlbi.nih.gov/portion>.

Use the Percent Daily Value (% Daily Value) column when possible: 5% Daily Value or less is low, 20% or more is high.

Reduce sodium (salt), increase potassium. Eating less than 2,300 mg of sodium (about 1 teaspoon of salt) per day may reduce the risk of high blood pressure. Most of the sodium people eat comes from processed foods, not from the salt-shaker. Foods high in potassium include fruits and vegetables.

Don't sugarcoat it. Since sugars contribute calories with few, if any, nutrients, look for foods and beverages low in added sugars. Read the ingredient list and make sure that added sugars are not one of the first few ingredients. Some names for added sugars include sucrose, glucose, high-fructose corn syrup, corn syrup, and fructose.

ADAPTED FROM US DEPARTMENT OF HEALTH AND HUMAN SERVICES, US DEPARTMENT OF AGRICULTURE. FINDING YOUR WAY TO A HEALTHIER YOU: BASED ON THE DIETARY GUIDELINES FOR AMERICANS. WWW.HEALTHIERUS.GOV/DIETARYGUIDELINES.



This information is provided by your physician and the *Cleveland Clinic Journal of Medicine*. It is not designed to replace a physician's medical assessment and judgment.

This page may be reproduced noncommercially to share with patients. Any other reproduction is subject to *Cleveland Clinic Journal of Medicine* approval. Bulk color reprints available by calling 216-444-2661.

For patient information on hundreds of health topics, see the Patient Education and Health Information web site, www.clevelandclinic.org/health