

SYED KASHIF MAHMOOD, MDDepartment of Internal Medicine,
Cleveland Clinic**J. WALTON TOMFORD, MD**Department of Infectious Diseases,
Cleveland Clinic**STEVEN ROSENBLATT, MD**Department of General Surgery,
Cleveland Clinic**STEVEN GORDON, MD**Chairman, Department of Infectious
Diseases, Cleveland Clinic

The Clinical Picture

Dropped gallstones disguised as a liver abscess



FIGURE 1. Computed tomography scan of the abdomen with contrast shows a possible hepatic lesion (arrow).



FIGURE 2. Computed tomography scan of the pelvis with contrast shows a possible hepatic lesion (arrow).

A 67-YEAR-OLD RETIRED MAN presents to his internist with a 3-month history of abdominal discomfort in the right upper quadrant on deep breathing. He has no other abdominal complaints, but he mentions that he underwent laparoscopic cholecystectomy 3 months ago for gallstone pancreatitis.

A physical examination and preliminary laboratory work are inconclusive, but the internist, concerned about the ongoing symptoms, orders a computed tomographic (CT) study of the abdomen (FIGURE 1) and pelvis (FIGURE 2), with contrast, and the resulting CT report mentions a possible hepatic lesion, which in turn raises the possibility of a hepatic abscess. However, on further review of the

scans with a radiologist, the lesion appears perihepatic rather than intrahepatic.

The surgeon who had performed the laparoscopic cholecystectomy is consulted and says that he had noted no hepatic or perihepatic lesion at the time of the operation. He adds, however, that the operation had been technically difficult because of inflammation, and that gallstones were dropped during retraction of the gallbladder and could not be retrieved, despite every effort. The presence of dropped gallstones therefore raises suspicion of abscess.

A biopsy specimen obtained with CT guidance shows chronic inflammation but is sterile on aerobic culture. There is no evidence of malignancy. Because of concern for



FIGURE 3. Pus was noted after incision of the abscess cavity.



FIGURE 4. The gallstone (arrow) was seen in the abscess cavity after evacuation of pus.

underlying infection, the infectious disease staff recommends empirical treatment with a 4-week course of ampicillin-sulbactam (Unasyn). At completion of the antibiotic course, the patient's symptoms have resolved.

In another case, a 66-year-old woman presented to the infectious disease department with a persistent subdiaphragmatic abscess 2 years after undergoing laparoscopic cholecystectomy. Despite CT-guided drainage of the abscess followed by several courses of antibiotics, the abscess did not resolve. The patient was then evaluated by a general surgeon who, considering the recurrent nature of her abscess, suspected that the inflammation might be a foreign-body reaction to a dropped gallstone. The patient was taken for surgical evacuation, during which a chronic abscess was found and was unroofed and drained of pus (FIGURE 3). A gallstone was found in the abscess cavity (FIGURE 4).

■ LAPAROSCOPY'S DRAWBACKS

In the United States, more than 700,000 laparoscopic cholecystectomies are performed each year,¹ and the number is growing. The key advantages of laparoscopic cholecystectomy over the open procedure are smaller incisions, less postoperative pain, and a shorter recovery time. On the other hand, limited visualization, pneumoperitoneum, and other technical challenges of laparoscopy increase the risk of

bile duct injury and dropped gallstones. As many as a third of all laparoscopic cholecystectomies are complicated by dropped gallstones.¹⁻⁴ Gallstones may also be dropped during open cholecystectomy, but the larger operating field makes them easier to retrieve.⁵

Complications of dropped stones, though rare, can include localized or systemic infection, inflammation, fibrosis, adhesion, cutaneous sinus formation, ileus, and abscess.^{1,6} Lohan et al¹ estimated that dropped stones produce an intra-abdominal abscess in 0.6% to 2.9% of cases of dropped stones and bile spillage, based on reports by Rice et al⁴ and Morrin et al.⁷ Dropped stones should be recognized as a potential cause of intra-abdominal abscess in any cholecystectomy patient months or even years after the surgery. Also, these abscesses are not necessarily confined to the right upper quadrant: they can occur anywhere in the abdominal cavity.^{5,7}

Given the ever-increasing popularity of laparoscopic cholecystectomy, the problem of intra-abdominal abscess due to dropped gallstones will only become a more common problem. Early diagnosis is the key to avoiding long and unnecessary treatment.

If dropped gallstones do become infected and eventually cause symptoms, they may require surgical or percutaneous removal in conjunction with antimicrobial therapy.⁸

**Infected
dropped
gallstones may
need surgical
removal plus
antibiotics**

REFERENCES

1. **Lohan D, Walsh S, McLoughlin R, Murphy J.** Imaging of the complications of laparoscopic cholecystectomy. *Eur Radiol* 2005; 15:904–912.
2. **Casillas S, Kittur DS.** Late abscess formation after spilled gallstones masquerading as a liver mass. *Surg Endosc* 2003; 17:833.
3. **Tumer AR, Yuksek YN, Yasti AC, Gozalan U, Kama NA.** Dropped gallstones during laparoscopic cholecystectomy: the consequences. *World J Surg* 2005; 29:437–440.
4. **Rice DC, Memon MA, Jamison RL, et al.** Long-term consequences of intraoperative spillage of bile and gallstones during laparoscopic cholecystectomy. *J Gastrointest Surg* 1997; 1:85–91.
5. **Sathesh-Kumar T, Saklani AP, Vinayagam R, Blackett RL.** Spilled gall stones during laparoscopic cholecystectomy: a review of the literature. *Postgrad Med J* 2004; 80:77–79.
6. **Horton M, Florence MG.** Unusual abscess patterns following dropped gallstones during laparoscopic cholecystectomy. *Am J Surg* 1998; 175:375–379.
7. **Morrin MM, Kruskal JB, Hochman MG, Saldinger PF, Kane RA.** Radiologic features of complications arising from dropped gallstones in laparoscopic cholecystectomy patients. *AJR Am J Roentgenol* 2000; 174:1441–1445.
8. **Akyar G, Aytac S, Yagci C, Akyar S.** Abscess formation due to dropped gallstone after laparoscopic cholecystectomy. *Eur J Radiol* 1997; 25:242–245.

ADDRESS: Steven Gordon, MD, Department of Infectious Diseases, S32, Cleveland Clinic, 9500 Euclid Avenue, Cleveland, OH 44195; e-mail gordons@ccf.org.



BRIEF ANSWERS
TO SPECIFIC
CLINICAL QUESTIONS

What questions do you want answered?

We want to know what questions you want addressed in “1-Minute Consult.”
All questions should be on practical, clinical topics.

PLEASE PRINT CLEARLY

Q: _____

NAME _____
 ADDRESS _____
 CITY _____
 STATE _____ ZIP _____
 PHONE _____ EMAIL _____