JOHAN DENOLLET, PhD

CoRPS–Center of Research on Psychology in Somatic diseases, Tilburg University, Tilburg, The Netherlands; Department of Cardiology, Antwerp University Hospital, Antwerp, Belgium

VIVIANE M. CONRAADS, MD, PhD

Department of Cardiology, Antwerp University Hospital, Antwerp, Belgium

Type D personality and vulnerability to adverse outcomes in heart disease

ABSTRACT

General distress, shared across depression, anxiety and anger, partly accounts for the link between mind and heart. The type D (distressed) personality profile identifies individuals who are particularly vulnerable to the adverse effect of general distress. Type D individuals frequently experience negative emotions and are socially inhibited. This profile is more stable than that associated with episodes of clinical depression and describes the chronic nature of distress in some patients. Type D may also partly account for the effect of emotional distress on cardiac prognosis. Type D is associated with a threefold increased risk of adverse cardiovascular outcomes, even after adjustment for depression. This relationship is less obvious in patients with heart failure. Plausible pathways linking type D to cardiovascular complications include hypothalamic-pituitary-adrenal-axis hyperreactivity, autonomic and inflammatory dysregulation, and increased oxidative stress. Research needs to further clarify these pathways and investigate whether type D patients may benefit from closer monitoring of risk factors and a personalized approach to behavioral intervention. The DS14 is a brief, well-validated measure of type D that could be incorporated into clinical research and practice to identify high-risk patients.

epression has been studied extensively in relation to cardiovascular disease.¹⁻³ In addition to depression, anger⁴ and anxiety⁵ also may promote coronary artery disease (CAD), suggesting that emotional distress in general may be related to increased cardiovascular risk. Evidence indicates that the general distress shared across depression, anger, and anxiety predicts CAD, even after controlling for each of these specific negative emotions.⁶

doi:10.3949/ccjm.78.s1.02

THE CONCEPT OF TYPE D PERSONALITY

Lately, there is a renewed interest in broad individual differences in general distress and heart disease.⁷ Since psychologic factors often cluster together in individual patients, biobehavioral research may benefit from the identification of discrete personality sub-types.⁸ This focus on the identification of psychologically vulnerable patients who are at increased risk for adverse outcomes has led to the introduction of the *distressed*⁹ or *type D*¹⁰ personality profile in cardiovas-cular research. This personality construct is defined as follows:

"The type D (distressed) personality profile refers to a general propensity to psychological distress that is characterized by the combination of negative affectivity and social inhibition."¹⁰

Negative affectivity, or the tendency to experience negative emotions across time and situations, is a major determinant of emotional distress in cardiac patients.^{9,10} Patients who score high on this trait frequently report feelings of dysphoria, worry, and tension. Social inhibition, or the tendency to inhibit the expression of emotions or behavior, is a major determinant of social distress.^{9,10} Patients who score high on this trait tend to avoid negative reactions from others.

Both traits define psychologically vulnerable patients and can be assessed with the type D scale (DS14).¹⁰ This brief measure consists of a seven-item negative affectivity subscale (eg, *I often feel unhappy*) and a seven-item inhibition subscale (eg, *I am inhibited in social interactions*), and has a clear two-factor structure and good reliability (Cronbach's α = .88 and .86). Patients are classified as type D if they score 10 or higher on both DS14 subscales.¹⁰ The prevalence of type D personality ranges between 20% and 40% across different types of cardiovascular conditions.

The type D construct was designed for the early identification of chronically distressed patients. This article reviews (1) the risk of adverse events associated with type D, (2) the extent to which type D is

CLEVELAND CLINIC JOURNAL OF MEDICINE VOLUME 78 \bullet SUPPLEMENT 1 AUGUST 2011 S13

Both authors reported that they have no financial relationships that pose a potential conflict of interest with this article.

This work was supported by the Netherlands Organization for Scientific Research (The Hague, The Netherlands) with a VICI grant (453-04-004) to Dr. Johan Denollet.

TABLE 1

Type [) and	risk	of	clinical	events	in	cardiovascula	r disease	patients
.,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,			•••	enneon.	0101100		call allo i ab calla	ansease	particites

Cardiovascular disease (n)	Clinical event (follow-up)	OR/HR (95% CI)	Meta-analytic review ²³
CAD			
CAD (303) ¹¹	Total mortality (6–10 y)	OR = 4.1 (1.9–8.8)*	Included in meta-analysis
CAD (319) ¹²	Cardiac death, MI (5 y)	OR = 8.9 (3.2–24.7) ⁺	Included in meta-analysis
CAD (337) ¹³	Total mortality, MI (5 y)	OR = 4.8 (1.4–16.5)*	Included in meta-analysis
CAD (875) ¹⁴	Total mortality, MI (9 mo)	OR = 5.3 (2.0–13.6) ⁺	Included in meta-analysis
CAD (358) ¹⁵	Total mortality, MI (2 y)	HR = 2.6 (1.1–6.0) [±]	Included in meta-analysis
CAD (473) ¹⁶	Cardiac death, MI (1.8 y)	HR = 2.2 (1.1–4.3) [±]	Not included in meta-analysis
Other			
PAD (184) ¹⁷	Total mortality (4 y)	HR = 3.5 (1.1–11.1) [±]	Included in meta-analysis
CHF (87) ¹⁸	Cardiac death, MI (6–10 y)	OR = 4.7 (1.9–11.8)*	Included in meta-analysis
CHF/HT (51) ¹⁹	Mortality, rejection (5.4 y)	OR = 6.8 (1.4–30.9) [±]	Included in meta-analysis
CHF (641) ²⁰	Cardiac death (3.1 y)	HR = 1.2 (0.6–2.1)	Not included in meta-analysis
ICD (391) ²¹	Ventricular arrhythmia (1 y)	HR = 1.9 (1.1–3.1) [±]	Not included in meta-analysis
ICD (371) ²²	Total mortality (1.7 y)	HR = 2.8 (1.2–6.2)*	Not included in meta-analysis

* *P* < .01; [†]*P* < .0001; [‡]*P* < .05

CAD = coronary artery disease; CHF = chronic heart failure; HR = hazard ratio; HT = heart transplantation; ICD = implantable cardioverter defibrillator; MI = myocardial infarction; OR = odds ratio; PAD = peripheral arterial disease

distinct from depression, (3) the biologic pathways of type D, and (4) the implications of the type D personality profile.

RISK ASSOCIATED WITH TYPE D

Several prospective studies from our group have examined the notion that type D patients are particularly vulnerable to adverse events **(Table 1)**. In patients with CAD, evidence indicates that type D personality is an independent predictor of adverse events, including (cardiac) death, myocardial infarction, and need for revascularization procedures.^{11–16} In these studies, type D also emerged as an independent predictor of adverse events after adjustment for anxiety,¹¹ stress,¹³ depression,¹⁶ disease severity,^{11–16} and type of invasive treatment.¹⁴ This increased risk associated with the type D profile was observed in the broader group of patients with CAD,^{11–15} as well as in patients who survived an initial myocardial infarction.¹⁶

The relationship between type D personality and adverse events has also been investigated in other cardiovascular conditions. Type D has been associated with poor prognosis in patients with peripheral arterial disease,¹⁷ but evidence for the prognostic role of type D in patients with chronic heart failure is mixed. In a study of patients with heart failure following myocardial infarction, type D predicted cardiac death independent of disease severity¹⁸; in a study of heart failure patients who underwent cardiac transplantation, type D was associated with early allograft rejection and increased mortality.¹⁹ However, type D was not associated with cardiac death in a recent, larger heart failure study.²⁰ The link between psychologic factors and heart failure is complex³ and may be less obvious than the type D-CAD link.²⁰ Type D has also been associated with the occurrence of life-threatening arrhythmias following implantable cardioverter defibrillator (ICD) treatment,²¹ and it has been shown to predict an increased risk for mortality in ICD patients, independent from shocks and disease severity.²²

The wide range in odds ratios and confidence intervals indicates disparity in data across these type D studies (**Table 1**). We recently performed a metaanalysis of prospective studies between 1996 and 2009 to provide a more reliable estimate of the risk associated with type D. In this analysis, type D was associated with a threefold increased risk of adverse events²³; the confidence interval of this pooled odds ratio ranged

S14 CLEVELAND CLINIC JOURNAL OF MEDICINE VOLUME 78 • SUPPLEMENT 1 AUGUST 2011

from 2.7 to 5.1. In addition, type D personality was associated with a threefold increased risk (range, 2.6 to 4.3) of emotional distress over time.²³ From the recent studies that were not included in this meta-analysis, one reported negative findings²⁰ and three others positive findings^{16,21,22} on the risk associated with type D.

COMPARING DEPRESSION AND TYPE D

Many studies report on depression and cardiac disease,¹⁻³ but both conceptual differences and clinical evidence indicate that type D and depression are distinct forms of distress (**Table 2**). Conceptually, type D focuses not only on depressive affect but also on the general distress shared across negative emotions,¹⁰ and it is based on the notion that social inhibition modulates the effect of negative emotions on cardiac prognosis.²⁴ While depression refers to an episodic distress factor (patients may go in and out of depressive episodes), the type D construct focuses on an underlying factor that predisposes patients to more chronic forms of distress.⁸

Clinical evidence shows that, after adjustment for depression, type D remained a predictor of adverse cardiac events in CAD.^{16,24,25} Following ICD implantation, anxious type D patients were at risk of ventricular arrhythmias, whereas depression did not predict arrhythmias.²¹ Type D also exerts an adverse effect on patients' health status following coronary bypass surgery,²⁶ heart failure,²⁷ or myocardial infarction,²⁸ adjusting for depressive symptoms. Type D is related to biomarkers of increased stress levels independent of depression^{29–31} and, unlike depression, type D is not confounded by the severity of cardiac disorder.³²

Following myocardial infarction, only one of four distressed patients met criteria for both type D and depression; most had one form of distress but not the other.³² Research in healthy³³ and in cardiac³⁴ populations confirmed that items from depression and type D scales reflect different distress factors. After adjustment for depression at baseline, type D also predicted the incidence,³⁵ persistence,³⁶ and severity^{37,38} of depression and anxiety. However, these findings do not imply that depression and type D are antonymous perspectives or that one perspective is better than the other in predicting outcomes; rather, we would like to argue that both constructs represent complementary perspectives that have added value.²³

BIOLOGIC PATHWAYS OF TYPE D

A number of biologic pathways have been suggested to explain the effect of type D (Table 3). Some have suggested dysregulation of the hypothalamic-

TABLE 2

Type D and depression are different forms of distress in cardiovascular disease patients

	Conceptual differences	
Emotional	Type D focuses on general distress shared across negative emotions (anxiety, irritability, and others ¹⁰) in addition to depressive affect	
Social	Social inhibition is a factor in type D that may moderate the expression of emotions and behaviors in social interaction ²⁴	
Duration	Emotional and social distress is a chronic factor (\geq 2 years) in type D, whereas it is an episodic factor (< 2 years) in depression ^{8–10}	
	Cardiovascular outcomes	
Clinical events	Type D personality predicts mortality and other clinical events in cardiac patients, even after adjustment for severity ¹⁶ and symptoms ^{24,25} of depression	
Health status	Type D personality independently predicts poor health status over time in cardiac patients, above and beyond symptoms of depression ^{26–28}	
Pathways of disease	Type D personality predicts increased oxida- tive stress and cortisol levels in cardiac patients after adjustment for depressive symptoms ²⁹⁻³¹	
	Psychologic outcomes	
Distinct diagnosis	There is only limited overlap between type D and depression classification ^{25,32} ; items in type D and depression scales reflect different distress factors	
Depressive symptoms	Type D personality predicts the onset and persistence of depressive symptoms in cardiac patients, controlling for depression at baseline ^{35,36}	
Anxiety symptoms	Type D personality predicts the occurrence and severity of anxiety in cardiac patients, above and beyond symptoms of depression ^{37,38}	

pituitary-adrenal axis in patients with type D personality.³⁹ In fact, type D has been associated with greater cortisol reactivity to stress in healthy individuals⁴⁰ and with higher awakening³⁰ and daytime³¹ cortisol levels in CAD patients. Autonomic dysregulation can also be inferred in type D individuals on the basis of a higher resting heart rate⁴¹ and cardiovascular hyperreactivity^{40,42} and decreased heart rate variability⁴³ in response to stress. In addition, type D has been related to reduced heart rate recovery after

AUGUST 2011

S15

CLEVELAND CLINIC JOURNAL OF MEDICINE VOLUME 78 • SUPPLEMENT 1

TABLE 3

Potential biologic mechanisms underlying type D

	Healthy individuals	Cardiovascular patients
HPA-axis dysregulation	Increased cortisol reactivity to stress ⁴⁰	Higher CAR ³⁰ ; higher daytime cortisol ³¹
Autonomic dysregulation	Higher HR ⁴¹ ; increased CV stress reactivity ^{40,42} ; decreased HRV ⁴³	Reduced HR recovery after exercise ⁴⁴
Inflammatory dysregulation	Higher concentration of CRP ⁴¹	Increased plasma levels of TNF-α, TNFR1, TNFR2 ^{46,47}
Reduced number of stem cells	Decreased EPC counts associated with NA ⁴⁸	Decreased EPC counts associated with type D ⁴⁹
Increased oxidative stress		Lower levels of Hsp70 and higher levels of XO ²⁹

CAR = cortisol awakening response; CRP = C-reactive protein; CV = cardio-vascular; EPC = bone marrow-derived endothelial progenitor cells; HPA = hypothalamic-pituitary-adrenal; HR = heart rate; HRV = heart rate variability; Hsp70 = heat shock protein 70; NA = negative affectivity; TNF- α = tumor necrosis factor- α ; TNFR1 = soluble TNF- α receptor 1; TNFR2 = soluble TNF- α receptor 2; XO = xanthine oxidase

exercise in patients with heart failure.⁴⁴ These indices of excessive sympathetic or inadequate parasympathetic modulation of heart rate predict poor cardiac prognoses.⁴⁵

Other studies found that type D was associated with inflammatory dysregulation. In healthy adults, type D has been related to higher concentrations of C-reactive protein.⁴¹ In heart failure patients, type D is associated with increased plasma levels of the proinflammatory cytokine tumor necrosis factor (TNF)-α and its soluble receptors 1 and 2.46,47 Increased TNF- α levels may cause suppression of bone-marrow-derived endothelial progenitor cells (EPCs) that play an important role in maintaining vascular integrity. The negative affectivity component of type D has been shown to predict decreased circulating EPC counts in healthy individuals⁴⁸; another study found that these EPC numbers were reduced by more than 50% in heart failure patients with a type D personality.⁴⁹ Type D personality is also associated with an increased oxidative stress burden in patients with chronic heart failure.²⁹ Studies on genetic linkage⁵⁰ and heritability⁵¹ further support biologic underpinnings of the type D construct.

Regarding pathways that may explain the effect

of type D, some issues are of special interest. First, genetic factors contribute to stability in type D personality, but environmental factors may induce changes in type D characteristics over time.⁵¹ Hence, given this role of environmental influences over time, behavioral intervention would be feasible and useful in type D patients. Second, type D can promote heart disease indirectly through behavioral pathways. Type D has been associated with a sedentary lifestyle,^{41,52} an unhealthy diet,⁵³ and a passive coping style.^{54,55} Poor adherence to medical treatment^{56,57} and reluctance to consult clinical staff⁵⁸ may jeopardize the working relationship with type D patients in clinical care. Intervention may focus on the management of these behavioral risk factors in type D patients. Third, many of these biologic^{40-43,48,50,51} and behavioral^{41,52-54} pathways have also been documented in healthy type D individuals, which suggests that these associations cannot be explained away by the confounding effect of underlying cardiovascular disease.

CLINICAL IMPLICATIONS OF TYPE D

The findings from type D research have a number of clinical implications. Type D is associated with an increased risk of adverse events,²³ chronic distress,^{35–38} and suicidal ideation.⁵⁹ Type D may also have an adverse effect on the outcome of invasive treatment.^{14,19,21,22,24,26,60}

Type D was associated with mortality and morbidity at 9 months¹⁴ and 2 years²⁴ following coronary artery stenting, and with impaired health status 1 year following bypass surgery.²⁶ Type D also predicted mortality and allograft rejection following heart transplantation,¹⁹ and an increased risk of ventricular arrhythmia²¹ and mortality²² in ICD patients. Researchers from the Cleveland Clinic have shown that type D is a risk factor for anxiety in ICD patients.⁶⁰

Regarding the DSM-IV classification by the American Psychiatric Association,⁶¹ type D qualifies for the diagnosis "psychological factors affecting medical condition" (Section 316). In keeping with this classification, the diagnostic category type D affects (1) the *course* of cardiovascular conditions,²³ (2) the *treatment* of these conditions,^{56,57} and (3) the *working relationship* with medical staff.⁵⁸ At present, no clinical trial has examined whether intervention for distress among type D patients alters their risk for adverse events. Nevertheless, some have argued that it is plausible for type D patients to learn new strategies to reduce their level of general distress.⁶² Previous research with patients suggests that psychotherapy,

S16 CLEVELAND CLINIC JOURNAL OF MEDICINE VOLUME 78 • SUPPLEMENT 1 AUGUST 2011

social skills training, stress management, and relaxation training may reduce stress in these patients and improve their ability to express their emotions to others.⁶² Others have suggested that stress management training, including communication skills and problem-solving, may further improve the risk profile and health in cardiac patients.⁶³

It is possible that type D patients may benefit from close monitoring of their clinical condition and from aggressive management of their risk factor profile to prevent adverse clinical events. Cardiac rehabilitation is an effective approach to treating risk factors and enhancing well-being in CAD.^{63,64} Ă few studies have examined the effect of cardiac rehabilitation in type D patients. One study found a significant decrease in the social inhibition component of type D following cardiac rehabilitation, but there was no change in the prevalence of type D at 1-year followup.65 Although the type D profile tends to remain stable during rehabilitation,^{65,66} evidence shows that type D patients who participate in cardiac rehabilitation improve in physical and mental health status.⁶⁶ Cardiac rehabilitation may also ward off further deterioration in negative affect,⁶⁷ which, in turn, has been associated with better survival in patients who participated in rehabilitation.68 Future studies need to examine the effect of cardiac rehabilitation and other personalized approaches to treatment in type D patients.

CONCLUSIONS

General distress shared across negative emotions^{6,23} may partly account for the role of depression, anxiety, and anger in cardiovascular disorders.^{1–5} Some cardiac patients are more likely to experience distress than others. Type D may identify these psychologically vulnerable patients who tend to experience general distress.²³ This propensity to general distress differs from depression, predicts adverse outcomes, is linked to plausible biologic pathways, and highlights the chronic nature of psychologic distress in some cardiac patients.

After adjustment for depression, type D remains significantly associated with an increased risk of adverse events in patients with CAD.^{16,24,25} However, this association is less obvious in patients with heart failure, and type D did not predict survival in one heart failure study.²⁰ Although initial findings suggest a number of plausible biologic and behavioral pathways, more research is needed to explain the adverse effect of type D on cardiovascular outcomes. Future research also needs to investigate whether type D patients may benefit from close monitoring of their risk factors and a more personalized approach to behavioral and cardiac treatment.

Overall, the current understanding of type D indicates that general distress should not be ignored in the link between mind and heart, and that cardiovascular patients who have a type D personality profile are particularly vulnerable to the adverse clinical effects of general distress. The DS14¹⁰ is a brief, well-validated measure of type D that could be incorporated into clinical research and practice to identify patients who are at risk of chronic distress and poor prognosis.

REFERENCES

- Pozuelo L, Zhang J, Franco K, Tesar G, Penn M, Jiang W. Depression and heart disease: what do we know, and where are we headed? Cleve Clin J Med 2009; 76:59–70.
- Davidson KW, Korin MR. Depression and cardiovascular disease: selected findings, controversies, and clinical implications from 2009. Cleve Clin J Med 2010; 77(suppl 3):S20–S26.
- Kop WJ, Synowski SJ, Gottlieb SS. Depression in heart failure: biobehavioral mechanisms. Heart Failure Clin 2011; 7:23–38.
- **4. Chida Y, Steptoe A.** The association of anger and hostility with future coronary heart disease: a meta-analytic review of prospective evidence. J Am Coll Cardiol 2009; 53:936–946.
- Roest AM, Martens EJ, de Jonge P, Denollet J. Anxiety and risk of incident coronary heart disease: a meta-analysis. J Am Coll Cardiol 2010; 56:38–46.
- 6. Kubzansky LD, Cole SR, Kawachi I, Vokonas P, Sparrow D. Shared and unique contributions of anger, anxiety, and depression to coronary heart disease: a prospective study in the normative aging study. Ann Behav Med 2006; 31:21–29.
- Steptoe A, Molloy GJ. Personality and heart disease. Heart 2007; 93:783–784.
- Denollet J. Biobehavioral research on coronary heart disease: where is the person? J Behav Med 1993; 16:115–141.
- Denollet J, Sys SU, Brutsaert DL. Personality and mortality after myocardial infarction. Psychosom Med 1995; 57:582–591.
- Denollet J. DS14: standard assessment of negative affectivity, social inhibition, and Type D personality. Psychosom Med 2005; 67:89–97.
- Denollet J, Sys SU, Stoobant N, Rombouts H, Gillebert TC, Brutsaert DL. Personality as independent predictor of long-term mortality in patients with coronary heart disease. Lancet 1996; 347:417–421.
- 12. Denollet J, Vaes J, Brutsaert DL. Inadequate response to treatment in coronary heart disease: adverse effects of type D personality and younger age on 5-year prognosis and quality of life. Circulation 2000; 102:630–635.
- Denollet J, Pedersen SS, Vrints CJ, Conraads VM. Usefulness of type D personality in predicting five-year cardiac events above and beyond concurrent symptoms of stress in patients with coronary heart disease. Am J Cardiol 2006; 97:970–973.
- 14. Pedersen SS, Lemos PA, van Vooren PR, et al. Type D personality predicts death or myocardial infarction after bare metal stent or sirolimus-eluting stent implantation: a Rapamycin-Eluting Stent Evaluated at Rotterdam Cardiology Hospital (RESEARCH) registry substudy. J Am Coll Cardiol 2004; 44:997–1001.
- Pedersen SS, Denollet J, Ong AT, et al. Adverse clinical events in patients treated with sirolimus-eluting stents: the impact of Type D personality. Eur J Cardiovasc Prev Rehabil 2007; 14:135–140.
- Martens EJ, Mols F, Burg MM, Denollet J. Type D personality predicts clinical events after myocardial infarction, above and beyond disease severity and depression. J Clin Psychiatry 2010; 71:778–783.
- 17. Aquarius AE, Smolderen KG, Hamming JF, De Vries J, Vriens

CLEVELAND CLINIC JOURNAL OF MEDICINE

PW, Denollet J. Type D personality and mortality in peripheral arterial disease: a pilot study. Arch Surg 2009; 144:728–733.

- Denollet J, Brutsaert DL. Personality, disease severity, and the risk of long-term cardiac events in patients with a decreased ejection fraction after myocardial infarction. Circulation 1998; 97:167–173.
- Denollet J, Holmes RV, Vrints CJ, Conraads VM. Unfavorable outcome of heart transplantation in recipients with type D personality. J Heart Lung Transplant 2007; 26:152–158.
- Pelle AJ, Pedersen SS, Schiffer AA, Szabó BM, Widdershoven JW, Denollet J. Psychological distress and mortality in systolic heart failure. Circ Heart Fail 2010; 3:261–267.
- van den Broek KC, Nyklíček I, van der Voort PH, Alings M, Meijer A, Denollet J. Risk of ventricular arrhythmia after implantable defibrillator treatment in anxious type D patients. J Am Coll Cardiol 2009; 54:531–537.
- 22. Pedersen SS, van den Broek KC, Erdman RA, Jordaens L, Theuns DA. Pre-implantation implantable cardioverter defibrillator concerns and Type D personality increase the risk of mortality in patients with an implantable cardioverter defibrillator. Europace 2010; 12:1446–1452.
- Denollet J, Schiffer AA, Spek V. A general propensity to psychological distress affects cardiovascular outcomes: evidence from research on the type D (distressed) personality profile. Circ Cardiovasc Qual Outcomes 2010; 3:546–557.
- Denollet J, Pedersen SS, Ong AT, Erdman RA, Serruys PW, van Domburg RT. Social inhibition modulates the effect of negative emotions on cardiac prognosis following percutaneous coronary intervention in the drug-eluting stent era. Eur Heart J 2006; 27:171–177.
- Denollet J, Pedersen SS. Prognostic value of Type D personality compared with depressive symptoms. Arch Intern Med 2008; 168:431–432.
- Al-Ruzzeh S, Athanasiou T, Mangoush O, et al. Predictors of poor mid-term health related quality of life after primary isolated coronary artery bypass grafting surgery. Heart 2005; 91:1557–1562.
- 27. Schiffer AA, Pedersen SS, Widdershoven JW, Denollet J. Type D personality and depressive symptoms are independent predictors of impaired health status in chronic heart failure. Eur J Heart Fail 2008; 10:802–810.
- Mols F, Martens EJ, Denollet J. Type D personality and depressive symptoms are independent predictors of impaired health status following acute myocardial infarction. Heart 2010; 96:30–35.
- Kupper N, Gidron Y, Winter J, Denollet J. Association between type D personality, depression, and oxidative stress in patients with chronic heart failure. Psychosom Med 2009; 71:973–980.
- Whitehead DL, Perkins-Porras L, Strike PC, Magid K, Steptoe A. Cortisol awakening response is elevated in acute coronary syndrome patients with type-D personality. J Psychosom Res 2007; 62:419–425.
- Molloy GJ, Perkins-Porras L, Strike PC, Steptoe A. Type-D personality and cortisol in survivors of acute coronary syndrome. Psychosom Med 2008; 70:863–868.
- Denollet J, de Jonge P, Kuyper A, et al. Depression and Type D personality represent different forms of distress in the Myocardial INfarction and Depression–Intervention Trial (MIND-IT). Psychol Med 2009; 39:749–756.
- 33. Kudielka BM, von Känel R, Gander ML, Fischer JE. The interrelationship of psychosocial risk factors for coronary artery disease in a working population: do we measure distinct or overlapping psychological concepts? Behav Med 2004; 30:35–43.
- Pelle AJ, Denollet J, Zwisler AD, Pedersen SS. Overlap and distinctiveness of psychological risk factors in patients with ischemic heart disease and chronic heart failure: are we there yet? J Affect Disord 2009; 113:150–156.
- 35. Pedersen SS, Ong AT, Sonnenschein K, Serruys PW, Erdman RA, van Domburg RT. Type D personality and diabetes predict the onset of depressive symptoms in patients after percutaneous coronary intervention. Am Heart J 2006; 151:367.e1–367.e6.
- 36. Martens EJ, Smith OR, Winter J, Denollet J, Pedersen SS. Cardiac history, prior depression and personality predict course

of depressive symptoms after myocardial infarction. Psychol Med 2008; 38:257–264.

- van Gestel YR, Pedersen SS, van de Sande M, et al. Type-D personality and depressive symptoms predict anxiety 12 months post-percutaneous coronary intervention. J Affect Disord 2007; 103:197–203.
- Schiffer AA, Pedersen SS, Broers H, Widdershoven JW, Denollet J. Type-D personality but not depression predicts severity of anxiety in heart failure patients at 1-year follow-up. J Affect Disord 2008; 106:73–81.
- Sher L. Type D personality: the heart, stress, and cortisol. QJM 2005; 98:323–329.
- Habra ME, Linden W, Anderson JC, Weinberg J. Type D personality is related to cardiovascular and neuroendocrine reactivity to acute stress. J Psychosom Res 2003; 55:235–245.
- Einvik G, Dammen T, Hrubos-Strøm H, et al. Prevalence of cardiovascular risk factors and concentration of C-reactive protein in type D personality persons without cardiovascular disease [published online ahead of print February 9, 2011]. Eur J Cardiovasc Prev Rehabil. PMID: 21450648.
- Williams L, O'Carroll RE, O'Connor RC. Type D personality and cardiac output in response to stress. Psychol Health 2009; 24:489–500.
- Martin LA, Doster JA, Critelli JW, et al. Ethnicity and Type D personality as predictors of heart rate variability. Int J Psychophysiol 2010; 76:118–121.
- 44. von Känel R, Barth J, Kohls S, et al. Heart rate recovery after exercise in chronic heart failure: role of vital exhaustion and type D personality. J Cardiol 2009; 53:248–256.
- Carney RM, Freedland KE. Depression and heart rate variability in patients with coronary heart disease. Cleve Clin J Med 2009; 76(suppl 2):S13–S17.
- 46. DenoÎlet J, Vrints CJ, Conraads VM. Comparing Type D personality and older age as correlates of tumor necrosis factor-α dysregulation in chronic heart failure. Brain Behav Immun 2008; 22:736–743.
- Denollet J, Schiffer AA, Kwaijtaal M, et al. Usefulness of Type D personality and kidney dysfunction as predictors of interpatient variability in inflammatory activation in chronic heart failure. Am J Cardiol 2009; 103:399–404.
- Fischer JC, Kudielka BM, von Känel R, Siegrist J, Thayer JF, Fischer JE. Bone-marrow derived progenitor cells are associated with psychosocial determinants of health after controlling for classical biological and behavioral cardiovascular risk factors. Brain Behav Immun 2009; 23:419–426.
- Van Craenenbroeck EM, Denollet J, Paelinck BP, et al. Circulating CD34+/KDR+ endothelial progenitor cells are reduced in chronic heart failure patients as a function of Type D personality. Clin Sci 2009; 117:165–172.
- Ladwig K-H, Emeny RT, Gieger C, et al. Single nucleotide polymorphism associations with type-D personality in the general population: findings from the KORA K-500-substudy. Psychosom Med 2009; 71:A-28. Abstract 1781.
- Kupper N, Boomsma DI, de Geus EJ, Denollet J, Willemsen G. Nine-year stability of type D personality: contributions of genes and environment. Psychosom Med 2011; 73:75–82.
- 52. Hausteiner C, Klupsch D, Emeny R, Baumert J, Ladwig KH; for the KORA Investigators. Clustering of negative affectivity and social inhibition in the community: prevalence of type D personality as a cardiovascular risk marker. Psychosom Med 2010; 72:163–171.
- Williams L, O'Connor RC, Howard S, et al. Type-D personality mechanisms of effect: the role of health-related behavior and social support. J Psychosom Res 2008; 64:63–69.
- Polman R, Borkoles E, Nicholls AR. Type D personality, stress, and symptoms of burnout: the influence of avoidance coping and social support. Br J Health Psychol 2010; 15:681–696.
- 55. Yu X-N, Chen Z, Zhang J, Liu X. Coping mediates the association between Type D personality and perceived health in Chinese patients with coronary heart disease. Int J Behav Med 2010; Oct 13 [Epub ahead of print].
- 56. Broström A, Strömberg A, Mårtensson J, Ulander M, Harder L,

S18 CLEVELAND CLINIC JOURNAL OF MEDICINE VOLUME 78 • SUPPLEMENT 1 AUGUST 2011

Svanborg E. Association of Type D personality to perceived side effects and adherence in CPAP-treated patients with OSAS. J Sleep Res 2007; 16:439–447.

- Williams L, O'Connor RC, Grubb N, O'Carroll R. Type D personality predicts poor medication adherence in myocardial infarction patients [published online ahead of print March 3, 2011]. Psychol Health. PMID: 21391133.
- Schiffer AA, Denollet J, Widdershoven JW, Hendriks EH, Smith OR. Failure to consult for symptoms of heart failure in patients with a type-D personality. Heart 2007; 93:814–818.
- Michal M, Wiltink J, Till Y, et al. Type D personality and depersonalization are associated with suicidal ideation in the German general population aged 35–74: results from the Gutenberg Heart Study. J Affect Disord 2010; 125:227–233.
- Pozuelo L, Panko M, Ching B, et al. Prevalence of anxiety and type-D personality in an outpatient ICD clinic. Circulation 2009; 120:S493–S494. Abstract 1385.
- American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders. Fourth Edition. Washington, DC: American Psychiatric Association, 2000.
- 62. Tulloch H, Pelletier R. Does personality matter after all? Type D personality and its implications for cardiovascular prevention and rehabilitation. Curr Issues Card Rehab Prevention 2008; 16:2–4.
- 63. Blumenthal JA, Wang JT, Babyak M, et al. Enhancing standard

cardiac rehabilitation with stress management training: background, methods, and design for the enhanced study. J Cardiopulm Rehabil Prev 2010; 30:77–84.

- Denollet J. Sensitivity of outcome assessment in cardiac rehabilitation. J Consult Clin Psychol 1993; 61:686–695.
- 65. Karlsson MR, Edström-Plüss C, Held C, Henriksson P, Billing E, Wallén NH. Effects of expanded cardiac rehabilitation on psychosocial status in coronary artery disease with focus on type D characteristics. J Behav Med 2007; 30:253–261.
- 66. Pelle AJ, Erdman RA, van Domburg RT, Spiering M, Kazemier M, Pedersen SS. Type D patients report poorer health status prior to and after cardiac rehabilitation compared to non-type D patients. Ann Behav Med 2008; 36:167–175.
- Denollet J, Brutsaert DL. Enhancing emotional well-being by comprehensive rehabilitation in patients with coronary heart disease. Eur Heart J 1995; 16:1070–1078.
- Denollet J, Brutsaert DL. Reducing emotional distress improves prognosis in coronary heart disease: 9-year mortality in a clinical trial of rehabilitation. Circulation 2001; 104:2018–2023.

Correspondence: Johan Denollet, PhD, CoRPS, Department of Medical Psychology and Neuropsychology, Tilburg University, P.O. Box 90153, 5000 LE Tilburg, The Netherlands; denollet@uvt.nl