

Q: Can we reduce the risk of readmission for a patient with an exacerbation of COPD?

MIRNA B. AYACHE, MD, MPH, FACP

Department of Hospital Medicine, Cleveland Clinic

SHAFIK BOYAJI, MD

Department of Hospital Medicine, Cleveland Clinic

JAMES C. PILE, MD

Department of Hospital Medicine, Cleveland Clinic; Associate Professor, Cleveland Clinic Lerner College of Medicine of Case Western Reserve University, Cleveland, OH; Deputy Editor, *Cleveland Clinic Journal of Medicine*

A: We think so. Some strategies to reduce readmission rates, such as coordinating care and managing comorbidities, apply to chronic diseases in general, while others are disease-specific. To reduce the need for hospital readmission for chronic obstructive pulmonary disease (COPD), coordinated efforts involving both inpatient and outpatient care are necessary. This can be achieved by using a checklist before discharge (TABLE 1) and by implementing outpatient COPD programs that continue patient education and provide rapid access to medical support if needed.

See related commentary, page 528

There is room for improvement. COPD is common and expensive, with high rates of hospital readmission,¹ and up to 70% of the money we spend on it goes for hospital care.² No wonder then that the Centers for Medicare and Medicaid Services has now expanded its Readmissions Reduction Program to include acute COPD exacerbations.³ Yet in a retrospective study, Yip et al⁴ found that fewer than half of patients hospitalized with acute exacerbation of COPD received appropriate vaccinations, counseling on smoking cessation, and long-acting inhalers—all of which are on our checklist.⁴

The following interventions have been demonstrated to be useful in reducing COPD hospital admissions and the risk of death.

doi:10.3949/ccjm.81a.13139

TABLE 1

Checklist for COPD patients before they leave the hospital

Counseling on smoking cessation if the patient is a smoker or is at risk of starting smoking again

Appropriate influenza and pneumonia vaccinations

Long-acting bronchodilators: an anticholinergic or a beta-agonist, or both (except GOLD group A patients, who have few symptoms and are at a low risk of exacerbations); inhaled steroids for those with previous exacerbations who are at high risk of relapse

Long-term oxygen therapy, if indicated, ie, in those with chronic resting hypoxemia (room air PaO₂ ≤ 55 mm Hg or ≤ 59 mm Hg with signs of right-sided heart strain or polycythemia)

Home exercise program or pulmonary rehabilitation referral after an exacerbation

Patient education regarding symptom monitoring and in-haler therapy, including proper inhaler technique (see http://my.clevelandclinic.org/disorders/Chronic_Obstructive_Pulmonary_Disease/hic_Understanding_COPD.aspx)

Consider home noninvasive ventilator support in select patients with recurrent hospitalization for acidotic exacerbation of COPD if the patient qualifies under Centers for Medicare and Medicaid Services guidelines,²⁰ ie:

- Paco₂ ≥ 52 mm Hg and
- Evidence of nocturnal hypoventilation based on nocturnal oximetry showing sustained desaturation to < 89% for ≥ 5 min on oxygen use and
- Sleep apnea excluded

COPD = chronic obstructive pulmonary disease; GOLD = Global Strategy for the Diagnosis, Management, and Prevention of Chronic Obstructive Lung Disease⁶

SMOKING CESSATION

Cigarette smoking is the most common and easily identifiable risk factor for COPD exacerbation.⁵

Au et al⁵ found that quitting smoking reduces the risk of COPD exacerbation (adjust-

ed hazard ratio 0.78, 95% confidence interval [CI] 0.75–0.87), and the risk keeps decreasing the longer the patient stays off tobacco.⁵

Whether counseling hospitalized patients on smoking cessation reduces the COPD readmission rate has not been well studied. However, a meta-analysis of nine randomized controlled trials, two of which were done in the hospital, revealed higher abstinence rates in COPD patients who received extensive counseling on smoking cessation.⁷ For these reasons, hospitalized COPD patients who smoke should be strongly encouraged to quit.⁶

■ PNEUMOCOCCAL AND INFLUENZA VACCINATIONS

In a large retrospective study,⁸ pneumococcal vaccination was associated with a significantly lower risk of hospitalization for pneumonia in patients with chronic lung disease, including those with COPD (relative risk [RR] 0.57, 95% CI 0.38–0.84). The benefit was even greater with pneumococcal and influenza vaccinations during the influenza season (RR 0.28, 95% CI 0.14–0.58).

Randomized controlled trials indicate that influenza vaccination may reduce the rate of COPD exacerbations, especially in epidemic years when the proportion of exacerbations due to influenza is higher.⁹

Wongsurakiat et al¹⁰ found a significant reduction in the incidence of influenza-related acute respiratory illness in COPD patients in a well-designed randomized, placebo-controlled trial (RR 0.24, $P = .005$).¹⁰

Similarly, in another randomized controlled trial, pneumococcal vaccination was effective in preventing community-acquired pneumonia in COPD patients under age 65 and in those with severe airflow obstruction, although no statistically significant differences were found among other groups of patients with COPD.¹¹

Therefore, influenza and pneumococcal vaccinations are recommended by major COPD guidelines, such as GOLD (Global Strategy for the Diagnosis, Management, and Prevention of Chronic Obstructive Lung Disease).⁶

■ INHALERS

Inhaler therapy is recommended based on COPD severity according to GOLD classifi-

cation, and appropriate inhaler therapy with proper inhaler technique reduces the number of COPD exacerbations and hospitalizations.⁶

Long-acting beta-agonists and anticholinergics reduce the risk of COPD exacerbation and hospitalization and so are preferred over short-acting formulations except for patients in GOLD group A, ie, those who have few symptoms and are at low risk of exacerbations.⁶

Long-term treatment with **inhaled corticosteroids with long-acting bronchodilators** is recommended for patients at high risk of exacerbations (ie, those with two or more exacerbations in the previous year or a forced expiratory volume in 1 second [FEV₁] less than 50% of predicted).⁶

■ OXYGEN THERAPY

Two older randomized controlled trials, the Nocturnal Oxygen Therapy Trial and the Medical Research Council study, reviewed by Stoller et al,¹² provided clear evidence that oxygen therapy reduces the death rate and improves quality of life in COPD patients who have chronic resting hypoxemia (room air Pao₂ ≤ 55 mm Hg, or ≤ 59 mm Hg with signs of right-sided heart strain or polycythemia).

■ PULMONARY REHABILITATION

Pulmonary rehabilitation likely reduces hospital admissions by improving exercise capacity.¹³ A systematic review of six trials in 230 patients found that respiratory rehabilitation after an acute COPD exacerbation reduced the risk of COPD hospital admission (RR 0.26, 95% CI 0.12–0.54) and the risk of death (RR 0.45, 95% CI 0.22–0.91).¹³

■ OTHER INTERVENTIONS

Home noninvasive ventilator support reduced hospital and intensive care unit readmissions in select patients recurrently hospitalized for acidotic exacerbations of COPD in one small study.¹⁴

Long-term antibiotic therapy. Although there is evidence that azithromycin, taken daily for 1 year, decreases the frequency of COPD exacerbations,¹⁵ concern persists that this approach promotes antibiotic resistance, and the GOLD guidelines do not recommend

Needed:
coordinated
efforts
involving both
inpatient and
outpatient care

routinely using antibiotics in patients with clinically stable COPD.⁶

Roflumilast. According to the GOLD guidelines, the phosphodiesterase-4 inhibitor roflumilast (Daliresp) may be useful in reducing exacerbations in patients who have an FEV₁ less than 50% of predicted, chronic bronchitis, and frequent exacerbations.⁶

Referral. Patients who have severe recurrent COPD exacerbations despite appropriate therapy will likely benefit from referral to a pulmonary specialist for other options such as theophylline, lung-reduction surgery, and lung transplantation.

■ PATIENT EDUCATION AND OUTPATIENT COPD PROGRAMS

There is growing evidence that outpatient programs that provide education and medical support significantly reduce the rate of hospitalizations for COPD.^{16–18} Patient education includes symptom monitoring, early recogni-

tion of an exacerbation, appropriate use of inhalers and nebulizers, and advice on smoking cessation.¹⁶

On the other hand, a Veterans Administration randomized controlled trial was stopped early because of a higher rate of death in the group that underwent a comprehensive care-management program of COPD education, an action plan for identification and treatment of exacerbations, and scheduled proactive telephone calls for case management.¹⁹

Further study is needed to investigate the cost-effectiveness and safety of COPD management programs and whether to adopt such programs on a systematic level.

In conclusion, COPD patients require a comprehensive approach based on studied interventions. This may be achieved through systematic methods that allow each patient to benefit from all possible interventions appropriate for him or her. Hospitalization of COPD patients provides an excellent opportunity to implement this comprehensive approach. ■

■ REFERENCES

1. Westert GP, Lagoe RJ, Keskimäki I, Leyland A, Murphy M. An international study of hospital readmissions and related utilization in Europe and the USA. *Health Policy* 2002; 61:269–278.
2. Halpern MT, Stanford RH, Borker R. The burden of COPD in the USA: results from the Confronting COPD survey. *Respir Med* 2003; 97(suppl C):S81–S89.
3. Centers for Medicare and Medicaid Services. Readmissions reduction program. www.cms.gov/Medicare/Medicare-Fee-for-Service-Payment/AcuteInpatientPPS/Readmissions-Reduction-Program.html. Accessed August 9, 2014.
4. Yip NH, Yuen G, Lazar EJ, et al. Analysis of hospitalizations for COPD exacerbation: opportunities for improving care. *COPD* 2010; 7:85–92.
5. Au DH, Bryson CL, Chien JW, et al. The effects of smoking cessation on the risk of chronic obstructive pulmonary disease exacerbations. *J Gen Intern Med* 2009; 24:457–463.
6. Vestbo J, Hurd SS, Agustí AG, et al. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease: GOLD executive summary. *Am J Respir Crit Care Med* 2013; 187:347–365.
7. Thabane M; COPD Working Group. Smoking cessation for patients with chronic obstructive pulmonary disease (COPD): an evidence-based analysis. *Ont Health Technol Assess Ser* 2012; 12:1–50.
8. Nichol KL, Baken L, Wuorenma J, Nelson A. The health and economic benefits associated with pneumococcal vaccination of elderly persons with chronic lung disease. *Arch Intern Med* 1999; 159:2437–2442.
9. Poole PJ, Chacko E, Wood-Baker RW, Cates CJ. Influenza vaccine for patients with chronic obstructive pulmonary disease. *Cochrane Database Syst Rev* 2006; 1:CD002733.
10. Wongsurakiat P, Maranetra KN, Wasi C, Kositanont U, Dejsomritruttai W, Charoenratanakul S. Acute respiratory illness in patients with COPD and the effectiveness of influenza vaccination: a randomized controlled study. *Chest* 2004; 125:2011–2020.
11. Alfageme I, Vazquez R, Reyes N, et al. Clinical efficacy of anti-pneumococcal vaccination in patients with COPD. *Thorax* 2006; 61:189–195.
12. Stoller JK, Panos RJ, Krachman S, Doherty DE, Make B; Long-term Oxygen Treatment Trial Research Group. Oxygen therapy for patients with COPD: current evidence and the long-term oxygen treatment trial. *Chest* 2010; 138:179–187.
13. Puhan MA, Scharplatz M, Troosters T, Steurer J. Respiratory rehabilitation after acute exacerbation of COPD may reduce risk for readmission and mortality—a systematic review. *Respir Res* 2005; 6:54.
14. Tuggey JM, Plant PK, Elliott MW. Domiciliary non-invasive ventilation for recurrent acidotic exacerbations of COPD: an economic analysis. *Thorax* 2003; 58:867–871.
15. Albert RK, Connert J, Bailey WC, et al; COPD Clinical Research Network. Azithromycin for prevention of exacerbations of COPD. *N Engl J Med* 2011; 365:689–698.
16. Lawlor M, Kealy S, Agnew M, et al. Early discharge care with ongoing follow-up support may reduce hospital readmissions in COPD. *Int J Chron Obstruct Pulmon Dis* 2009; 4:55–60.
17. Gadoury MA, Schwartzman K, Rouleau M, et al; Chronic Obstructive Pulmonary Disease axis of the Respiratory Health Network, Fonds de la Recherche en Santé du Québec (FRSQ). Self-management reduces both short- and long-term hospitalisation in COPD. *Eur Respir J* 2005; 26:853–857.
18. Rice KL, Dewan N, Bloomfield HE, et al. Disease management program for chronic obstructive pulmonary disease: a randomized controlled trial. *Am J Respir Crit Care Med* 2010; 182:890–896.
19. Fan VS, Gaziano JM, Lew R, et al. A comprehensive care management program to prevent chronic obstructive pulmonary disease hospitalizations: a randomized, controlled trial. *Ann Intern Med* 2012; 156:673–683.
20. COPD Working Group. Noninvasive positive pressure ventilation for chronic respiratory failure patients with stable chronic obstructive pulmonary disease (COPD): an evidence-based analysis. *Ont Health Technol Assess Ser* 2012; 12(9):1–51.

ADDRESS: James C. Pile, MD, Department of Hospital Medicine, M2-Annex, Cleveland Clinic, 9500 Euclid Avenue, Cleveland, OH 44195; e-mail: pilej@ccl.org