### REVIEW



**EDUCATIONAL OBJECTIVE:** Readers will consider the possibility that patients with either asthma or chronic obstructive pulmonary disease may also have obstructive sleep apnea, and vice versa

#### SUMITA B. KHATRI, MD, MS

Co-Director, Asthma Center, Respiratory Institute, Cleveland Clinic; Associate Professor, Cleveland Clinic Lerner College of Medicine of Case Western Reserve University, Cleveland, OH OCTAVIAN C. IOACHIMESCU, MD, PhD Medical Director, Sleep Medicine Center, and Chief, Sleep Medicine Section, Atlanta VA Medical Center, Atlanta, GA; Associate Professor of Medicine, Emory University, Atlanta, GA

# The intersection of obstructive lung disease and sleep apnea

### ABSTRACT

Chronic obstructive pulmonary disease (COPD) and obstructive sleep apnea (OSA) have synergistic detrimental effects. Their comorbid association leads to compromised gas exchange (hypoxia and hypercapnia) and higher rates of morbidity and death. As our understanding of the pathophysiologic processes of sleep evolves, the relationship between OSA and obstructive lung diseases such as COPD ("overlap syndrome") or asthma ("alternative overlap syndrome") has become more apparent. The pathophysiology of the combined conditions and optimal management are still being defined, but the effect on quality of life and morbidity underscore the importance of proper diagnosis and appropriately tailored management in these patients.

### **KEY POINTS**

Obstructive lung diseases and OSA are both common and may exacerbate each other.

When assessing a patient with COPD, it may be prudent to think about whether the patient also has OSA, and vice versa.

Oxygen therapy lowers the risk of death in patients with COPD but may worsen hypercapnia and apneic episodes in those with OSA.

Continuous positive airway pressure is the first line of therapy for overlap syndrome. Daytime hypercapnia and nocturnal hypoxemia despite supplemental oxygen therapy are indications for nocturnal bilevel positive airway pressure therapy, regardless of the presence of OSA. **M**ANY PATIENTS who have obstructive lung disease, ie, chronic obstructive pulmonary disease (COPD) or asthma, also have obstructive sleep apnea (OSA), and vice versa.

The combination of COPD and OSA was first described almost 30 years ago by Flenley, who called it "overlap syndrome."<sup>1</sup> At that time, he recommended that a sleep study be considered in all obese patients with COPD who snore and in those who have frequent headaches after starting oxygen therapy. In the latter group, he doubted that nocturnal oxygen was the correct treatment. He also believed that the outcomes in patients with overlap syndrome were worse than those in patients with COPD or OSA alone. These opinions remain largely valid today.

We now also recognize the combination of asthma and OSA (alternative overlap syndrome) and collectively call both combinations obstructive lung disease-obstructive sleep apnea (OLDOSA) syndrome.<sup>2</sup> Interestingly, these relationships are likely bidirectional, with one condition aggravating or predisposing to the other.

Knowing that a patient has one of these overlap syndromes, one can initiate continuous positive airway pressure (CPAP) therapy, which can improve clinical outcomes.<sup>3-6</sup> Therefore, when evaluating a patient with asthma or COPD, one should consider OSA using a validated questionnaire and, if the findings suggest the diagnosis, polysomnography. Conversely, it is prudent to look for comorbid obstructive lung disease in patients with OSA, as interactions between upper and lower airway dysfunction may lead to distinctly different treatment and outcomes.

Here, we briefly review asthma and COPD, explore shared risk factors for sleep-disordered

doi:10.3949/ccjm.83a.14104

### TABLE 1

### Main differentiating and overlapping features of asthma and COPD

Feature	Asthma	COPD	Exceptions and overlapping features
Risk factors	Allergens	Smoking	Smoking may be an aggravating factor in asthma; children with early-life wheezing have a higher risk of developing asthma as teenagers and chronic obstructive pulmonary disease (COPD) as adults (the "Dutch hypothesis"); repeated airway infections may trigger development of either asthma or COPD, depending on the nature of infection and the intrinsic genetic predisposition (the "British hypothesis")
Anatomic segment involved	Airways	Airways and parenchyma	Severe asthma exacerbations may lead to adjacent pa- renchymal destruction, severe hyperinflation, and loss of scaffolding, or to radial or axial traction, or both, on the small airways
Atopy	Yes	No	Atopy may be present in some patients with COPD
Airway inflammation	Yes	Yes	Absent in mild asthma between exacerbations
Peripheral eosinophilia	Yes	No	Some patients with COPD may have eosinophilia; some patients with asthma may have neutrophilic infiltration (eg, severe asthma, asthma coexistent with gastroesophageal reflux and chronic rhinitis)
Airway eosinophilia	Yes	No	20%–40% of patients with COPD have airway eosinophilia (if one excludes eosinophilic bronchitis, up to 20% patients with COPD can still have some eosinophilic airway infiltration)
Exhaled gas (lower-airway) nitric oxide concentration	High	Low	May be reduced in some asthma phenotypes, eg, neutrophilic or paucigranulocytic; may be present in some patients with COPD
Diffusing lung capacity for carbon monoxide	Normal	Reduced (in emphysema)	Normal in chronic bronchitis; increased in acute asthma
Airway resistance	High	High	Normal in mild asthma between exacerbations; normal or slightly increased in emphysema
Airway hyperresponsiveness	Yes	No	May be present in COPD
Elastic recoil of the airways and parenchyma	Normal	Reduced (in emphysema)	Reduced in acute exacerbations of severe asthma; normal in chronic bronchitis
Symptoms	Episodic	Persistent	May be episodic in some patients with COPD, especially during significant environmental exposures
Comorbidities	Atopic dermatitis, allergic rhinitis	Cardiovascular disease	Obstructive sleep apnea may complicate both conditions

128 CLEVELAND CLINIC JOURNAL OF MEDICINE VOLUME 83 • NUMBER 2 FEBRUARY 2016

### **KHATRI AND IOACHIMESCU**





**Expiration** 

**Radial coupling** between airways and lung parenchyma during inspiration (left) and expiration (right) is affected by differences in upper airway resistance from obstructive sleep apnea;

(-) signifies subatmospheric pressure or pressure lower than mouth-level barometric pressure during inspiration;

(+) signifies supra-atmospheric pressure or pressure higher than mouth-level barometric pressure during expiration.



### **FIGURE 1.**

breathing and obstructive lung diseases, describe potential pathophysiologic mechanisms explaining these associations, and highlight the importance of recognizing and individually treating the overlaps of OSA and COPD or asthma.

### COPD AND ASTHMA ARE VERY COMMON

About 10% of the US population have COPD,<sup>7</sup> a preventable and treatable disease mainly caused by smoking, and a leading cause of sickness and death worldwide.<sup>8,9</sup>

CLEVELAND CLINIC JOURNAL OF MEDICINE VOLUME 83 • NUMBER 2 FEBRUARY 2016 129

About 8% of Americans have asthma,<sup>7</sup> which has become one of the most common chronic conditions in the Western world, affecting about 1 in 7 children and about 1 in 12 adults. The World Health Organization estimates that 235 million people suffer from asthma worldwide, and by 2025 this number is projected to rise to 400 million.<sup>10,11</sup>

The prevalence of these conditions in a particular population depends on the frequency of risk factors and associated morbidities, including OSA. These factors may allow asthma or COPD to arise earlier or have more severe manifestations.<sup>8,12</sup>

### Asthma and COPD: similarities and differences

Asthma and COPD share several features. Both are inflammatory airway conditions triggered or perpetuated by allergens, viral infection, tobacco smoke, products of biomass or fossil fuel combustion, and other substances. In both diseases, airflow is "obstructed" or limited, with a low ratio of forced expiratory volume in 1 second to forced vital capacity (FEV<sub>1</sub>/FVC). Symptoms can also be similar, with dyspnea, cough, wheezing, and chest tightness being the most frequent complaints. The similarities support the theory proposed by Orie et al<sup>13</sup> (the "Dutch hypothesis") that asthma and COPD may actually be manifestations of the same disease.

But there are also differences. COPD is strongly linked to cigarette smoking and has at least three phenotypes:

- Chronic bronchitis, defined clinically by cough and sputum production for more than 3 months per year for 2 consecutive years
- Emphysema, characterized anatomically by loss of lung parenchyma, as seen on tomographic imaging or examination of pathologic specimens
- A mixed form with bronchitic and emphysematous features, which is likely the most common.

Particularly in emphysematous COPD, smoking predisposes patients to gas-exchange abnormalities and low diffusing capacity for carbon monoxide.

In asthma, symptoms may be more episodic, the age of onset is often younger, and atopy is common, especially in allergic asthma. These episodic symptoms may correlate temporally with measurable airflow reversibility ( $\ge 12\%$  and  $\ge 200$  mL improvement in FVC or in FEV<sub>1</sub> after bronchodilator challenge).

However, the current taxonomy does not unequivocally divide obstructive lung diseases into asthma and COPD, and major features such as airway hyperresponsiveness, airflow reversibility, neutrophilic or CD8 lymphocytic airway inflammation, and lower concentration of nitric oxide in the exhaled air may be present in different phenotypes of both conditions (**Table 1**).

### AIRFLOW IN OBSTRUCTIVE LUNG DISEASES AND DURING SLEEP

Normal airflow involves a complex interplay between airway resistance and elastic recoil of the entire respiratory system, including the airways, the lung parenchyma, and the chest wall (Figure 1).

In asthma and COPD, resistance to airflow is increased, predominantly in the upper airways (nasal passages, pharynx, and larynx) and in the first three or four subdivisions of the tracheobronchial tree. The problem is worse during exhalation, when elastic recoil of the lung parenchyma and chest wall also increases airway resistance, reduces airway caliber, and possibly even constricts the bronchi. This last effect may occur either due to mass loading of the bronchial smooth muscles or to large intrathoracic transmural pressure shifts that may increase extravasation of fluid in the bronchial walls, especially with higher vascular permeability in inflammatory conditions.

Furthermore, interactions between the airway and parenchyma and between the upper and lower airways, as well as radial and axial coupling of these anatomic and functional components, contribute to complex interplay between airway resistance and parenchymalchest wall elastic energy—stretch or recoil.

The muscles of the upper and lower airway may not work together due to the loss of normal lung parenchyma (as in emphysema) or to the acute inflammation in the small airways and adjacent parenchyma (as in severe asthma exacerbations). This loss of coordination makes the upper airway more collapsible,

About 10% of the US population have COPD, and 8% have asthma

### **KHATRI AND IOACHIMESCU**



**FIGURE 2.** The main overlap syndromes. Sizes of circles roughly correspond to prevalences of the diseases they represent. COPD = chronic obstructive pulmonary disease; OLD = obstructive lung disease; OLDOSA = obstructive lung disease and obstructive sleep apnea; OSA = obstructive sleep apnea. OLD overlap syndrome has also been called asthma-COPD overlap syndrome.

Based on loachimescu OC, Teodorescu M. Integrating the overlap of obstructive lung disease and obstructive sleep apnoea: OLDOSA syndrome. Respirology 2013; 18:421–431; with permission from John Wiley & Sons, Inc.

a feature of OSA.

Additionally, obesity, gastroesophageal reflux, disease chronic rhinitis, nasal polyposis, and acute exacerbations of chronic systemic inflammation all contribute to more complex interactions between obstructive lung diseases and OSA.<sup>6</sup>

Sleep affects breathing, particularly in patients with respiratory comorbidities, and sleep-disordered breathing causes daytime symptoms and worsens quality of life.<sup>1,13–15</sup> During sleep, respiratory centers become less sensitive to oxygen and carbon dioxide; breathing becomes more irregular, especially during rapid eye movement (REM) sleep; the chest wall moves less, so that the tidal volume and functional residual capacity are lower; sighs, yawns, and deep breaths become limited; and serum carbon dioxide concentration may rise.

### OBSTRUCTIVE SLEEP APNEA

The prevalence of OSA, a form of sleep-disordered breathing characterized by limitation of inspiratory and (to a lesser degree) expiratory flow, has increased significantly in recent years, in parallel with the prevalence of its major risk factor, obesity.

OSA is generally defined as an apnea-hypopnea index of 5 or higher, ie, five or more episodes of apnea or hypopnea per hour.

OSA syndrome, ie, an apnea-hypopnea index of 5 or higher and excessive daytime sleepiness (defined by an Epworth Sleepiness Scale score > 10) was found in the initial analysis of the Wisconsin Sleep cohort in 1993 to be present in about 2% of women and 4% of men.<sup>16</sup> A more recent longitudinal analysis showed a significant increase—for example, in people 50 to 70 years old the prevalence was up to 17.6% in men and 7.5% in women.<sup>17</sup>

Upper airway resistance syndrome, a milder form of sleep-disordered breathing, is now included under the diagnosis of OSA, as its pathophysiology is not significantly different.<sup>18</sup>

In the next section, we discuss what happens when OSA overlaps with COPD (overlap syndrome) and with asthma ("alternative overlap syndrome")<sup>2,8</sup> (Figure 2).

### **OSA AND COPD (OVERLAP SYNDROME)**

Flenley<sup>1</sup> hypothesized that patients with COPD in whom supplemental oxygen worsened hypercapnia may also have OSA and called this association overlap syndrome.

#### How common is overlap syndrome?

Since both COPD and OSA are prevalent conditions, overlap syndrome may also be common.

The reported prevalence of overlap syndrome varies widely, depending on the population studied and the methods used. In various studies, COPD was present in 9% to 56% of patients with OSA,<sup>19–23</sup> and OSA was found in 5% to 85% of patients with COPD.<sup>24–27</sup>

hypothesis': asthma and COPD are manifestations of the same disease

The 'Dutch

Based on the prevalence of COPD in the general population (about 10%12) and that of sleep-disordered breathing (about 5% to 10%<sup>17</sup>), the expected prevalence of overlap syndrome in people over age 40 may be 0.5% to 1%.<sup>28</sup> In a more inclusive estimate with "subclinical" forms of overlap syndrome—ie, OSA defined as an apnea-hypopnea index of 5 or more (about 25% of the population<sup>17</sup>) and COPD Global initiative for Chronic Obstructive Lung Disease (GOLD) stage 1 (16.8% in the National Health and Nutrition Education Survey<sup>12</sup>)—the expected prevalence of overlap is around 4%. Some studies found a higher prevalence of COPD in OSA patients than in the general population,<sup>21,29</sup> while others did not.<sup>22,28,30</sup> The studies differed in how they defined sleep-disordered breathing.

Larger studies are needed to better assess the true prevalence of sleep-disordered breathing in COPD. They should use more sensitive measures of airflow and standardized definitions of sleep-disordered breathing and should include patients with more severe COPD. Fatigue and insomnia are common in COPD

Fatigue is strongly correlated with declining lung function, low exercise tolerance, and impaired quality of life in COPD.<sup>31</sup> Factors that contribute to fatigue include dyspnea, depression, and impaired sleep.<sup>32</sup> Some suggest that at least half of COPD patients have sleep complaints such as insomnia, sleep disruption, or sleep fragmentation.<sup>33</sup> Insomnia, difficulty falling asleep, and early morning awakenings are the most common complaints (30%–70% of patients) and are associated with daytime fatigue.<sup>34</sup> Conversely, comorbid OSA can contribute to fatigue and maintenance-type insomnia (ie, difficulty staying asleep and returning to sleep).

### Multiple mechanisms of hypoxemia in overlap syndrome

Oxygenation abnormalities and increased work of breathing contribute to the pathophysiology of overlap syndrome. In patients with COPD, oxygenation during wakefulness is a strong predictor of gas exchange during sleep.<sup>35</sup> Further, patients with overlap syndrome tend to have more severe hypoxia during sleep than patients with isolated COPD or OSA at rest or during exercise.<sup>36</sup>

In overlap syndrome, hypoxemia is the result of several mechanisms:

• Loss of upper airway muscle tone from intermittent episodes of obstructive apnea and hypopnea leads to upper airway collapse during sleep, particularly during REM sleep, increasing the severity of OSA.<sup>37</sup>

• Reductions in functional residual capacity from lying in the recumbent position and during REM sleep render patients with COPD more vulnerable, as compensatory use of accessory muscles to maintain near-normal ventilation in a hyperinflated state becomes impaired.<sup>37</sup>

• Alterations in pulmonary ventilation-perfusion matching may lead to altered carbon dioxide homeostasis and impaired oxygenation in patients with emphysema.

• Circadian variation in lower airway caliber may also be observed, in parallel with the bronchoconstriction caused by increased nocturnal vagotonia.

• Hypercapnia ( $Paco_2 \ge 45 \text{ mm Hg}$ ) may lead to overall reduced responsiveness of re-

spiratory muscles and to a blunted response of respiratory centers to low oxygen and high carbon dioxide levels.<sup>38</sup> Thus, hypercapnia is a better predictor of the severity of nocturnal hypoxemia than hypoxemia developing during exercise.<sup>39</sup>

In a person who is at near-maximal ventilatory capacity, even a mild increase in upper airway resistance (as seen with snoring, upper airway resistance syndrome, or OSA) increases the work of breathing. This phenomenon can lead to early arousals even before significant oxyhemoglobin desaturation occurs.

Normally, inspiratory flow limitation is counteracted by increasing inspiratory time to maintain ventilation. Patients with COPD may not be able to do this, however, as they need more time to breathe out due to narrowing of their lower airways.<sup>40</sup> The inability to compensate for upper airway resistance, similar to the increased work of breathing seen with exercise, may lead to early arousals and increased sleep fragmentation.

### **Consequences of overlap syndrome**

Patients with overlap syndrome appear to have higher morbidity and mortality rates than those with COPD or sleep-disordered breathing alone.

**Cor pulmonale.** Nighttime hypoxia is more severe and persistent in overlap syndrome than with COPD or OSA alone. This may contribute to more significant pulmonary hypertension and to the development of cor pulmonale, in which the right ventricle is altered in structure (eg, hypertrophied, dilated) or reduced in function, or both, from severe pulmonary hypertension.

In contrast to right ventricular failure due to disorders of the left heart, cor pulmonale is a result of diseases of the vasculature (eg, idiopathic pulmonary arterial hypertension), lung parenchyma (eg, COPD), upper airway (eg, OSA), or chest wall (eg, severe kyphoscoliosis). COPD is the most common cause of cor pulmonale in the United States, accounting for up to 30% of cases of cor pulmonale.<sup>41–45</sup> In OSA, cor pulmonale is seen in up to 20% of cases,<sup>43</sup> while in overlap syndrome cor pulmonale is encountered even more often (ie, in up to 80%); these patients have a dismal 5-year survival rate of about 30%.46

**Obesity hypoventilation syndrome** is characterized by obesity (body mass index  $\geq$ 30 kg/m<sup>2</sup>) and daytime hypercapnia (Paco<sub>2</sub>  $\geq$  45 mm Hg) that cannot be fully attributed to an underlying cardiopulmonary or neurologic condition.<sup>18</sup> Hypercapnia worsens during sleep (especially during REM sleep) and is often associated with severe arterial oxygen desaturation. Up to 90% of patients with obesity hypoventilation syndrome have comorbid OSA, and the rest generally have sleep-related hypoventilation, particularly during REM sleep.

In patients with obesity hypoventilation syndrome, daytime hypercapnia may improve or even normalize with adequate positive airway pressure treatment and sustained adherence to treatment.<sup>18</sup> Many patients with obesity hypoventilation syndrome respond to CPAP or bilevel positive airway pressure (BPAP), with improvement in daytime Paco<sub>2</sub>. However, normalization of daytime Paco<sub>2</sub> occurs only in a subgroup of patients. In contrast, treatment with oxygen therapy alone may worsen hypercapnia.

### Oxygen therapy for pure COPD, but maybe not for overlap syndrome

Continuous oxygen therapy reduces mortality in COPD,<sup>47,48</sup> but the duration and severity of hypoxemia that warrant oxygen therapy are less clear. Oxygen therapy in hypoxemic patients has been shown to improve sleep quality and reduce arousals.<sup>49</sup>

Indications for oxygen treatment of nocturnal hypoxemia are generally based on Medicare guidelines:

- At least 5 minutes of sleep with peripheral oxygen saturation ≤ 88% or Pao<sub>2</sub> ≤ 55 mm Hg, or
- A decrease in Pao<sub>2</sub> of more than 10 mm Hg or in peripheral oxygen saturation of more than 5% for at least 5 minutes of sleep and associated with signs or symptoms reasonably attributable to hypoxemia (group I criteria), or
- At least 5 minutes of sleep with peripheral oxygen saturation ≥ 89% or Pao<sub>2</sub> 56 to 59 mm Hg and pedal edema, pulmonary hypertension, cor pulmonale, or erythrocytosis (group II criteria).<sup>50</sup>

The prevalence of obstructive sleep apnea has increased in parallel with that of obesity Approximately 47% of COPD patients who are hypoxemic during the day spend about 30% of sleep time with an oxygen saturation less than 90%, even while on continuous oxygen therapy.<sup>51</sup> Current recommendations for nocturnal oxygen therapy are to increase the oxygen concentration by 1 L/minute above the baseline oxygen flow rate needed to maintain an oxygen saturation higher than 90% during resting wakefulness, using a nasal cannula or face mask.<sup>52</sup>

**Caveat.** In overlap syndrome, supplemental oxygen may prolong the duration of apnea episodes and worsen hypercapnia.

### Positive airway pressure for OSA

Positive airway pressure therapy improves cardiovascular outcomes in OSA.<sup>53</sup> Several studies<sup>54–58</sup> compared the effectiveness of CPAP vs BPAP as initial therapy for OSA but did not provide enough evidence to favor one over the other in this setting. Similarly, the results are mixed for the use of fixed or auto-adjusting BPAP as salvage therapy in patients who cannot tolerate CPAP.<sup>59–61</sup>

In overlap syndrome, CPAP or BPAP with or without supplemental oxygen has been investigated in several studies.<sup>26,62–65</sup> In general, the mortality rate of COPD patients who require oxygen therapy is quite high.<sup>47,66</sup> In hypoxemic COPD patients with moderate to severe sleep-disordered breathing, the 5-year survival rate was 71% in those treated with CPAP plus oxygen, vs 26% in those on oxygen alone, independent of baseline postbronchodilator FEV<sub>1</sub>.<sup>67</sup>

There is no specific  $\text{FEV}_1$  cutoff for prescribing CPAP. In general, daytime hypercapnia and nocturnal hypoxemia despite supplemental oxygen therapy are indications for BPAP therapy, regardless of the presence of OSA. Whether noninvasive nocturnal ventilation for COPD patients who do not have OSA improves long-term COPD outcomes is not entirely clear.<sup>65,68,69</sup>

Adding nocturnal BPAP in spontaneous timed mode to pulmonary rehabilitation for severe hypercapnic COPD was found to improve quality of life, mood, dyspnea, gas exchange, and decline in lung function.<sup>70</sup> Other studies noted that COPD patients hospitalized with respiratory failure who were random-

ized to noninvasive nocturnal ventilation plus oxygen therapy as opposed to oxygen alone experienced improvement in health-related quality of life and reduction in intensive-careunit length of stay but no difference in mortality or subsequent hospitalizations.<sup>69</sup> In stable hypercapnic COPD patients without OSA, there is no clear evidence that nocturnal noninvasive ventilation lessens the risk of death despite improved daytime gas exchange,<sup>71,72</sup> but additional long-term studies are needed.

Lung volume reduction surgery, a procedure indicated for highly selected patients with severe COPD, has been shown to reduce hyperinflation, improve nocturnal hypoxemia, and improve total sleep time and sleep efficiency in patients without sleep-disordered breathing.<sup>73</sup> More studies are needed to determine if reduction in lung hyperinflation has an impact on the occurrence of OSA and on morbidity related to sleep-disordered breathing.

#### Benefit of CPAP in overlap syndrome

In a nonrandomized study, Marin et al<sup>62</sup> found that overlap syndrome is associated with an increased risk of death and hospitalization due to COPD exacerbations. CPAP therapy was associated with improved survival rates and decreased hospitalization rates in these patients.

Stanchina et al,<sup>74</sup> in a post hoc analysis of an observational cohort, assessed the outcomes of 227 patients with overlap syndrome. Greater use of CPAP was found to be associated with lower mortality rates.

Jaoude et al<sup>75</sup> found that hypercapnic patients with overlap syndrome who were adherent to CPAP therapy had a lower mortality rate than nonadherent hypercapnic patients (P = .04). In a multivariate analysis, the comorbidity index was the only independent predictor of mortality in normocapnic patients with overlap syndrome, while CPAP adherence was associated with improved survival.

Lastly, patients with overlap syndrome tend to need more healthcare and accrue higher medical costs than patients with COPD alone. An analysis of a state Medicaid database that included COPD patients showed that beneficiaries with overlap syndrome spent at least

At near-maximal ventilatory capacity, even a mild increase in upper airway resistance increases the work of breathing \$4,000 more in medical expenditures than beneficiaries with "lone" COPD.<sup>24</sup>

In conclusion, CPAP is the first line of therapy for overlap syndrome, while daytime hypercapnia or nocturnal hypoxemia despite supplemental oxygen therapy are indications for nocturnal BPAP therapy, regardless of whether patients have OSA.

### OSA AND ASTHMA (ALTERNATIVE OVERLAP SYNDROME)

### **Epidemiology and clinical features**

The coexistence of asthma and OSA can begin in childhood and continue throughout adult life. A higher prevalence of lifetime asthma and OSA has been noted in children of racial and ethnic minorities, children of lower socioeconomic status, and those with atopy.<sup>76</sup>

In a pediatric asthma clinic, it was noted that 12 months into structured asthma management and optimization, children with sleep-disordered breathing were nearly four times more likely to have severe asthma at follow-up, even after adjusting for obesity, race, and gender.<sup>77</sup>

In adult patients with OSA, the prevalence of asthma is about 35%.<sup>78</sup> Conversely, people with asthma are at higher risk of OSA. High risk of OSA was more prevalent in a group of patients with asthma than in a general medical clinic population (39.5% vs 27.2%, *P* < .05).<sup>79</sup>

Analysis of a large prospective cohort found that asthma was a risk factor for newonset OSA. The incidence of OSA over 4 years in patients with self-reported asthma was 27%, compared with 16% without asthma. The relative risk adjusted for risk factors such as body mass index, age, and gender was 1.39 (95% confidence interval [CI] 15%–19%).<sup>80</sup>

Patients with asthma who are at high risk of OSA are more likely to have worse daytime and nighttime asthma symptoms. Interestingly, patients who are diagnosed with OSA and treated with CPAP seem to have better asthma control.

Patients with asthma who are more likely to have OSA are women (odds ratio [OR] 2.1), have greater asthma severity (OR 1.6), have gastroesophageal reflux disease (OR 2.7), and use inhaled corticosteroids (OR 4.0).<sup>81</sup> These associations are different than the traditional, population-wide risk factors for OSA, such as male sex, excess body weight, and nocturnal nasal congestion.<sup>82</sup>

OSA also worsens asthma control. Teodorescu et al<sup>15</sup> found that severe asthma was more frequent in older asthma patients (ages 60–75, prevalence 49%) than in younger patients (ages 18–59, 39%). Older adults with OSA were seven times as likely to have severe asthma (OR 6.6), whereas young adults with sleep apnea were only three times as likely (OR 2.6).

In a group of patients with difficult-to-treat asthma, OSA was significantly associated with frequent exacerbations (OR 3.4), an association similar in magnitude to that of psychological conditions (OR 10.8), severe sinus disease (OR 3.7), recurrent respiratory tract infections (OR 6.9), and gastroesophageal reflux disease (OR 4.9).<sup>83</sup> More than half of the patients had at least three of these comorbid conditions.

Sleep quality can greatly affect asthma control, and its importance is often underestimated. Patients with severe asthma have worse sleep quality than patients with milder asthma or nonasthmatic patients, even after excluding patients with a high risk of OSA, patients on CPAP therapy, and patients with a history of gastroesophageal reflux disease. Furthermore, regardless of asthma severity, sleep quality is a significant predictor of asthmarelated quality of life, even after accounting for body mass index, daytime sleepiness, and gastroesophageal reflux disease.<sup>84</sup>

### Pathophysiology of alternative overlap syndrome

Sleep significantly affects respiratory pathophysiology in asthma. The underlying mechanisms include physical and mechanical stressors, neurohormonal changes, hypoxia, confounding medical conditions, and local and systemic inflammatory changes.

Patients with nocturnal asthma experience more pronounced obstruction when sleep-deprived, suggesting that sleep loss may contribute to worsening airflow limitation.<sup>14</sup> Although changes in pulmonary mechanics and lung volumes may also have a role, volume-dependent airway narrowing does not appear to account for all observed nocturnal increases in airway resistance. Intrathoracic

Overlap syndrome with cor pulmonale typically has a poor prognosis; one study found a 5-year survival rate of 30%

### TABLE 2

## Proposed ABCD-3P-PQRST characterization system for asthma phenotypes, genotypes, and endotypes

ABCD	3P model <sup>a</sup>	PQRST		
<b>A</b> sthmatic symptoms	Paroxysmal (< 1 month) Persistent (1–6 months) Permanent (> 6 months)	<b>P</b> recipitating factors	Exertion (during, after), allergens, odors, drugs, tobacco smoke	
		<b>Q</b> ualitative description	Cough, wheezing, chest tightness, dyspnea	
		<b>R</b> eversibility to bronchodilator	Absolute change, percent change	
		<b>S</b> everity	Mild intermittent, mild, moderate, severe, or very severe persistent	
		Туре	Onset (adulthood or childhood), race and ethnicity, gender (male or female, prepubertal, adult, postmenopausal)	
<b>B</b> iomarkers	<b>P</b> redominant cells	Effector cells: eosinophils, neutrophils, mast cells, mixed cellularity, paucicellular Lymphocytes: T-helper 1 (Th1), Th2, Th17, regulatory T lymphocyes, natural killer cells		
	<b>P</b> redominant cytokines, immunoglobulins, molecules	Interleukin (IL) 4, IL-5, IL-9, IL-13, IL-33, periostin; interferon gamma, IL-15, IL-17, IL-18, IL-21, IL-22; immunoglobulin E or G; nitric oxide		
	<b>P</b> redisposing genotype	Specific gene polymorphisms, mutations		
<b>C</b> onditions associated	<b>P</b> renatal risk factors	Maternal smokin	Maternal smoking, diet, nutrition, antibiotic use, stress	
	Postnatal risk factors	Early-life wheezing, breastfeeding, early tobacco smoke exposure, viral infections, vitamin D deficiency, contact with animals, occupational exposures		
	<b>P</b> athogenically linked conditions	Atopy, allergic rhinitis, chronic rhinosinusitis, nasal polyposis, obstructive sleep apnea, gastroesophageal reflux disease, obesity, bronchiectases		
<b>D</b> rugs used	Anticholinergics (short-acting a ticosteroids (inhaled, oral), leu genase inhibitors), phosphodie stabilizers, anti–IL-5 agents (eg tralokinumab), anti-IgE (eg, on	and long-acting), be kotriene pathway-m sterase inhibitors (e g, mepolizumab, ber nalizumab), anti-tun	eta-2 adrenergic agonists (short-acting and long-acting), cor- nodifying agents (eg, leukotriene receptor agonists, lipooxy- eg, theophylline, roflumilast), cromolyn, nedocromil, mast cell nralizumab, reslizumab), anti–IL-13 agents (eg, lebrikizumab, nor necrosis factor alpha (golimumab)	
<sup>a</sup> 3P model for d of nitric oxide).	isease persistence should be characte	rized clinically, functio	onally, and biologically (eg, including exhaled gas concentration	

136 CLEVELAND CLINIC JOURNAL OF MEDICINE VOLUME 83 • NUMBER 2 FEBRUARY 2016

blood pooling may also contribute to nocturnal bronchoconstriction through stimulation of pulmonary C fibers and increased bronchial wall edema, a mechanism that may be similar to the "cardiac asthma" seen in left ventricular dysfunction.

Early studies of sleep-disordered breathing demonstrated that patients with asthma were breathing more irregularly (with hypopnea, apnea, and hyperpnea) in REM sleep than those without asthma.<sup>85</sup> Interestingly, REM-related hypoxia has also been noted in children with asthma.<sup>86</sup> This may be related to the increased cholinergic outflow that occurs during REM sleep, which in turn modulates the caliber and reactivity of the lower airways.

Physical changes such as upper airway collapse and reduced pharyngeal cross-sectional area may cause further mechanical strain.87 This can further propagate airway inflammation, alter airway mucosal muscle fibers, and stimulate neural reflexes, thereby increasing cholinergic tone and bronchoconstriction. Furthermore, heightened negative intrathoracic pressure during obstructive episodes can increase nocturnal pulmonary blood pooling.<sup>14</sup> Hypoxia itself can augment airway hyperresponsiveness via vagal pathways or carotid body receptors, increasing reactive oxygen species and inflammatory mediators. Local inflammation can "spill over" into systemic inflammatory changes, while alterations in airway inflammatory markers in asthma seem to follow a circadian rhythm, in parallel with the nocturnal worsening of the asthma symptoms.<sup>88</sup> Finally, altered sleep may be related to other comorbid conditions, such as gastroesophageal reflux disease, insomnia, and restless leg syndrome.

### Management and outcomes of alternative overlap syndrome

Despite optimization of asthma management, OSA can still significantly affect asthma control and symptoms.<sup>84</sup>

Interestingly, medications that reduce airway inflammation (eg, corticosteroids) may promote OSA. This occurrence cannot be fully explained by an increase in body mass, as more respiratory disturbances occur during sleep with continuous corticosteroid treatment even without increases in body mass index.<sup>87</sup> Therefore, these associations may be related to upper airway myopathy caused by the treatment, a small pharynx, facial dysmorphisms, or fat deposition.<sup>89</sup>

### Does CPAP improve asthma?

OSA is often unrecognized in patients with asthma, and treating it can have an impact on asthma symptoms.

CPAP therapy has not been shown to significantly change airway responsiveness or lung function, but it has been noted to significantly improve both OSA-related and asthma-related quality of life and reduce the use of rescue bronchodilators.<sup>3,90</sup> CPAP has demonstrated improvement of quality of life that positively correlated with body weight and apnea-hypopnea index at baseline, suggesting that asthmatic patients with greater obesity or worse OSA may benefit most from aggressive management.<sup>90</sup>

However, CPAP should be used only if the patient has confirmed OSA. Empiric use of CPAP without a diagnosis of OSA was poorly tolerated and failed to improve asthma symptoms or lung function.<sup>91</sup> More importantly, using CPAP in a patient who does not have OSA may contribute to further sleep disruption.<sup>91</sup>

Second-line treatments such as mandibular advancing devices and airway or bariatric surgery have not yet been studied in alternative overlap syndrome.

#### A multidimensional assessment of asthma

The Western world is experiencing an epidemic of obesity and of asthma. Obesity contributes to the pathogenesis of OSA by altering the anatomy and collapsibility of the upper airway, affecting ventilatory control and increasing respiratory workload. Another paradigm, supported by some evidence, is that OSA itself may contribute to the development of obesity. Both OSA and obesity lead to activation of inflammatory biologic cascades, which are likely the pathogenic mechanisms for their cardiovascular and metabolic consequences. As such, early recognition of OSA is important, as effective treatments are available.

In some patients, obesity may cause asthma, as obesity precedes the onset of asthma in a significant proportion of patients, and bariatric surgery for morbid obesity may resolve In overlap syndrome, oxygen may prolong the duration of apnea episodes and worsen hypercapnia asthma. The obese asthma phenotype seems to include chronic rhinosinusitis, gastroesophageal reflux disease, poorer asthma control, limited responsiveness to corticosteroids, and even different sets of biomarkers (eg, neutrophilic airway inflammation). A cohort of obese patients with poor asthma control demonstrated significant improvement in asthma symptoms, quality of life, and airway reactivity after weight loss from bariatric surgery.<sup>92</sup>

To improve our knowledge about airway disease phenotypes and endotypes and their response to therapy, we propose taking a multidimensional, structured assessment of all patients with asthma, using a schema we call "ABCD-3P-PQRST" (Table 2).

The purpose of using this type of system in

#### REFERENCES

- 1. Flenley DC. Sleep in chronic obstructive lung disease. Clin Chest Med 1985; 6:651–661.
- Ioachimescu OC, Teodorescu M. Integrating the overlap of obstructive lung disease and obstructive sleep apnoea: OLDOSA syndrome. Respirology 2013; 18:421–431.
- Ciftci TU, Ciftci B, Guven SF, Kokturk O, Turktas H. Effect of nasal continuous positive airway pressure in uncontrolled nocturnal asthmatic patients with obstructive sleep apnea syndrome. Respiratory Med 2005; 99:529–534.
- Kim MY, Jo EJ, Kang SY, et al. Obstructive sleep apnea is associated with reduced quality of life in adult patients with asthma. Ann Allergy Asthma Immunol 2013; 110:253–257.
- Teodorescu M, Polomis DA, Teodorescu MC, et al. Association of obstructive sleep apnea risk or diagnosis with daytime asthma in adults. J Asthma 2012; 49:620–628.
- Puthalapattu S, loachimescu OC. Asthma and obstructive sleep apnea: clinical and pathogenic interactions. J Investig Med 2014; 62:665–675.
- National Institutes of Health. National Heart, Lung, and Blood Institute. NHLBI Factbook, Fiscal Year 2007. Chapter 4. Disease Statistics. www. nhlbi.nih.gov/about/factbook-07/chapter4.htm. Accessed November 11, 2015.
- Vestbo J, Hurd SS, Agusti AG, et al. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease: GOLD executive summary. Am J Respir Crit Care Med 2013; 187:347–365.
- WHO. World Health Organization: Asthma. Fact sheet No 307. www. who.int/mediacentre/factsheets/fs307/en/. Accessed November 11, 2015.
- Masoli M, Fabian D, Holt S, Beasley R. The global burden of asthma: executive summary of the GINA Dissemination Committee report. Allergy 2004; 59:469–478.
- WHO. Chronic obstructive pulmonary disease (COPD). Fact sheet No 315. www.who.int/mediacentre/factsheets/fs315/en/index.html. 2011. Accessed November 11, 2015.
- Ford ES, Mannino DM, Wheaton AG, Giles WH, Presley-Cantrell L, Croft JB. Trends in the prevalence of obstructive and restrictive lung function among adults in the United States: findings from the National Health and Nutrition Examination surveys from 1988–1994 to 2007–2010. Chest 2013; 143:1395–1406.
- Orie N, Sluiter H, de Vries K, Tammeling G, Witkop J. The host factor in bronchitis. Paper presented at: Bronchitis—an international symposium 1961; Assen, Netherlands.
- Ballard RD. Sleep, respiratory physiology, and nocturnal asthma. Chronobiol Int 1999; 16:565–580.
- 15. Teodorescu M, Polomis DA, Gangnon RE, et al. Asthma control and its

clinics and research is to capture the multidimensionality of the disease and better develop future individualized therapeutic strategies by employing the latest advances in systems biology and computational methods such as cluster and principal component analysis.

Multidimensional assessments addressing airway problems such as asthma, COPD, OSA, other comorbidities and risk factors, and personalized management plans will need to be the basis of future therapeutic interventions. Increased attention to the complications of asthma and obstructive airway and lung diseases in our patients is imperative, specifically to develop effective systems of care, appropriate clinical guidelines, and research studies that lead to improved health outcomes.

relationship with obstructive sleep apnea (OSA) in older adults. Sleep Disord 2013; 2013:251567.

- Young T, Palta M, Dempsey J, Skatrud J, Weber S, Badr S. The occurrence of sleep-disordered breathing among middle-aged adults. N Engl J Med 1993; 328:1230–1235.
- Peppard PE, Young T, Barnet JH, Palta M, Hagen EW, Hla KM. Increased prevalence of sleep-disordered breathing in adults. Am J Epidemiol 2013; 177:1006–1014.
- International Classification of Sleep Disorders, 3rd ed.: Diagnostic and coding manual. Darien, Illinois: American Academy of Sleep Medicine. 2014.
- Lopez-Acevedo MN, Torres-Palacios A, Elena Ocasio-Tascon M, Campos-Santiago Z, Rodriguez-Cintron W. Overlap syndrome: an indication for sleep studies? A pilot study. Sleep Breath 2009; 13:409–413.
- 20. Scharf C, Li P, Muntwyler J, et al. Rate-dependent AV delay optimization in cardiac resynchronization therapy. PACE 2005; 28:279–284.
- Chaouat A, Weitzenblum E, Krieger J, Ifoundza T, Oswald M, Kessler R. Association of chronic obstructive pulmonary disease and sleep apnea syndrome. Am J Respir Crit Care Med 1995;151:82–86.
- Bednarek M, Plywaczewski R, Jonczak L, Zielinski J. There is no relationship between chronic obstructive pulmonary disease and obstructive sleep apnea syndrome: a population study. Respiration 2005; 72:142–149.
- Fletcher EC. Chronic lung disease in the sleep apnea syndrome. Lung 1990; 168(suppl):751–761.
- 24. Shaya FT, Lin PJ, Aljawadi MH, Scharf SM. Elevated economic burden in obstructive lung disease patients with concomitant sleep apnea syndrome. Sleep Breath 2009; 13:317–323.
- Larsson LG, Lindberg A, Franklin KA, Lundbäck B; Obstructive Lung Disease in Northern Sweden Studies. Obstructive sleep apnoea syndrome is common in subjects with chronic bronchitis. Report from the Obstructive Lung Disease in Northern Sweden studies. Respiration 2001; 68:250–255.
- Machado MC, Vollmer WM, Togeiro SM, et al. CPAP and survival in moderate-to-severe obstructive sleep apnoea syndrome and hypoxaemic COPD. Eur Resp J 2010; 35:132–137.
- 27. Guilleminault C, Cummiskey J, Motta J. Chronic obstructive airflow disease and sleep studies. Am Rev Respir Dis 1980; 122:397–406.
- Weitzenblum E, Chaouat A, Kessler R, Canuet M. Overlap syndrome: obstructive sleep apnea in patients with chronic obstructive pulmonary disease. Proc Am Thorac Soc 2008; 5:237–241.
- Bradley TD, Rutherford R, Lue F, et al. Role of diffuse airway obstruction in the hypercapnia of obstructive sleep apnea. Am Rev Respir Dis 1986; 134:920–924.
- Sanders MH, Newman AB, Haggerty CL, et al. Sleep and sleep-disordered breathing in adults with predominantly mild obstructive airway disease. Am J Respir Crit Care Med 2003; 167:7–14.
- 31. Breslin E, van der Schans C, Breukink S, et al. Perception of fatigue and
- 138 CLEVELAND CLINIC JOURNAL OF MEDICINE VOLUME 83 NUMBER 2 FEBRUARY 2016

quality of life in patients with COPD. Chest 1998; 114:958-964.

- Kapella MC, Larson JL, Patel MK, Covey MK, Berry JK. Subjective fatigue, influencing variables, and consequences in chronic obstructive pulmonary disease. Nurs Res 2006; 55:10–17.
- Klink M, Quan SF. Prevalence of reported sleep disturbances in a general adult population and their relationship to obstructive airways diseases. Chest 1987; 91:540–546.
- Bellia V, Catalano F, Scichilone N, et al. Sleep disorders in the elderly with and without chronic airflow obstruction: the SARA study. Sleep 2003; 26:318–323.
- Connaughton JJ, Catterall JR, Elton RA, Stradling JR, Douglas NJ. Do sleep studies contribute to the management of patients with severe chronic obstructive pulmonary disease? Am Rev Respir Dis 1988; 138:341–344.
- Mulloy E, McNicholas WT. Ventilation and gas exchange during sleep and exercise in severe COPD. Chest 1996; 109:387–394.
- Johnson MW, Remmers JE. Accessory muscle activity during sleep in chronic obstructive pulmonary disease. J Appl Physiol 1984; 57:1011– 1017.
- Douglas NJ, White DP, Pickett CK, Weil JV, Zwillich CW. Respiration during sleep in normal man. Thorax 1982; 37:840–844.
- Mulloy E, Fitzpatrick M, Bourke S, O'Regan A, McNicholas WT. Oxygen desaturation during sleep and exercise in patients with severe chronic obstructive pulmonary disease. Respir Med 1995; 89:193–198.
- 40. Herpel LB, Brown CD, Goring KL, et al. COPD cannot compensate for upper airway obstruction during sleep (abstract). Am J Respir Crit Care Med 2007; 175:A71.
- MacNee W. Pathophysiology of cor pulmonale in chronic obstructive pulmonary disease. Part 2. Am J Respir Crit Care Med 1994; 150:1158–1168.
- 42. MacNee W. Pathophysiology of cor pulmonale in chronic obstructive pulmonary disease. Part 1. Am J Respir Crit Care Med 1994; 150:833–852.
- Budev MM, Arroliga AC, Wiedemann HP, Matthay RA. Cor pulmonale: an overview. Semin Respir Crit Care Med 2003; 24:233–244.
- Simonneau G, Gatzoulis MA, Adatia I, et al. Updated clinical classification of pulmonary hypertension. J Am Coll Cardiol 2013; 62(25 suppl):D34–D41.
- 45. Naeije R. Pulmonary hypertension and right heart failure in chronic obstructive pulmonary disease. Proc Am Thorac Soc 2005; 2:20–22.
- Rasche K, Orth M, Kutscha A, Duchna HW. [Pulmonary diseases and heart function]. In German. Internist (Berl) 2007; 48:276–282.
- Continuous or nocturnal oxygen therapy in hypoxemic chronic obstructive lung disease: a clinical trial. Nocturnal Oxygen Therapy Trial Group. Ann Intern Med 1980; 93:391–398.
- Newman AB, Foster G, Givelber R, Nieto FJ, Redline S, Young T. Progression and regression of sleep-disordered breathing with changes in weight: the Sleep Heart Health Study. Arch Intern Med 2005; 165:2408–2413.
- Calverley PM, Brezinova V, Douglas NJ, Catterall JR, Flenley DC. The effect of oxygenation on sleep quality in chronic bronchitis and emphysema. Am Rev Respir Dis 1982; 126:206–210.
- Centers for Medicare and Medicaid Services. National coverage determination (NCD) for home use of oxygen (240.2). www.cms.gov/medicarecoverage-database/details/ncd-details.aspx?NCDId=169&ncdver=1&NCAI d=169&NcaName=Home+Use+of+Oxygen&lsPopup=y&bc=AAAAAAAAI AAA&. Accessed November 11, 2015.
- Plywaczewski R, Sliwinski P, Nowinski A, Kaminski D, Zielinski J. Incidence of nocturnal desaturation while breathing oxygen in COPD patients undergoing long-term oxygen therapy. Chest 2000; 117:679–683.
- Mokhlesi B, Tulaimat A, Faibussowitsch I, Wang Y, Evans AT. Obesity hypoventilation syndrome: prevalence and predictors in patients with obstructive sleep apnea. Sleep Breathing 2007; 11:117–124.
- Marin JM, Carrizo SJ, Vicente E, Agusti AG. Long-term cardiovascular outcomes in men with obstructive sleep apnoea-hypopnoea with or without treatment with continuous positive airway pressure: an observational study. Lancet 2005; 365:1046–1053.
- 54. Marin JM, DeAndres R, Alonso J, Sanchez A, Carrizo S. Long term mortality in the overlap syndrome. Eur Resp J 2008; 32(suppl 52):P865.
- Reeves-Hoche MK, Hudgel DW, Meck R, Witteman R, Ross A, Zwillich CW. Continuous versus bilevel positive airway pressure for obstructive sleep apnea. Am J Respir Crit Care Med 1995; 151:443–449.

- Gay PC, Herold DL, Olson EJ. A randomized, double-blind clinical trial comparing continuous positive airway pressure with a novel bilevel pressure system for treatment of obstructive sleep apnea syndrome. Sleep 2003; 26:864–869.
- 57. Blau A, Minx M, Peter JG, et al. Auto bi-level pressure relief-PAP is as effective as CPAP in OSA patients—a pilot study. Sleep Breath 2012; 16:773–779.
- Randerath WJ, Galetke W, Ruhle KH. Auto-adjusting CPAP based on impedance versus bilevel pressure in difficult-to-treat sleep apnea syndrome: a prospective randomized crossover study. Med Sci Monit 2003; 9:CR353–CR358.
- Schwartz SW, Rosas J, Iannacone MR, Foulis PR, Anderson WM. Correlates of a prescription for bilevel positive airway pressure for treatment of obstructive sleep apnea among veterans. J Clin Sleep Med 2013; 9:327–335.
- Gentina T, Fortin F, Douay B, et al. Auto bi-level with pressure relief during exhalation as a rescue therapy for optimally treated obstructive sleep apnoea patients with poor compliance to continuous positive airways pressure therapy--a pilot study. Sleep Breathing 2011; 15:21–27.
- Ballard RD, Gay PC, Strollo PJ. Interventions to improve compliance in sleep apnea patients previously non-compliant with continuous positive airway pressure. J Clinical Sleep Med 2007; 3:706–712.
- Marin JM, Soriano JB, Carrizo SJ, Boldova A, Celli BR. Outcomes in patients with chronic obstructive pulmonary disease and obstructive sleep apnea: the overlap syndrome. Am J Respir Crit Care Med 2010; 182:325–331.
- de Miguel J, Cabello J, Sanchez-Alarcos JM, Alvarez-Sala R, Espinos D, Alvarez-Sala JL. Long-term effects of treatment with nasal continuous positive airway pressure on lung function in patients with overlap syndrome. Sleep Breath 2002; 6:3–10.
- Mansfield D, Naughton MT. Effects of continuous positive airway pressure on lung function in patients with chronic obstructive pulmonary disease and sleep disordered breathing. Respirology 1999; 4:365–370.
- McEvoy RD, Pierce RJ, Hillman D, et al. Nocturnal non-invasive nasal ventilation in stable hypercapnic COPD: a randomised controlled trial. Thorax 2009; 64:561–566.
- Long term domiciliary oxygen therapy in chronic hypoxic cor pulmonale complicating chronic bronchitis and emphysema. Report of the Medical Research Council Working Party. Lancet 1981; 1:681–686.
- Machado MC, Vollmer WM, Togeiro SM, et al. CPAP and survival in moderate-to-severe obstructive sleep apnoea syndrome and hypoxaemic COPD. Eur Respir J 2010; 35:132–137.
- Casanova C, Celli BR, Tost L, et al. Long-term controlled trial of nocturnal nasal positive pressure ventilation in patients with severe COPD. Chest 2000; 118:1582–1590.
- Clini E, Sturani C, Rossi A, et al. The Italian multicentre study on noninvasive ventilation in chronic obstructive pulmonary disease patients. Eur Respir J 2002; 20:529–538.
- Duiverman ML, Wempe JB, Bladder G, et al. Two-year home-based nocturnal noninvasive ventilation added to rehabilitation in chronic obstructive pulmonary disease patients: a randomized controlled trial. Respir Res 2011; 12:112.
- Gay PC, Hubmayr RD, Stroetz RW. Efficacy of nocturnal nasal ventilation in stable, severe chronic obstructive pulmonary disease during a 3-month controlled trial. Mayo Clin Proc 1996; 71:533–542.
- Meecham Jones DJ, Paul EA, Jones PW, Wedzicha JA. Nasal pressure support ventilation plus oxygen compared with oxygen therapy alone in hypercapnic COPD. Am J Respir Crit Care Med 1995; 152:538–544.
- Krachman SL, Chatila W, Martin UJ, et al. Effects of lung volume reduction surgery on sleep quality and nocturnal gas exchange in patients with severe emphysema. Chest 2005; 128:3221–3228.
- Stanchina ML, Welicky LM, Donat W, Lee D, Corrao W, Malhotra A. Impact of CPAP use and age on mortality in patients with combined COPD and obstructive sleep apnea: the overlap syndrome. J Clin Sleep Med 2013; 9:767–772.
- 75. Jaoude P, Kufel T, El-Solh AA. Survival benefit of CPAP favors hypercapnic patients with the overlap syndrome. Lung 2014; 192:251–258.
- Ramagopal M, Mehta A, Roberts DW, et al. Asthma as a predictor of obstructive sleep apnea in urban African-American children. J Asthma 2009; 46:895–899.

- 77. Ross KR, Storfer-Isser A, Hart MA, et al. Sleep-disordered breathing is associated with asthma severity in children. J Ped 2012; 160:736–742.
- Alharbi M, Almutairi A, Alotaibi D, Alotaibi A, Shaikh S, Bahammam AS. The prevalence of asthma in patients with obstructive sleep apnoea. Prim Care Respir J 2009; 18:328–330.
- Auckley D, Moallem M, Shaman Z, Mustafa M. Findings of a Berlin Questionnaire survey: comparison between patients seen in an asthma clinic versus internal medicine clinic. Sleep Med 2008; 9:494–499.
- Teodorescu M, Barnet JH, Hagen EW, Palta M, Young TB, Peppard PE. Association between asthma and risk of developing obstructive sleep apnea. JAMA 2015; 313:156–164.
- Teodorescu M, Polomis DA, Hall SV, et al. Association of obstructive sleep apnea risk with asthma control in adults. Chest 2010; 138:543–550.
- Larsson LG, Lindberg A, Franklin KA, Lundback B. Gender differences in symptoms related to sleep apnea in a general population and in relation to referral to sleep clinic. Chest 2003; 124:204–211.
- ten Brinke A, Sterk PJ, Masclee AA, et al. Risk factors of frequent exacerbations in difficult-to-treat asthma. Eur Respir J 2005; 26:812–818.
- Luyster FS, Teodorescu M, Bleecker E, et al. Sleep quality and asthma control and quality of life in non-severe and severe asthma. Sleep Breath 2012; 16:1129–1137.
- Catterall JR, Douglas NJ, Calverley PM, et al. Irregular breathing and hypoxaemia during sleep in chronic stable asthma. Lancet 1982; 1:301–304.

- Perez GF, Gutierrez MJ, Huseni S, et al. Oximetry signal processing identifies REM sleep-related vulnerability trait in asthmatic children. Sleep Disord 2013; 2013:406157.
- Yigla M, Tov N, Solomonov A, Rubin AH, Harlev D. Difficult-to-control asthma and obstructive sleep apnea. J Asthma 2003; 40:865–871.
- Kelly EA, Houtman JJ, Jarjour NN. Inflammatory changes associated with circadian variation in pulmonary function in subjects with mild asthma. Clin Exper Allergy 2004; 34:227–233.
- Bohadana AB, Hannhart B, Teculescu DB. Nocturnal worsening of asthma and sleep-disordered breathing. J Asthma 2002; 39:85–100.
- Lafond C, Series F, Lemiere C. Impact of CPAP on asthmatic patients with obstructive sleep apnoea. Eur Respir J 2007; 29:307–311.
- Martin RJ, Pak J. Nasal CPAP in nonapneic nocturnal asthma. Chest 1991; 100:1024–1027.
- Dixon AE, Pratley RE, Forgione PM, et al. Effects of obesity and bariatric surgery on airway hyperresponsiveness, asthma control, and inflammation. J Allergy Clin Immunol 2011; 128:508–515 e501–502.

ADDRESS: Sumita B. Khatri, MD MS, Respiratory Institute, A90, Cleveland Clinic, 9500 Euclid Avenue, Cleveland, OH 44195; e-mail: khatris@ccf.org; and Octavian C. Ioachimescu, MD, PhD, Atlanta VA Clinic-Sleep Medicine Center, 250 North Arcadia Avenue, Decatur, GA 30033; e-mail: oioac@yahoo.com