

REVIEW

LUKE D. KIM, MD, FACP, CMD

Assistant Professor of Medicine, Cleveland Clinic Lerner College of Medicine of Case Western Reserve University; Center for Geriatric Medicine, Medicine Institute, Cleveland Clinic

KENNETH KONCILJA, MD

Geriatric Medicine Fellow, University of California San Francisco, Division of Geriatrics, Department of Medicine

CRAIG NIELSEN, MD, FACP

Associate Professor of Medicine, Cleveland Clinic Lerner College of Medicine of Case Western Reserve University; Vice Chairman, Department of Internal Medicine, Medicine Institute, Cleveland Clinic; Deputy Editor, *Cleveland Clinic Journal of Medicine*

Medication management in older adults

ABSTRACT

Managing medications is a major part of providing care to older adults. Polypharmacy is common in the elderly and is fraught with risks. A careful and systematic approach is needed for managing drug therapy in these patients, recognizing the patient's specific goals.

KEY POINTS

Statins, anticholinergics, benzodiazepines, antipsychotics, and proton pump inhibitors are widely prescribed.

In older patients, a periodic comprehensive medication review is needed to reevaluate the risks and the benefits of current medications in light of goals of care, life expectancy, and the patient's preferences.

The Beers criteria and the Screening Tool of Older Persons' Potentially Inappropriate Prescriptions provide valuable guidance for safe prescribing in older adults.

MEDICATIONS STARTED FOR appropriate indications in middle age may need to be monitored more closely as the patient ages. Some drugs may become unnecessary or even dangerous as the patient ages, functional status and renal function decline, and goals of care change.

See related editorial, page 136

Older adults tend to have multiple illnesses and therefore take more drugs, and polypharmacy increases the risk of poor outcomes. The number of medications a person uses is a risk factor for adverse drug reactions, nonadherence, financial burden, drug-drug interactions, and worse outcomes.¹

The prevalence of polypharmacy increased from an estimated 8.2% to 15% from 1999 to 2011 based on the National Health and Nutrition Examination Survey.² Guideline-based therapy for specific diseases may lead to the addition of more medications to reach disease targets.³ Most older adults in the United States compound the risk of prescribed medications by also taking over-the-counter medications and dietary supplements.⁴

In addition, medications are often used in older adults based on studies of younger persons without significant comorbidities. Applying clinical guidelines based on these studies to older adults with comorbidity and functional impairment is challenging.⁵ Age-related pharmacokinetic and pharmacodynamic changes increase the risk of adverse drug reactions.⁶

In this article, we review commonly used medications that are potentially inappropriate based on clinical practice. We also review tools to evaluate appropriate drug therapy in older adults.

doi:10.3949/ccjm.85a.16109

TABLE 1

Drugs with strong anticholinergic properties

Class	Examples
Antihistamines	Diphenhydramine, hydroxyzine, meclizine
Antiparkinsonian agents	Benzotropine
Skeletal muscle relaxants	Cyclobenzaprine, methocarbamol
Antidepressants	Amitriptyline, imipramine, nortriptyline, paroxetine
Antipsychotics	Olanzapine
Antiarrhythmics	Disopyramide
Antimuscarinics (for urinary incontinence)	Oxybutynin, tolterodine, trospium
Antiemetics	Prochlorperazine, promethazine
Antispasmodics	Hyoscyamine, scopolamine

DRUGS THAT ARE COMMONLY USED, BUT POTENTIALLY INAPPROPRIATE

Statins

Statins are effective when used as *secondary* prevention in older adults,⁷ but their efficacy when used as *primary* prevention of atherosclerotic cardiovascular disease in people age 75 and older is questionable.⁸ Nevertheless, they are widely used for this purpose. For example, before the 2013 joint guidelines of the American College of Cardiology and the American Heart Association (ACC/AHA) were released, 22% of patients age 80 and older in the Geisinger health system were taking a statin for primary prevention.⁹

The 2013 ACC/AHA guidelines included a limited recommendation for statins for primary prevention of atherosclerotic cardiovascular disease in adults age 75 and older.¹⁰ The guideline noted, however, that few data were available to support this recommendation.¹⁰

In a systematic review of 18 randomized clinical trials of statins for primary prevention of atherosclerotic cardiovascular disease, the mean age was 57, yet conclusions were extrapolated to an older patient population.¹¹ The estimated 10-year risk of atherosclerotic cardiovascular disease based on pooled cohort risk equations of adults age 75 and older always exceeds the 7.5% treatment threshold recommended by the guidelines.⁸

Myopathy is a common adverse effect of statins. In addition, statins interact with other

drugs that inhibit the cytochrome P450 3A4 isoenzyme, such as amlodipine, amiodarone, and diltiazem.^{8,12} If statin therapy caused no functional limitation due to muscle pain or weakness, statins for primary prevention would be cost-effective, but even a small increase in adverse effects in an elderly patient can offset the cardiovascular benefit.¹³ A recent post hoc secondary analysis found no benefit of pravastatin for primary prevention in adults age 75 and older.¹⁴

Thus, statin treatment for primary prevention in older patients should be individualized, based on life expectancy, function, and cardiovascular risk. Statin therapy does not replace modification of other risk factors.

Anticholinergics

Drugs with anticholinergic properties are commonly prescribed in the elderly for conditions such as muscle spasm, overactive bladder, psychiatric disorders, insomnia, extrapyramidal symptoms, vertigo, pruritus, peptic ulcer disease, seasonal allergies, and even the common cold,¹⁵ and many of the drugs often prescribed have strong anticholinergic properties (Table 1). Taking multiple medications with anticholinergic properties results in a high “anticholinergic burden,” which is associated with falls, impulsive behavior, poor physical performance, loss of independence, dementia, delirium, and brain atrophy.^{15–18}

The 2014 American College of Physicians guideline on nonsurgical management of uri-

With statins, even a small increase in adverse effects in an elderly patient can offset cardiovascular benefit

TABLE 2

The 2015 Beers criteria: Selected drugs to avoid in older adults

Drug class (example)	Recommendation	Rationale	Quality of evidence	Strength of recommendation
First-generation antihistamines (diphenhydramine)	Avoid	Highly anticholinergic	Moderate	Strong
Antiparkinsonian agents (benztropine)	Avoid	Not recommended for prevention of extrapyramidal symptoms with antipsychotics; more-effective agents available for treatment of Parkinson disease	Moderate	Strong
Antispasmodics (hyoscyamine)	Avoid	Highly anticholinergic	Moderate	Strong
Antidepressants (amitriptyline)	Avoid	Highly anticholinergic	High	Strong
Antipsychotics (conventional or atypical)	Avoid except for schizophrenia, bipolar disorder, or short-term use as antiemetic during chemotherapy	Increased risk of stroke and death in persons with dementia	Moderate	Strong
Skeletal muscle relaxants (methocarbamol)	Avoid	Most muscle relaxants are poorly tolerated by older adults	Moderate	Strong
Benzodiazepine (lorazepam)	Avoid	All benzodiazepines increase risk of cognitive impairment, delirium, falls, fractures, and motor vehicle crashes in older adults	Moderate	Strong
Nonbenzodiazepine and benzodiazepine hypnotics (zolpidem)	Avoid	Adverse events similar to those of benzodiazepines in older adults	Moderate	Strong
Proton pump inhibitors	Avoid using for > 8 weeks unless for high-risk patients	Risk of <i>Clostridium difficile</i> infection, bone loss, and fractures	High	Strong

Based on information in reference 58.

nary incontinence in women recommends pharmacologic treatment for urgency and stress urinary incontinence after failure of nonpharmacologic therapy,¹⁹ and many drugs for these urinary symptoms have anticholinergic properties. If an anticholinergic is necessary, an agent that results in a lower anticholinergic burden should be considered in older patients.

A pharmacist-initiated medication review and intervention may be another way to adjust medications to reduce the patient's anticholinergic burden.^{20,21} The common use of anticholinergic drugs in older adults reminds us to monitor their use closely.²²

Benzodiazepines and nonbenzodiazepines

Benzodiazepines are among the most commonly prescribed psychotropics in developed countries and are prescribed mainly by primary care physicians rather than psychiatrists.²³

In 2008, 5.2% of US adults ages 18 to 80 used a benzodiazepine, and long-term use was more prevalent in older patients (ages 65–80).²³

Benzodiazepines are prescribed for anxiety,²⁴ insomnia,²⁵ and agitation. They can cause withdrawal²⁶ and have potential for abuse.²⁷ Benzodiazepines are associated with cognitive decline,²⁸ impaired driving,²⁹ falls,³⁰ and hip fractures³¹ in older adults.

TABLE 3

The Screening Tool of Older Persons' Potentially Inappropriate Prescriptions (STOPP): Selected warnings and recommendations

Tricyclic antidepressants	Avoid use with dementia, narrow-angle glaucoma, cardiac conduction abnormality, prostatism, or history of urinary retention because of risk of worsening these conditions
Benzodiazepines	No indication for treatment beyond 4 weeks Risk of prolonged sedation, confusion, impaired balance, falls, traffic accidents Withdraw gradually if taken for > 2 weeks because of risk of withdrawal syndrome
Proton pump inhibitors	For uncomplicated peptic ulcer disease or erosive peptic esophagitis, discontinue full therapeutic dosage before 8 weeks, or reduce dosage
Hypnotic "z" drugs (eg, zopiclone, zolpidem, zaleplon)	May cause protracted daytime sedation, ataxia
Antimuscarinic drugs	For overactive bladder with concurrent dementia or chronic cognitive impairment, there is risk of increased confusion, agitation For narrow-angle glaucoma, there is risk of acute exacerbation of glaucoma For chronic prostatism, there is risk of urinary retention

Based on information in references 60 and 61.

Nondrug therapies for reflux such as weight loss and elevation of the head of the bed may improve symptoms

In addition, use of nonbenzodiazepine hypnotics (eg, zolpidem) is on the rise,³² and these drugs are known to increase the risk of hip fracture in nursing home residents.³³

The American Geriatrics Society, through the American Board of Internal Medicine's Choosing Wisely campaign, recommends avoiding benzodiazepines as a first-line treatment for insomnia, agitation, or delirium in older adults.³⁴ Yet prescribing practices with these drugs in primary care settings conflict with guidelines, partly due to lack of training in constructive strategies regarding appropriate use of benzodiazepines.³⁵ Educating patients on the risks and benefits of benzodiazepine treatment, especially long-term use, has been shown to reduce the rate of benzodiazepine-associated secondary events.³⁶

Antipsychotics

Off-label use of antipsychotics is common and is increasing in the United States. In 2008, off-label use of antipsychotic drugs accounted for an estimated \$6 billion.³⁷ A common off-label use is to manage behavioral symptoms of dementia, despite a black-box warning about

an increased risk of death in patients with dementia who are treated with antipsychotics.^{38,39} The Choosing Wisely campaign recommends against prescribing antipsychotics as a first-line treatment of behavioral and psychological symptoms of dementia.³⁴

Antipsychotic drugs are associated with risk of acute kidney injury,⁴⁰ as well as increased risk of falls and fractures (eg, a 52% higher risk of a serious fall, and a 50% higher risk of a nonvertebral osteoporotic fracture).⁴¹

Patients with dementia often exhibit aggression, resistance to care, and other challenging or disruptive behaviors. In such instances, antipsychotic drugs are often prescribed, but they provide limited and inconsistent benefits, while causing oversedation and worsening of cognitive function and increasing the likelihood of falling, stroke, and death.^{38,39,41}

Because pharmacologic treatments for dementia are only modestly effective, have notable risks, and do not treat some of the behaviors that family members and caregivers find most distressing, nonpharmacologic

measures are recommended as first-line treatment.⁴² These include caregiver education and support, training in problem-solving, and targeted therapy directed at the underlying causes of specific behaviors (eg, implementing nighttime routines to address sleep disturbances).⁴² Nonpharmacologic management of behavioral symptoms in dementia can significantly improve quality of life for patients and caregivers.⁴² Use of antipsychotic drugs in patients with dementia should be limited to cases in which nonpharmacologic measures have failed and patients pose an imminent threat to themselves or others.⁴³

Proton pump inhibitors

Proton pump inhibitors are among the most commonly prescribed medications in the United States, and their use has increased significantly over the decade. It has been estimated that between 25% and 70% of these prescriptions have no appropriate indication.⁴⁴

There is considerable excess use of acid suppressants in both inpatient and outpatient settings.^{45,46} In one study, at discharge from an internal medicine service, almost half of patients were taking a proton pump inhibitor.⁴⁷

Evidence-based guidelines recommend these drugs to treat gastroesophageal reflux disease, nonerosive reflux disease, erosive esophagitis, dyspepsia, and peptic ulcer disease. However, long-term use (ie, beyond 8 weeks) is recommended only for patients with erosive esophagitis, Barrett esophagus, a pathologic hypersecretory condition, or a demonstrated need for maintenance treatment for reflux disease.⁴⁸

Although proton pump inhibitors are highly effective and have low toxicity, there are reports of an association with *Clostridium difficile* infection,⁴⁹ community-acquired pneumonia,⁵⁰ hip fracture,⁵¹ vitamin B₁₂ deficiency,⁵² atrophic gastritis,⁵³ kidney disease,⁵⁴ and dementia.⁵⁵

Nondrug therapies such as weight loss and elevation of the head of the bed may improve esophageal pH levels and reflux symptoms.⁵⁶

Deprescribing.org has practical advice for healthcare providers, patients, and caregivers on how to discontinue proton pump inhibitors, including videos, algorithms, and guidelines.

TOOLS TO EVALUATE APPROPRIATE DRUG THERAPY

Beers criteria

The Beers criteria (Table 2), developed in 1991 by a geriatrician as an approach to safer, more effective drug therapy in frail elderly nursing home patients,⁵⁷ were updated by the American Geriatrics Society in 2015 for use in any clinical setting.⁵⁸ (The criteria are also available as a smartphone application through the American Geriatrics Society at www.americangeriatrics.org.)

The Beers criteria offer evidence-based recommendations on drugs to avoid in the elderly, along with the rationale for use, the quality of evidence behind the recommendation, and the graded strength of the recommendation. The Beers criteria should be viewed through the lens of clinical judgment to offer safer nonpharmacologic and pharmacologic treatments.

The Joint Commission recommends medication reconciliation at every transition of care.⁵⁹ The Beers criteria are a good starting point for a comprehensive medication review.

STOPP/START criteria

Another tool to aid safe prescribing in older adults is the Screening Tool of Older Persons' Potentially Inappropriate Prescriptions (STOPP), used in conjunction with the Screening Tool to Alert Doctors to Right Treatment (START). The STOPP/START criteria^{60,61} are based on an up-to-date literature review and consensus (Table 3).

THE BOTTOM LINE

Physicians caring for older adults need to diligently weigh the benefits of drug therapy and consider the patient's care goals, current level of functioning, life expectancy, values, and preferences. Statin therapy for primary prevention, anticholinergics, benzodiazepines, antipsychotics, and proton pump inhibitors are widely used without proper indications, pointing to the need for a periodic comprehensive review of medications to reevaluate the risks vs the benefits of the patient's medications. The Beers criteria and the STOPP/START criteria can be useful tools for this purpose.

For older patients, the Joint Commission recommends medication reconciliation at every transition of care

REFERENCES

- Steinman MA. Polypharmacy—time to get beyond numbers. *JAMA Intern Med* 2016; 176:482–483.
- Kantor ED, Rehm CD, Haas JS, Chan AT, Giovannucci EL. Trends in prescription drug use among adults in the United States from 1999–2012. *JAMA* 2015; 314:1818–1831.
- Tinetti ME, Bogardus ST Jr, Agostini JV. Potential pitfalls of disease-specific guidelines for patients with multiple conditions. *N Engl J Med* 2004; 351:2870–2874.
- Qato DM, Wilder J, Schumm LP, Gillet V, Alexander GC. Changes in prescription and over-the-counter medication and dietary supplement use among older adults in the United States, 2005 vs 2011. *JAMA Intern Med* 2016; 176:473–482.
- Boyd CM, Darer J, Boult C, et al. Clinical practice guidelines and quality of care for older patients with multiple comorbid diseases: implications for pay for performance. *JAMA* 2005; 294:716–724.
- Atkin PA, Veitch PC, Veitch EM, Ogle SJ. The epidemiology of serious adverse drug reactions among the elderly. *Drugs Aging* 1999; 14:141–152.
- Collins R, Reith C, Emberson J, et al. Interpretation of the evidence for the efficacy and safety of statin therapy. *Lancet* 2016; 338:2532–2561.
- Gurwitz JH, Go AS, Fortman SP. Statins for primary prevention in older adults: uncertainty and the need for more evidence. *JAMA* 2016; 316:1971–1972.
- Chokshi NP, Messerli FH, Sutlin D, Supariwala AA, Shah NR. Appropriateness of statins in patients aged ≥ 80 years and comparison to other age groups. *Am J Cardiol* 2012; 110:1477–1481.
- Stone NJ, Robinson J, Lichtenstein AH, et al; American College of Cardiology/American Heart Association Task Force on Practice Guidelines. 2013 ACC/AHA guideline on the treatment of blood cholesterol to reduce atherosclerotic cardiovascular risk in adults: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *Circulation* 2014; 129(suppl 2):S1–S45.
- Taylor F, Huffman MD, Macedo AF, et al. Statins for the primary prevention of cardiovascular disease. *Cochrane Database Syst Rev* 2013; 1:CD004816.
- Chatzizisis YS, Koskinas KC, Misirli G, Vaklavas C, Hatzitolios A, Giannoglou GD. Risk factors and drug interactions predisposing to statin-induced myopathy: implications for risk assessment, prevention and treatment. *Drug Saf* 2010; 33:171–187.
- Odden MC, Pletcher MJ, Coxson PG, et al. Cost-effectiveness and population impact of statins for primary prevention in adults aged 75 years or older in the United States. *Ann Intern Med* 2015; 162:533–541.
- Han BH, Sutlin D, Williamson JD, et al. Effect of statin treatment vs usual care on primary cardiovascular prevention among older adults. The ALLHAT-LLT randomized clinical trial. *JAMA Intern Med* 2017; 177:955–965.
- Gray SL, Anderson ML, Dublin S, et al. Cumulative use of strong anticholinergics and incident dementia: a prospective cohort study. *JAMA Intern Med* 2015; 175:401–407.
- Rudolph JL, Salow MJ, Angelini MC, McGlinchey RE. The anticholinergic risk scale and anticholinergic adverse effects in older persons. *Arch Intern Med* 2008; 168:508–513.
- Hilmer SN, Mager DE, Simonsick EM, et al. A drug burden index to define the functional burden of medications in older people. *Arch Intern Med* 2007; 167:781–787.
- Risacher SL, McDonald BC, Tallman EF, et al; Alzheimer's Disease Neuroimaging Initiative. Association between anticholinergic medication use and cognition, brain metabolism, and brain atrophy in cognitively normal older adults. *JAMA Neurol* 2016; 73:721–732.
- Qaseem A, Dallas P, Forciea MA, Starkey M, Denberg TD, Shekelle P; Clinical Guidelines Committee of the American College of Physicians. Nonsurgical management of urinary incontinence in women: a clinical practice guideline from the American College of Physicians. *Ann Intern Med* 2014; 161:429–440.
- Efjestad AS, Molden E, Oksengard AR. Pharmacist-initiated management of antagonistic interactions between anticholinergic drugs and acetyl cholinesterase inhibitors in individuals with dementia. *J Am Geriatr Soc* 2013; 61:1624–1625.
- Kersten H, Molden E, Tolo IK, Skovlund E, Engedal K, Wyller TB. Cognitive effects of reducing anticholinergic drug burden in a frail elderly population: a randomized controlled trial. *J Gerontol A Biol Sci Med Sci* 2013; 68:271–278.
- Curtis LH, Østbye T, Sendersky V, et al. Inappropriate prescribing for elderly Americans in a large outpatient population. *Arch Intern Med* 2004; 164:1621–1625.
- Olfson M, King M, Schoenbaum M. Benzodiazepine use in the United States. *JAMA Psychiatry* 2015; 72:136–142.
- Martin JL, Sainz-Pardo M, Furukawa TA, Martín-Sánchez E, Seoane T, Galán C. Benzodiazepines in generalized anxiety disorder: heterogeneity of outcomes based on a systematic review and meta-analysis of clinical trials. *J Psychopharmacol* 2007; 21:774–782.
- Buscemi N, Vandermeer B, Friesen C, et al. The efficacy and safety of drug treatments for chronic insomnia in adults: a meta-analysis of RCTs. *J Gen Intern Med* 2007; 22:1335–1350.
- Rickels K, Schweizer E, Case WG, Greenblatt DJ. Long-term therapeutic use of benzodiazepines. I. Effects of abrupt discontinuation. *Arch Gen Psychiatry* 1990; 47:899–907.
- Fenton MC, Keyes KM, Martins SS, Hasin DS. The role of a prescription in anxiety medication use, abuse, and dependence. *Am J Psychiatry* 2010; 167:1247–1253.
- Billoti de Gage S, Moride Y, Ducruet T, et al. Benzodiazepine use and risk of Alzheimer's disease: case-control study. *BMJ* 2014; 349:g5205.
- Smink BE, Egberts AC, Lushhof KJ, Uges DR, de Gier JJ. The relationship between benzodiazepine use and traffic accidents: a systemic literature review. *CNS Drugs* 2010; 24:639–653.
- Tinetti ME, Speechley M, Ginter S. Risk factors for falls among elderly persons living in the community. *N Engl J Med* 1988; 319:1701–1707.
- Zint K, Haefeli WE, Glynn RJ, Mogun H, Avorn J, Stürmer T. Impact of drug interactions, dosage, and duration of therapy on the risk of hip fracture associated with benzodiazepine use in older adults. *Pharmacoepidemiol Drug Saf* 2010; 19:1248–1255.
- Briesacher BA, Soumerai SB, Field TS, Fouayzi H, Gurwitz JH. Medicare Part D's exclusion of benzodiazepines and fracture risk in nursing homes. *Arch Intern Med* 2010; 170:693–698.
- Berry SD, Lee Y, Cai S, Dore DD. Nonbenzodiazepine sleep medication use and hip fractures in nursing home residents. *JAMA Intern Med* 2013; 173:754–761.
- American Geriatrics Society. Choosing Wisely. Ten things clinicians and patients should question. www.choosingwisely.org/societies/american-geriatrics-society/. Accessed December 3, 2017.
- Cook JM, Marshall R, Masci C, Coyne JC. Physicians' perspectives on prescribing benzodiazepines for older adults: a qualitative study. *J Gen Intern Med* 2007; 22:303–307.
- Tannenbaum C, Martin P, Tamblyn R, Benedetti A, Ahmed S. Reduction of inappropriate benzodiazepine prescriptions among older adults through direct patient education: the EMPOWER cluster randomized trial. *JAMA Intern Med* 2014; 174:890–898.
- Alexander GC, Gallagher SA, Mascola A, Moloney RM, Stafford RS. Increasing off-label use of antipsychotic medications in the United States, 1995–2008. *Pharmacoepidemiol Drug Saf* 2011; 20:177–184.
- Gill SS, Bronskill SE, Normand SL, et al. Antipsychotic drug use and mortality in older adults with dementia. *Ann Intern Med* 2007; 146:775–786.
- US Food and Drug Administration (FDA). Public health advisory: deaths with antipsychotics in elderly patients with behavioral disturbances. www.fda.gov/Drugs/DrugSafety/PostmarketDrugSafetyInformationforPatientsandProviders/ucm053171.htm. Accessed December 4, 2017.
- Hwang YJ, Dixon SN, Reiss JP, et al. Atypical antipsychotic drugs and the risk for acute kidney injury and other adverse outcomes in older

- adults. *Ann Intern Med* 2014; 161:242–248.
41. Fraser L, Liu K, Naylor KL, et al. Falls and fractures with atypical antipsychotic medication use: a population-based cohort study. *JAMA Intern Med* 2015; 175:450–452.
 42. Gitlin LN, Kales HC, Lyketsos CG. Nonpharmacologic management of behavioral symptoms in dementia. *JAMA* 2012; 308:2020–2029.
 43. Schneider LS, Tariot PN, Dagerman KS, et al; CATIE-AD Study Group. Effectiveness of atypical antipsychotic drugs in patients with Alzheimer's disease. *N Engl J Med* 2006; 355:1525–1538.
 44. Forgacs I, Loganayagam A. Overprescribing proton pump inhibitors. *BMJ* 2008; 336:2–3.
 45. Mazer-Amirshahi M, Mullins PM, van den Anker J, Meltzer A, Pines JM. Rising rates of proton pump inhibitor prescribing in US emergency departments. *Am J Emerg Med* 2014; 32:618–622.
 46. Heidelbaugh JJ, Goldberg KL, Inadomi JM. Magnitude and economic effect of overuse of antisecretory therapy in the ambulatory care setting. *Am J Manag Care* 2010; 16:e228–e324.
 47. Pham CQ, Regal RE, Bostwich TR, Knauf KS. Acid suppressive therapy used on an inpatient internal medicine service. *Ann Pharmacother* 2006; 40:1261–1266.
 48. Kahrilas PJ, Shaheen NJ, Vaezi MF, et al; American Gastroenterological Association. American Gastroenterological Association medical position statement on the management of gastroesophageal reflux disease. *Gastroenterology* 2008; 135:1383–1391.e1–e5.
 49. Howell MD, Novack V, Grgurich P, et al. Iatrogenic gastric acid suppression and the risk of nosocomial *Clostridium difficile* infection. *Arch Intern Med* 2010; 170:784–790.
 50. Gulmez SE, Holm A, Frederiksen H, Jensen TG, Pedersen C, Hallas J. Use of proton pump inhibitors and the risk of community-acquired pneumonia: a population-based case-control study. *Arch Intern Med* 2007; 167:950–955.
 51. Yang YX, Lewis JD, Epstein S, Metz DC. Long-term proton pump inhibitor therapy and risk of hip fracture. *JAMA* 2006; 296:2947–2953.
 52. Lam JR, Schneider JL, Zhao W, Corley DA. Proton pump inhibitor and histamine 2 receptor antagonist use and vitamin B₁₂ deficiency. *JAMA* 2013; 310:2435–2442.
 53. Kuipers EJ, Lundell L, Klinkenberg-Knol EC, et al. Atrophic gastritis and *Helicobacter pylori* infection in patients with reflux esophagitis treated with omeprazole or fundoplication. *N Engl J Med* 1996; 334:1018–1022.
 54. Lazarus B, Chen Y, Wilson FP, et al. Proton pump inhibitor use and the risk of chronic kidney disease. *JAMA Intern Med* 2016; 176:238–246.
 55. Gomm W, von Holt K, Thomé F, et al. Association of proton pump inhibitors with risk of dementia: a pharmacoepidemiological claims data analysis. *JAMA Neurol* 2016; 73:410–416.
 56. Kaltenbach T, Crockett S, Gerson LB. Are lifestyle measures effective in patients with gastroesophageal reflux disease? An evidence-based approach. *Arch Intern Med* 2006; 166:965–971.
 57. Beers MH, Ouslander JG, Rollinger I, Reuben DB, Brooks J, Beck JC. Explicit criteria for determining inappropriate medication use in nursing home residents. *Arch Intern Med* 1991; 151:1825–1832.
 58. American Geriatrics Society 2015 Beers Criteria Update Expert Panel. American Geriatrics Society 2015 updated Beers criteria for potentially inappropriate medication use in older adults. *J Am Geriatr Soc* 2015; 63:2227–2246.
 59. Joint Commission. Sentinel event alert, Issue 35: using medication reconciliation to prevent errors. www.jointcommission.org/sentinel_event_alert_issue_35_using_medication_reconciliation_to_prevent_errors/. Accessed August 18, 2017.
 60. Gallagher P, Ryan C, Byrne S, Kennedy J, O'Mahony D. STOPP (Screening Tool of Older Person's Prescriptions) and START (Screening Tool to Alert doctors to Right Treatment). Consensus validation. *Int J Clin Pharmacol Ther* 2008; 46:72–83.
 61. O'Mahony D, O'Sullivan D, Byrne S, O'Connor MN, Ryan C, Gallagher P. STOPP/START criteria for potentially inappropriate prescribing in older people: version 2. *Age Ageing* 2015; 44:213–218.

ADDRESS: Luke D. Kim, MD, Center for Geriatric Medicine, Medicine Institute, X10, Cleveland Clinic, 9500 Euclid Avenue, Cleveland, OH 44195; kiml2@ccf.org