

# Anxiety and Depression in Chronic Obstructive Pulmonary Disease: Recognition and Management

Abebaw Mengistu Yohannes, PhD; Alan Kaplan, MD; and Nicola A. Hanania, MD, MS

## Introduction

Anxiety and depression are common in patients with chronic obstructive pulmonary disease (COPD), occurring more frequently than in the general population<sup>1-4</sup> or patients with other chronic diseases such as hypertension, diabetes, cancer, or musculoskeletal disorders.<sup>5,6</sup> Their presence is associated with worse outcomes of COPD, and increased morbidity, mortality, disability, and health care expenditure.<sup>6-8</sup> In

spite of this, both anxiety and depression are frequently overlooked and undertreated in patients with COPD,<sup>9</sup> and symptoms of anxiety and depression can overlap significantly, as well as overlap with COPD symptoms.<sup>7,10</sup>

Comorbid depressive disorders that may occur in patients with COPD include major depressive disorder, dysthymias (chronic depressive symptoms of mild severity), and minor depression.<sup>11</sup> Depressive disorders are characterized by feelings of sadness, emptiness, and/or irritability, along with cognitive and somatic symptoms, which have a detrimental effect on the patient's ability to function.<sup>11</sup> Anxiety disorders include generalized anxiety disorder (GAD), phobias, and panic disorders.<sup>11</sup> The main features of anxiety disorders, such as excessive fear and anxiety, may be accompanied by behavioral disturbances related to these symptoms, such as panic attacks and avoidance.<sup>11,12</sup>

The reported prevalence of depression in COPD varies widely between studies, owing to differences in sampling methods and degrees of illness severity used in assessment of depression<sup>6</sup>; rates have been reported to range from 10% to 42% in patients with stable COPD,<sup>6,13</sup> and from 10% to 86% in patients with acute COPD exacerbation.<sup>14</sup> Individuals with severe COPD are twice as likely to develop depression than patients with mild COPD.<sup>10</sup>

Prevalence rates for clinical anxiety in COPD range from 13% to 46% in outpatients and 10% to 55% among inpatients. GAD, panic disorders, and specific phobias are reported most frequently.<sup>15</sup> Patients with COPD are 85% more likely to develop anxiety disorders compared with matched controls without COPD,<sup>4</sup> and panic disorder is reported with a prevalence that is up to 10-fold higher than in the general population.<sup>16</sup>

Global prevalence rates of anxiety and depression are 1.8- and 1.4-fold higher in women than men, respectively<sup>17</sup>; the same gender difference is observed in patients with COPD.<sup>6</sup> The higher prevalence rates of anxiety and depression in women are thought to be a result of sex differences in brain structure, function, and stress responses, as well as differences in exposure to reproductive hormones, social

---

**Abebaw Mengistu Yohannes, PhD**, Azusa Pacific University, Department of Physical Therapy, Azusa, California

**Alan Kaplan, MD**, Family Physician Airways Group of Canada, University of Toronto, Toronto, Ontario, Canada

**Nicola A. Hanania, MD, MS**, Pulmonary and Critical Care Medicine, Baylor College of Medicine, Houston, Texas

## DISCLOSURES

Dr. Yohannes has no conflicts to disclose.

Dr. Kaplan has served as a consultant or speaker for AstraZeneca; Boehringer Ingelheim, GmbH; GlaxoSmithKline; Grifols; Johnson & Johnson; Novartis Pharmaceuticals Corporation; Novo Nordisk Inc.; Paladin Labs Inc.; Pfizer Inc.; Purdue Pharma L.P.; and Trudell Medical International.

Dr. Hanania received honoraria for serving as a consultant or advisory board member for AstraZeneca; Boehringer Ingelheim, GmbH; F. Hoffmann-La Roche Ltd; GlaxoSmithKline; Mylan Laboratories Inc.; Novartis Pharmaceuticals Corporation; Teva Pharmaceutical Industries Ltd.; and Sanofi-Regeneron. His institution has received research grant support from AstraZeneca; Boehringer Ingelheim, GmbH; Cheisi USA, Inc.; F. Hoffmann-La Roche Ltd; and GlaxoSmithKline.

Funding for this article was provided by AstraZeneca LP (Wilmington, DE, USA). Medical writing support was provided by Hannah Burke, BSc, of Core (London, UK) and editorial support was provided by Maryam Vahdat, PGDip, of Core (London, UK), which were funded by AstraZeneca LP (Wilmington, DE, USA), in accordance with Good Publication Practice guidelines (Battisti WP et al. *Ann Intern Med*. 2015;163:461-464. doi: 10.7326/M15-0288).

---

This article is being co-published in *The Journal of Family Practice* and *Cleveland Clinic Journal of Medicine*.  
doi: 10.3949/ccjm.85.s1.03

constraints, and experiences between women and men.<sup>18</sup> However, psychologic comorbidity is an issue for both men and women with COPD, so it is important that clinicians are vigilant in recognizing anxiety and depression in both sexes, and are careful not to underestimate the burden in the male patient population.

It is also important to note that depression and anxiety often occur simultaneously in patients with COPD, with prevalence estimates of 26% to 43%.<sup>9,19,20</sup> COPD patients with both depression and anxiety are at a heightened risk of suicidal ideation, increased physical disability, and chronic depressive symptoms versus those with either disorder alone.<sup>10,15</sup> It is therefore important that comorbid anxiety and depression is not overlooked in patients with COPD.

Ensuring that anxiety and depression are recognized and treated effectively in patients with COPD is essential for optimizing outcomes. Primary care practitioners are well placed to diagnose anxiety and depression, and to ensure these conditions are suitably managed alongside treatments of COPD.

### Potential mechanisms of anxiety and depression in COPD

Growing evidence suggests that the relationship between mood disorders—particularly depression—and COPD is bidirectional, meaning that mood disorders adversely impact prognosis in COPD, whereas COPD increases the risk of developing depression.<sup>21</sup> For example, in a study of 60 stable patients with COPD, elevated dyspnea and reduced exercise capacity were the predominant mechanisms leading to anxiety and depression symptoms associated with the condition.<sup>22</sup> In addition, the risk of new-onset depression was increased in COPD patients with moderate-to-severe dyspnea in a 3-year follow-up study.<sup>23</sup> Conversely, depression has been shown to be a significant risk factor for disabling dyspnea (modified Medical Research Council score  $\geq 2$ ) in patients with COPD.<sup>24</sup>

COPD can lead to feelings of hopelessness, social isolation, reduced physical functioning, and sedentary lifestyle, all of which are associated with an increased level of depressive symptoms.<sup>25</sup> Similarly, inadequate social support increases the risk of anxiety in patients with COPD.<sup>26</sup> Therefore, ensuring that patients with COPD have high-quality support is very important for reducing anxiety and depressive symptoms.<sup>27</sup>

The exact mechanisms for the association between mood disorders and COPD remain unclear.<sup>7,10</sup> Research to date indicates that the relationship between depression and impaired pulmonary function may be partly mediated by chronic inflammation<sup>7,10</sup>; systemic inflammation has been

associated with other comorbidities of COPD (eg, muscle wasting and osteoporosis),<sup>28</sup> and emerging data appear to show that proinflammatory cytokines partly mediate the association between depressive symptoms and pulmonary function.<sup>29</sup> Smoking and hypoxemia may also influence the prevalence of depression in COPD, but symptom severity and impaired quality of life remain the most important determinants.<sup>6,30</sup>

Clinical studies have demonstrated that a number of patient-related factors, including female gender, younger age, current smoking, greater severity of airflow limitation, and lower socioeconomic status, are associated with a higher prevalence and/or increased risk of depression and/or anxiety in COPD.<sup>3,4,30,31</sup> Frequent episodes of rehospitalization, and comorbidities such as hypertension, arthritis, cancer, and heart disease, have been found to increase the risk of anxiety and depression in patients with COPD.<sup>3,32</sup> Risk of anxiety has been shown to increase with greater dyspnea severity.<sup>4</sup> Pain, a frequently overlooked symptom in COPD, has been shown to be associated with symptoms of both anxiety and depression in patients with COPD.<sup>33</sup> This is driven by worsened quality of life and sleep quality, decreased physical activity, and an increased fear of movement that occur as a result of pain.<sup>34</sup>

### The impact of anxiety and depression in COPD

Comorbid anxiety and depression have a significant detrimental impact on morbidity and mortality in patients with COPD. Both disorders have been associated with an increased risk of death in COPD.<sup>13,35-37</sup> Indeed, of 12 comorbidities proposed to be predictors of mortality in a cohort of 187 female outpatients with COPD, anxiety was associated with the highest risk of death.<sup>35,36</sup>

In addition, patients with COPD and anxiety and/or depression have a higher risk of COPD exacerbations,<sup>4,8,23,36,38-40</sup> hospitalization,<sup>41,42</sup> rehospitalization,<sup>14,36,43</sup> longer hospital stays,<sup>37,41,44</sup> and mortality after exacerbations,<sup>14,36,41</sup> compared with patients without these comorbidities. Patients with COPD who have elevated anxiety symptoms also often experience their first hospitalization earlier in the natural course of COPD than those without anxiety.<sup>36</sup>

Psychologic comorbidities are also associated with worse lung function, dyspnea, and respiratory symptom burden in patients with COPD.<sup>37,40</sup> Patients with COPD and anxiety are more likely to experience greater dyspnea at an earlier stage of disease than those without anxiety.<sup>36</sup> Persistent smoking at 6 months after hospitalization for an acute exacerbation of COPD is also more likely to be seen in patients with depression.<sup>37</sup>

Patient-centered outcomes are worse in COPD patients with mood disorders. Both anxiety and depression have been

shown to correlate with significantly reduced health-related quality of life (HRQoL), poorer physical health status, functional limitations, and reduced exercise capacity.<sup>4,23,37,40,45</sup> The presence of either anxiety or depression at baseline has been shown to correlate with reduced HRQoL at 1-year follow-up, but depression appears to be the stronger predictor of low future HRQoL than anxiety.<sup>45</sup>

Additionally, mood disorders—particularly depression—reduce physical activity in patients with COPD.<sup>46,47</sup> Emotional responses to COPD symptoms, such as dyspnea, can further decrease activity and worsen deconditioning, resulting in a downward spiral of reduced inactivity, social isolation, fear, anxiety, and depression.<sup>48</sup>

COPD patients with any comorbidity exhibit lower rates of medication adherence than those without comorbidities.<sup>49-51</sup> Clinical studies have demonstrated that anxiety and depression are significant predictors of poor adherence to COPD interventions, including pulmonary rehabilitation (PR).<sup>51-55</sup> Nonadherence to COPD therapies is associated with poor clinical outcomes, including higher hospitalization rates and increased emergency department visits, and increased costs.<sup>56,57</sup> Health care expenditure, in terms of both specific COPD-related costs and general “all-cause” costs, is significantly higher in COPD patients with anxiety and/or depression than in those without.<sup>8</sup>

### Diagnosis of anxiety and depression in patients with COPD

The underdiagnosis and undertreatment of anxiety and depression in this population is common and can adversely affect patient outcomes.<sup>6,7,9,10,58</sup> Hence, it is crucial that anxiety and depression are identified and more effectively managed in clinical practice.<sup>10</sup>

Primary care practitioners are the main point of contact for many patients with COPD,<sup>6,59,60</sup> and so can play a key role in screening for and early identification of anxiety and depression. However, detection of mood disorders by primary care practitioners is challenging for several reasons. These include the lack of a standardized approach in diagnosis, and inadequate knowledge or confidence in assessing psychological status (particularly given the number of strategies available for screening patients for mood disorders),<sup>6</sup> as well as factors associated with time constraints, such as competing agendas, duration of visits, and high patient load.<sup>6,61</sup> Furthermore, system-level barriers, such as lack of electronic medical records and adequate health insurance, as well as any communication gaps between primary care and mental health care, may hinder the detection and management of anxiety and depression.<sup>6</sup> In addition, patients themselves may have a limited understanding of these comorbidities, or

may be hesitant to discuss symptoms of anxiety or depression with their primary care practitioner owing to stigma around mental illness.<sup>6</sup>

Patients with COPD should be screened and assessed for anxiety and depression, and the United States Preventive Services Task Force recommends that clinicians screen for depression in all adults.<sup>6,62</sup> There are several validated screening tools suitable for clinical use:

- Anxiety Inventory for Respiratory (AIR) Disease scale: a brief, easy-to-use tool for screening and measuring anxiety in COPD.<sup>63,64</sup> It is a self-administered scale, and takes approximately 2 minutes to complete. The AIR scale is responsive to PR.<sup>64</sup>
- COPD Anxiety Questionnaire (CAF): a reliable tool for early identification of COPD-related anxiety.<sup>65</sup>
- Primary Care Evaluation of Mental Disorders (PRIME-MD) Patient Health Questionnaire (PHQ; available at: <http://www.phqscreener.com/select-screener/>): the PRIME-MD comprises 26 yes/no questions on the 5 most common psychiatric disorders, including depression and anxiety.<sup>66,67</sup> This is not a diagnostic tool, but a high number of positive responses from a patient in any given module indicates that they require further clinical evaluation.
- PHQ-2 and PHQ-9 (**TABLE 1**; PHQ-9 available at <http://www.phqscreener.com/select-screener/>): widely-used self-administered 2- and 9-item versions of the PRIME-MD, specific to depression; similarly, the 3-item PHQ-3 is available for anxiety assessment (**TABLE 2**).<sup>6,67,68</sup> In a study investigating tools used by family physicians in England to assess depression, over 75% used PHQ-9.<sup>69</sup>
- Generalized Anxiety Disorder 7-item (GAD-7) scale: an efficient, self-report scale that scores 7 common anxiety symptoms and can be used for screening and severity assessment of GAD in clinical practice.<sup>70</sup>
- Hospital Anxiety and Depression Scale (HADS) and General Health Questionnaire-version 20 (GHQ-20): both can be used to screen for psychologic distress in patients with COPD.<sup>71</sup>
- The Beck Anxiety Inventory (BAI) and Beck Depression Inventory (BDI): two 21-item self-report questionnaires that are widely used in the United States to evaluate anxiety and depression.<sup>72</sup>

In addition to specific anxiety and depression questionnaires (**TABLES 1 and 2**), more general COPD assessments tools, such as the COPD Assessment Test and the COPD Clinical Questionnaire, also incorporate questions that may be indicative of symptoms of these comorbidities in patients with COPD.<sup>73</sup>

**TABLE 1 PHQ-2 and PHQ-9 screening questionnaires for depression<sup>6,67</sup>**

PHQ-2	In the past month, have you been bothered a lot by: <ol style="list-style-type: none"> <li>1. Little interest or pleasure in doing things?</li> <li>2. Feeling down, depressed, or hopeless?</li> </ol>
PHQ-9	Over the past 2 weeks, how often have you been bothered by any of the following problems (not at all, for several days, for more than half the days, or nearly every day)? <ol style="list-style-type: none"> <li>1. Little interest or pleasure in doing things</li> <li>2. Feeling down, depressed, or hopeless</li> <li>3. Trouble falling or staying asleep, or sleeping too much</li> <li>4. Feeling tired or having little energy</li> <li>5. Poor appetite or overeating</li> <li>6. Feeling bad about yourself – or that you are a failure or have let yourself or your family down</li> <li>7. Trouble concentrating on things, such as reading the newspaper or watching television</li> <li>8. Moving or speaking so slowly that other people could have noticed? Or the opposite – being so fidgety or restless that you have been moving around a lot more than usual</li> <li>9. Thoughts that you would be better off dead or hurting yourself in some way</li> </ol>

Reprinted from *Chest*,134(4 Suppl):43S–56S. Maurer J, Rebbapragada V, Borson S, et al. ACCP Workshop Panel on Anxiety and Depression in COPD. Anxiety and depression in COPD: current understanding, unanswered questions, and research needs. Copyright 2008, with permission from Elsevier. PHQ-9 questionnaire developed by Drs. Robert L. Spitzer, Janet B.W. Williams, Kurt Kroenke and colleagues, with an educational grant from Pfizer Inc.

### Management of anxiety and depression in COPD

Even though anxiety and depression are among the most common and burdensome comorbid conditions in COPD, less than one-third of patients with these comorbidities receive effective intervention.<sup>6,10</sup> Primary care providers have an excellent opportunity to impact this care gap.

It is important that all health care professionals involved in the care of patients with COPD are vigilant for anxiety and depressive symptoms, as well as the possibility of a major anxiety or depressive disorder. Communication with other multidisciplinary team members is central to ensuring appropriate psychiatric treatment in patients with COPD, particularly sharing key information about medication history, warning signs of depression and anxiety, and any indication of suicide ideation.<sup>74</sup> Referral to palliative care teams can also help to manage these psychological comorbidities in patients with severe COPD at advanced stages.<sup>75</sup>

As in non-COPD patients, comorbid depression and anxiety may be treated with nonpharmacologic and/or pharmacologic interventions (**FIGURE 1**).<sup>76</sup>

### Nonpharmacologic interventions

Evidence to date suggests that nonpharmacologic interventions such as behavioral therapy are as effective as antidepressants, and may be preferred by patients with mood disorders.<sup>12</sup>


**TABLE 2 PHQ-3 screening questionnaire for anxiety<sup>6,67</sup>**

PHQ-3	In the past month, have you been bothered a lot by: <ol style="list-style-type: none"> <li>1. ‘Nerves,’ or feeling anxious or ‘on edge’?</li> <li>2. Worrying about a lot of different things?</li> </ol> During the past month: <ol style="list-style-type: none"> <li>3. Have you had an anxiety attack (ie, suddenly feeling fear or panic)?</li> </ol>
-------	--

Reprinted from *Chest*,134(4 Suppl):43S–56S. Maurer J, Rebbapragada V, Borson S, et al. ACCP Workshop Panel on Anxiety and Depression in COPD. Anxiety and depression in COPD: current understanding, unanswered questions, and research needs. Copyright 2008, with permission from Elsevier.

Cognitive behavioral therapy (CBT), which is typically administered by psychologists/psychiatrists, may be effective in treating COPD-related anxiety and depression, especially in conjunction with exercise and education.<sup>12,76,77</sup> Individualized or group CBT is the treatment of choice for addressing thinking patterns that contribute to anxiety and depression to change a patient’s behavior and emotional state.<sup>76</sup> PR programs involve several components,

**FIGURE 1** Recommendations for the treatment of psychiatric symptoms in patients with COPD<sup>76</sup>



Republished with permission of Dove Medical Press Limited. Strategies to improve anxiety and depression in patients with COPD: a mental health perspective. Tselebis A et al. *Neuropsychiatric Disease and Treatment*. 2016;12:297–328; permission conveyed through Copyright Clearance Center, Inc.

including aerobic exercise, lung function training, and psycho-education.<sup>62,76</sup> PR is suitable for most patients with COPD, and provides multiple benefits, including reduced hospitalizations in patients who have had a recent exacerbation, and improved dyspnea, exercise tolerance, and health status in patients with stable disease,<sup>62</sup> as well as clinically and statistically significant improvements in depression and anxiety, irrespective of age.<sup>7,78,79</sup> Exercise-based forms of PR appear to be the most effective for reducing mood symptoms,<sup>12,76</sup> and incorporating psychotherapy may also improve psychologic outcomes.<sup>80</sup> Stress reduction (relaxation) therapy aims to reduce anxiety-related physiologic changes, and includes a variety of techniques (eg, breathing exercises, sequential muscle relaxation, hypnosis, mindfulness meditation), some of which may be included in PR or used alongside other treatments (eg, CBT).<sup>76</sup> Limited data indicate that such therapy may be beneficial for reducing anxiety and depression, as well as respiratory symptoms and dyspnea, in patients with COPD.<sup>12,76</sup>

Self-management techniques improve clinical outcomes in patients with COPD, but data on the management of depression or anxiety are inconclusive.<sup>7,12</sup> A minimal,

home-based, nurse-led, psycho-educational intervention was designed to encourage more open-ended, descriptive discussions of thoughts, emotions, behaviors, and bodily sensations in patients with COPD.<sup>81</sup> The intervention, which involved nurses attending a 1-hour face-to-face session in the patients' homes with a 15-minute telephone "booster" session 2 weeks later, helped patients with advanced COPD to self-manage their condition and provide relief from anxiety.<sup>81,82</sup> However, it should be noted that there is currently a lack of high-quality data evaluating psychologic interventions in the COPD population.<sup>83</sup>

In addition, it is important that caregivers are supported in the management of patients with COPD and comorbid anxiety and/or depression; areas in which caregivers can be assisted in their role may include disease education and counseling, where appropriate.<sup>84</sup>

Given that smoking cessation is a key recommendation for patients with COPD,<sup>44,62</sup> practitioners should be aware that patients with comorbid depression and anxiety may experience greater difficulty in smoking cessation, and worsened mood during nicotine withdrawal.<sup>44</sup> Clinicians should therefore carefully monitor current smokers with

COPD and comorbid depression/anxiety (using the tools described previously<sup>63,68,70,71</sup>) when they are attempting to quit smoking.

### Pharmacologic interventions

Pharmacologic therapy of anxiety and depression has so far only been investigated in patients with COPD in small studies.<sup>76</sup> However, the available evidence does not indicate that COPD patients with anxiety and depression should be managed any differently from individuals without COPD.<sup>62</sup> As such, pharmacologic interventions are particularly important for patients with acute or severe anxiety or depression.

Antidepressant agents are categorized according to their mechanism of action, and most commonly include selective serotonin-reuptake inhibitors (SSRIs), selective norepinephrine-reuptake inhibitors, bupropion (a norepinephrine- and dopamine-reuptake inhibitor; also approved for smoking cessation<sup>85</sup>), and mirtazapine (a norepinephrine and serotonin modulator), among others.<sup>86</sup> SSRIs are the current first-line drug treatment for depression, and have been shown to significantly improve depression and anxiety in patients with COPD in some, but not all, trials published to date.<sup>76</sup> However, it is important to note that a diagnosis of bipolar disorder must be ruled out before initiating standard antidepressant therapy.<sup>87</sup> In addition to antidepressants, atypical antipsychotics have also been shown to be useful for treating anxiety, either as monotherapy or combination therapy, and possibly as an adjunctive therapy for the management of depression.<sup>88,89</sup>

Primary care practitioners can refer to existing guidelines on the management of anxiety and depression in patients with COPD,<sup>86,90</sup> while taking certain factors into consideration. Any pharmacologic management strategy for the treatment of COPD may increase the risk of drug–drug or drug–disease interactions.<sup>76</sup> For example, it is important to avoid medications that cause respiratory depression (eg, benzodiazepines [unless used with extreme caution], particularly in patients who are already CO<sub>2</sub> retainers) or sedation; chosen drugs should have minimal other adverse effects.<sup>76</sup> Moreover, SSRIs may also be associated with troublesome adverse effects during treatment initiation, such as gastrointestinal upset, headache, tremor, psychomotor activation, and sedation<sup>76</sup>; in addition, dry mouth is an adverse effect associated with both SSRI treatment and several inhaled therapies, so may be particularly problematic in patients with COPD.<sup>91,92</sup> Currently, data are particularly scarce for the management of anxiety in patients with COPD, with inconclusive or contradictory findings reported for SSRIs, azapirones (including buspirone), and tricyclic antidepressants.<sup>76</sup>

In addition to monitoring adherence to COPD therapies, primary care practitioners should carefully monitor patients treated with antidepressants and anxiolytics for adherence. A meta-analysis of 18,245 individuals with chronic diseases showed that depressed patients had a 76% significantly higher risk of nonadherence to medication compared with those without depressive symptoms.<sup>93</sup>

Targeting dyspnea is key to the management of anxiety and depression in COPD, as breathlessness is frequently associated with the onset of both comorbidities.<sup>21,22</sup> Therapeutic approaches to alleviating dyspnea include PR, optimizing respiratory mechanics and muscle function (with bronchodilator therapy), and reducing central neural drive to respiratory muscles with supplemental oxygen or opioid medication.<sup>94</sup>

Although bronchodilator therapy for COPD has not been shown to have significant direct effects on depression or anxiety,<sup>95</sup> it can be assumed that the beneficial effects on dyspnea are likely to alleviate associated emotional and mood symptoms.

Further research into effective screening, diagnosis, and management of comorbid anxiety and depressive disorders in COPD is warranted, including evaluation of a broad range of nonpharmacologic and drug-based interventions, alone and in combination.<sup>76</sup>

### Conclusions

Anxiety and depression are common, yet frequently overlooked, comorbidities in COPD. The impact of these psychological comorbidities is significant; their consequences are evident in morbidity and mortality data, as well as in patient-reported outcomes. As key points of contact for patients with COPD, it is essential that primary care practitioners are vigilant in monitoring for anxiety and depression in their patients with COPD, making the most of the available tools that can support them in doing so, and maintain an ongoing line of communication with other members of the multidisciplinary team. Treatment of anxiety and depression in COPD should adopt a holistic approach that incorporates both nonpharmacologic and pharmacologic interventions. However, the impact of effective screening, diagnosis, and management of anxiety and depression on COPD burden in patients requires further investigation. ●

### REFERENCES

1. Chaudhary SC, Nanda S, Tripathi A, et al. Prevalence of psychiatric comorbidities in chronic obstructive pulmonary disease patients. *Lung India*. 2016;33(2):174-178.
2. Zhang MW, Ho RC, Cheung MW, Fu E, Mak A. Prevalence of depressive symptoms in patients with chronic obstructive pulmonary disease: a systematic review, meta-analysis and meta-regression. *Gen Hosp Psychiatry*. 2011;33(3):217-223.
3. Tsai TY, Livneh H, Lu MC, Tsai PY, Chen PC, Sung FC. Increased risk and related

- factors of depression among patients with COPD: a population-based cohort study. *BMC Public Health*. 2013;13:976.
4. Eisner MD, Blanc PD, Yelin EH, et al. Influence of anxiety on health outcomes in COPD. *Thorax*. 2010;65(3):229-234.
  5. Marsh S, Guck TP. Anxiety and depression: easing the burden in COPD patients. *J Fam Pract*. 2016;65(4):246-256.
  6. Maurer J, Rebbapragada V, Borson S, et al; ACCP Workshop Panel on Anxiety and Depression in COPD. Anxiety and depression in COPD: current understanding, unanswered questions, and research needs. *Chest*. 2008;134(4 Suppl):43S-56S.
  7. Pumar MI, Gray CR, Walsh JR, Yang IA, Rolls TA, Ward DL. Anxiety and depression—important psychological comorbidities of COPD. *J Thorac Dis*. 2014;6(11):1615-1631.
  8. Dalal AA, Shah M, Lunacek O, Hanania NA. Clinical and economic burden of depression/anxiety in chronic obstructive pulmonary disease patients within a managed care population. *COPD*. 2011;8(4):293-299.
  9. Kunik ME, Roundy K, Veazey C, et al. Surprisingly high prevalence of anxiety and depression in chronic breathing disorders. *Chest*. 2005;127(4):1205-1211.
  10. Yohannes AM, Alexopoulos GS. Depression and anxiety in patients with COPD. *Eur Respir Rev*. 2014;23(133):345-349.
  11. American Psychiatric Association. *Depressive Disorders*. Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition. Arlington, VA: American Psychiatric Association; 2013.
  12. Panagioti M, Scott C, Blakemore A, Coventry PA. Overview of the prevalence, impact, and management of depression and anxiety in chronic obstructive pulmonary disease. *Int J Chron Obstruct Pulmon Dis*. 2014;9:1289-1306.
  13. Laforest L, Roche N, Devouassoux G, et al. Frequency of comorbidities in chronic obstructive pulmonary disease, and impact on all-cause mortality: a population-based cohort study. *Respir Med*. 2016;117:33-39.
  14. Lecheler L, Richter M, Franzen DP, et al. The frequent and underrecognised co-occurrence of acute exacerbated COPD and depression warrants screening: a systematic review. *Eur Respir Rev*. 2017;26(144):pii: 170026.
  15. Willgoss TG, Yohannes AM. Anxiety disorders in patients with COPD: a systematic review. *Respir Care*. 2013;58(5):858-866.
  16. Livermore N, Sharpe L, McKenzie D. Panic attacks and panic disorder in chronic obstructive pulmonary disease: a cognitive behavioral perspective. *Respir Med*. 2010;104(9):1246-1253.
  17. World Health Organization. *Depression and other common mental disorders: global health estimates*. Geneva, Switzerland; World Health Organization: 2017.
  18. Altemus M, Sarvaiya N, Neill Epperson C. Sex differences in anxiety and depression clinical perspectives. *Front Neuroendocrinol*. 2014;35(3):320-330.
  19. Biswas D, Mukherjee S, Chakraborty R, et al. Occurrence of anxiety and depression among stable COPD patients and its impact on functional capability. *J Clin Diagn Res*. 2017;11(2):OC24-OC27.
  20. Yohannes AM, Baldwin R, Connolly MJ. Depression and anxiety in elderly outpatients with chronic obstructive pulmonary disease: prevalence, and validation of the BASDEC screening questionnaire. *Int J Geriatr Psychiatry*. 2000;15(12):1090-1096.
  21. Atlantis E, Fahey P, Cochrane B, Smith S. Bidirectional associations between clinically relevant depression or anxiety and COPD: a systematic review and meta-analysis. *Chest*. 2013;144(3):766-777.
  22. Tetikurt C, Ozdemir I, Tetikurt S, Yilmaz N, Ertan T, Bayar N. Anxiety and depression in COPD patients and correlation with sputum and BAL cytology. *Multidiscip Respir Med*. 2011;6(4):226-231.
  23. Yohannes AM, Müllerová H, Hanania NA, et al. Long-term course of depression trajectories in patients with COPD: a 3-year follow-up analysis of the evaluation of COPD longitudinally to identify predictive surrogate endpoints cohort. *Chest*. 2016;149(4):916-926.
  24. Sundh J, Ekström M. Persistent disabling breathlessness in chronic obstructive pulmonary disease. *Int J Chron Obstruct Pulmon Dis*. 2016;11:2805-2812.
  25. Kirkil G, Deveci F, Deveci SE, Atmaca M. Anxiety and depression symptoms in patients with chronic obstructive pulmonary disease. *Bulletin Clin Psychopharmacol*. 2015;25(2):151-161.
  26. Fuller-Thomson E, Lacombe-Duncan A. Understanding the association between chronic obstructive pulmonary disease and current anxiety: a population-based study. *COPD*. 2016;13(5):622-631.
  27. Yohannes AM. Is it quality or quantity of social support needed for patients with chronic medical illness? *J Psychosom Res*. 2013;74(2):87-88.
  28. Fabbri LM, Luppi F, Beghe B, Rabe KF. The multiple components of COPD. In: Hanania NA, Sharafkhaneh A, eds. *COPD: a guide to diagnosis and clinical management*. Humana Press; 2010.
  29. Lu Y, Feng L, Feng L, Nyunt MS, Yap KB, Ng TP. Systemic inflammation, depression and obstructive pulmonary function: a population-based study. *Respir Res*. 2013;14:53.
  30. Hanania NA, Müllerová H, Locantore NW, et al. Determinants of depression in the ECLIPSE chronic obstructive pulmonary disease cohort. *Am J Respir Crit Care Med*. 2011;183(5):604-611.
  31. Zhang Q, Liao J, Liao X, et al. Disease knowledge level is a noteworthy risk factor of anxiety and depression in patients with chronic obstructive pulmonary disease: a cross-sectional study. *BMC Pulm Med*. 2014;14:92.
  32. Jose AK, Chelangara DP, Shaji KS. Factors associated with anxiety and depression in chronic obstructive pulmonary disease. *Int J Res Med Sci*. 2016;4(4):1074-1079.
  33. van Dam van Isselt EF, Groenewegen-Sipkema KH, Spruit-van Eijk M, et al. Pain in patients with COPD: a systematic review and meta-analysis. *BMJ Open*. 2014;4(9):e005898.
  34. Lee AL, Harrison SL, Goldstein RS, Brooks D. Pain and its clinical associations in individuals with COPD: a systematic review. *Chest*. 2015;147(5):1246-1258.
  35. Divo M, Cote C, de Torres JP, et al; BODE Collaborative Group. Comorbidities and risk of mortality in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med*. 2012;186(2):155-161.
  36. Hillas G, Perlikos F, Tsiligianni I, Tzanakis N. Managing comorbidities in COPD. *Int J Chron Obstruct Pulmon Dis*. 2015;10:95-109.
  37. Ng TP, Niti M, Tan WC, Cao Z, Ong KC, Eng P. Depressive symptoms and chronic obstructive pulmonary disease: effect on mortality, hospital readmission, symptom burden, functional status, and quality of life. *Arch Intern Med*. 2007;167(1):60-67.
  38. Montserrat-Capdevila J, Godoy P, Marsal JR, et al. Overview of the impact of depression and anxiety in chronic obstructive pulmonary disease. *Lung*. 2017;195(1):77-85.
  39. Laurin C, Moullec G, Bacon SL, Lavoie KL. Impact of anxiety and depression on chronic obstructive pulmonary disease exacerbation risk. *Am J Respir Crit Care Med*. 2012;185(9):918-923.
  40. Martínez Rivera C, Costan Galicia J, Alcázar Navarrete B, et al. Factors associated with depression in COPD: a multicenter study. *Lung*. 2016;194(3):335-343.
  41. Pooler A, Beech R. Examining the relationship between anxiety and depression and exacerbations of COPD which result in hospital admission: a systematic review. *Int J Chron Obstruct Pulmon Dis*. 2014;9:315-330.
  42. Dahlén I, Janson C. Anxiety and depression are related to the outcome of emergency treatment in patients with obstructive pulmonary disease. *Chest*. 2002;122(5):1633-1637.
  43. Gudmundsson G, Gislason T, Janson C, et al. Risk factors for rehospitalisation in COPD: role of health status, anxiety and depression. *Eur Respir J*. 2005;26(3):414-419.
  44. Mikkelsen RL, Middelboe T, Pisinger C, Stage KB. Anxiety and depression in patients with chronic obstructive pulmonary disease (COPD). A review. *Nord J Psychiatry*. 2004;58(1):65-70.
  45. Blakemore A, Dickens C, Guthrie E, et al. Depression and anxiety predict health-related quality of life in chronic obstructive pulmonary disease: systematic review and meta-analysis. *Int J Chron Obstruct Pulmon Dis*. 2014;9:501-512.
  46. Dueñas-Espín I, Demeyer H, Gimeno-Santos E, et al. Depression symptoms reduce physical activity in COPD patients: a prospective multicenter study. *Int J Chron Obstruct Pulmon Dis*. 2016;11:1287-1295.
  47. Lee SH, Kim KU, Lee H, Kim YS, Lee MK, Park HK. Factors associated with low-level physical activity in elderly patients with chronic obstructive pulmonary disease [published online ahead of print June 7, 2017]. *Korean J Intern Med*. 2017;doi: 10.3904/kjim.2016.090.
  48. Hill K, Geist R, Goldstein RS, Lacasse Y. Anxiety and depression in end-stage COPD. *Eur Respir J*. 2008;31(3):667-677.
  49. George J, Kong DC, Thoman R, Stewart K. Factors associated with medication non-adherence in patients with COPD. *Chest*. 2005;128(5):3198-3204.
  50. Morrison D, Agur K, Mercer S, Eiras A, González-Montalvo JI, Gruffydd-Jones K. Managing multimorbidity in primary care in patients with chronic respiratory conditions. *NPJ Prim Care Respir Med*. 2016;26:16043.
  51. Khmour MR, Hawwa AF, Kidney JC, Smyth BM, McElroy JC. Potential risk factors for medication non-adherence in patients with chronic obstructive pulmonary disease (COPD). *Eur J Clin Pharmacol*. 2012;68(10):1365-1373.
  52. Busch AM, Scott-Sheldon LA, Pierce J, et al. Depressed mood predicts pulmonary rehabilitation completion among women, but not men. *Respir Med*. 2014;108(7):1007-1013.
  53. Heerema-Poelman A, Stuive I, Wempe JB. Adherence to a maintenance exercise program 1 year after pulmonary rehabilitation: what are the predictors of dropout? *J Cardiopulm Rehabil Prev*. 2013;33(6):419-426.
  54. DiMatteo MR, Lepper HS, Croghan TW. Depression is a risk factor for noncompliance with medical treatment: meta-analysis of the effects of anxiety and depression on patient adherence. *Arch Intern Med*. 2000;160(14):2101-2107.
  55. Fan VS, Giardino ND, Blough DK, Kaplan RM, Ramsey SD; Nett Research Group. Costs of pulmonary rehabilitation and predictors of adherence in the National Emphysema Treatment Trial. *COPD*. 2008;5(2):105-116.
  56. Bourbeau J, Bartlett SJ. Patient adherence in COPD. *Thorax*. 2008;63(9):831-838.
  57. van Boven JF, Chavannes NH, van der Molen T, Rutten-van Mölken MP, Postma MJ, Vegter S. Clinical and economic impact of non-adherence in COPD: a systematic review. *Respir Med*. 2014;108(1):103-113.
  58. Dury R. COPD and emotional distress: not always noticed and therefore untreated. *Br J Community Nurs*. 2016;21(3):138-141.
  59. Price D, Crockett A, Arne M, et al. Spirometry in primary care case-identification, diagnosis and management of COPD. *Prim Care Respir J*. 2009;18(3):216-223.
  60. van Boven JF, Ryan D, Eakin MN, Canonica GW, Barot A, Foster JM; Respiratory Effectiveness Group. Enhancing respiratory medication adherence: the role of health care professionals and cost-effectiveness considerations. *J Allergy Clin Immunol Pract*. 2016;4(5):835-846.

61. Wittchen HU, Mühlig S, Beesdo K. Mental disorders in primary care. *Dialogues Clin Neurosci*. 2003;5(2):115-128.
62. Global Initiative for Chronic Obstructive Lung Disease. GOLD 2017 Global Strategy for the Diagnosis, Management and Prevention of COPD. <http://goldcopd.org/gold-2017-global-strategy-diagnosis-management-prevention-copd/>. Accessed June 2017.
63. Willgoss TG, Goldbart J, Fatoye F, Yohannes AM. The development and validation of the anxiety inventory for respiratory disease. *Chest*. 2013;144(5):1587-1596.
64. Yohannes AM, Dryden S, Hanania NA. The responsiveness of the anxiety inventory for respiratory disease scale following pulmonary rehabilitation. *Chest*. 2016;150(1):188-195.
65. Kühl K, Kuhn C, Kenn K, Rief W. [The COPD-Anxiety-Questionnaire (CAF): a new instrument to assess illness specific anxiety in COPD patients]. *Psychother Psychosom Med Psychol*. 2011;61(1):e1-e9. German.
66. Tamburrino MB, Lynch DJ, Nagel RW, Smith MK. Primary care evaluation of mental disorders (PRIME-MD) screening for minor depressive disorder in primary care. *Prim Care Companion J Clin Psychiatry*. 2009;11(6):339-343.
67. Spitzer RL, Kroenke K, Williams JB. Validation and utility of a self-report version of PRIME-MD: the PHQ primary care study. Primary Care Evaluation of Mental Disorders. Patient Health Questionnaire. *JAMA*. 1999;282(18):1737-1744.
68. Arroll B, Goodyear-Smith F, Crengle S, et al. Validation of PHQ-2 and PHQ-9 to screen for major depression in the primary care population. *Ann Fam Med*. 2010;8(4):348-353.
69. Yohannes AM, Hann M, Sibbald B. The management of depressive symptoms in patients with COPD: a postal survey of general practitioners. *Prim Health Care Res Dev*. 2011;12(3):237-244.
70. Spitzer RL, Kroenke K, Williams JB, Löwe B. A brief measure for assessing generalized anxiety disorder: the GAD-7. *Arch Intern Med*. 2006;166(10):1092-1097.
71. Bratås O, Grønning K, Forbord T. Psychometric properties of the Hospital Anxiety and Depression Scale and The General Health Questionnaire-20 in COPD inpatients. *Scand J Caring Sci*. 2014;28(2):413-420.
72. Lovibond PF, Lovibond SH. The structure of negative emotional states: comparison of the Depression Anxiety Stress Scales (DASS) with the Beck Depression and Anxiety Inventories. *Behav Res Ther*. 1995;33(3):335-343.
73. Sundh J, Stållberg B, Lisspers K, Kämpe M, Janson C, Montgomery S. Comparison of the COPD Assessment Test (CAT) and the Clinical COPD Questionnaire (CCQ) in a clinical population. *COPD*. 2016;13(1):57-65.
74. Cantor L, Jacobson R. COPD: How to manage comorbid depression and anxiety. *Curr Psychiatry*. 2003;2(11):45-54.
75. Yohannes AM. Palliative care provision for patients with chronic obstructive pulmonary disease. *Health Qual Life Outcomes*. 2007;5:17.
76. Tselebis A, Pachi A, Ilias I, et al. Strategies to improve anxiety and depression in patients with COPD: a mental health perspective. *Neuropsychiatr Dis Treat*. 2016;12:297-328.
77. Doyle C, Bhar S, Fearn M, et al. The impact of telephone-delivered cognitive behaviour therapy and befriending on mood disorders in people with chronic obstructive pulmonary disease: a randomized controlled trial. *Br J Health Psychol*. 2017;22(3):542-556.
78. Alsarareh FA, Aloush SA. Does pulmonary rehabilitation alleviate depression in older patients with chronic obstructive pulmonary disease. *Saudi Med J*. 2017;38(5):491-496.
79. Bennett D, Bowen B, McCarthy P, Subramaniam A, O'Connor M, Henry MT. Outcomes of pulmonary rehabilitation for COPD in older patients: a comparative study. *COPD*. 2017;14(2):170-175.
80. Smith SM, Sonogo S, Ketcheson L, Larson JL. A review of the effectiveness of psychological interventions used for anxiety and depression in chronic obstructive pulmonary disease. *BMJ Open Respir Res*. 2014;1(1):e000042.
81. Bove DG, Overgaard D, Lomborg K, Lindhardt BØ, Midtgaard J. Efficacy of a minimal home-based psychoeducative intervention versus usual care for managing anxiety and dyspnoea in patients with severe chronic obstructive pulmonary disease: a randomised controlled trial protocol. *BMJ Open*. 2015;5(7):e008031.
82. Bove DG, Lomborg K, Jensen AK, Overgaard D, Lindhardt BØ, Midtgaard J. Efficacy of a minimal home-based psychoeducative intervention in patients with advanced COPD: a randomised controlled trial. *Respir Med*. 2016;121:109-116.
83. Usmani ZA, Carson KV, Heslop K, Esterman AJ, De Soyza A, Smith BJ. Psychological therapies for the treatment of anxiety disorders in chronic obstructive pulmonary disease. *Cochrane Database Syst Rev*. 2017;3: CD010673. doi:10.002/14651858.CD010673.pub2.
84. Cafarella P, Effing T, Frith P. An evaluation of the needs of carers of people with COPD. *Eur Res J*. 2012;40(Suppl 56).
85. Zyban [package insert]. Research Triangle Park, NC: GlaxoSmithKline; 2017. Available at: [https://www.gsksource.com/pharma/content/dam/GlaxoSmithKline/US/en/Prescribing\\_Information/Zyban/pdf/ZYBAN-PI-MG.PDF](https://www.gsksource.com/pharma/content/dam/GlaxoSmithKline/US/en/Prescribing_Information/Zyban/pdf/ZYBAN-PI-MG.PDF). Accessed June 2017.
86. Gelenberg AJ, Freeman MP, Markowitz JC, et al. Practice guideline for the treatment of patients with major depressive disorder. *Am J Psychiatry*. 2010;167(10):1.
87. Pary R, Matuschka PR, Lewis S, Lippmann S. Managing bipolar depression. *Psychiatry (Edgmont)*. 2006;3(2):30-41.
88. Blier P. Atypical antipsychotics for mood and anxiety disorders: safe and effective adjuncts? *J Psychiatry Neurosci*. 2005;30(4):232-233.
89. Vulink NC, Figee M, Denys D. Review of atypical antipsychotics in anxiety. *Eur Neuropsychopharmacol*. 2011;21(6):429-449.
90. Locke AB, Kirst N, Shultz CG. Diagnosis and management of generalized anxiety disorder and panic disorder in adults. *Am Fam Physician*. 2015;91(9):617-624.
91. Kew KM, Dias S, Cates CJ. Long-acting inhaled therapy (beta-agonists, anticholinergics and steroids) for COPD: a network meta-analysis. *Cochrane Database Syst Rev*. 2014;(3):CD010844. doi:10.1002/14651858.CD010844.pub2.
92. Scully C. Drug effects on salivary glands: dry mouth. *Oral Dis*. 2003;9(4):165-176.
93. Grenard JL, Munjas BA, Adams JL, et al. Depression and medication adherence in the treatment of chronic diseases in the United States: a meta-analysis. *J Gen Intern Med*. 2011;26(10):1175-1182.
94. O'Donnell DE, Webb KA, Harle I, Neder JA. Pharmacological management of breathlessness in COPD: recent advances and hopes for the future. *Expert Rev Respir Med*. 2016;10(7):823-834.
95. Hyun MK, Lee NR, Jang EJ, Yim JJ, Lee CH. Effect of inhaled drugs on anxiety and depression in patients with chronic obstructive pulmonary disease: a prospective observational study. *Int J Chron Obstruct Pulmon Dis*. 2016;11:747-754.