

1-MINUTE CONSULT

ZIAD SAYEDAHMAD, MD

Beaumont Hospital, Department of Internal Medicine, Royal Oak, MI

FAHED DARMOCH, MD

Department of Internal Medicine, Cleveland Clinic

YASSER AL-KHADRA, MD

Department of Internal Medicine, Cleveland Clinic

AMJAD KABACH, MD

Department of Cardiovascular Medicine, Creighton University, School of Medicine, Omaha, NE

M. CHADI ALRAIES, MD

Department of Cardiovascular Medicine, Wayne State University/Detroit Medical Center, Detroit, MI



BRIEF ANSWERS
TO SPECIFIC
CLINICAL
QUESTIONS

Q: Does early repolarization on ECG increase the risk of cardiac death in healthy people?




Figure 1. Early repolarization with and without QRS notch or slur.

A: No. The early repolarization pattern on electrocardiography (ECG) in asymptomatic patients is nearly always a benign incidental finding. However, in a patient with a history of idiopathic ventricular fibrillation or a family history of sudden cardiac death, the finding warrants further evaluation.

DEFINING EARLY REPOLARIZATION

Published studies differ in their definitions of the early repolarization pattern. In 2016, Patton et al described it as ST-segment elevation in the absence of chest pain, with terminal QRS slur or terminal QRS notch.¹ However, Mcfarlane et al² described it as a J-point elevation of at least 0.1 mV in 2 or more contiguous leads on 12-lead ECG, excluding leads V₁ to V₃, with the presence of terminal QRS notch or slur and QRS duration less than 120 msec. They defined the J point as either the peak of QRS notch or the beginning of QRS slur (Figure 1).² J-point elevation and QRS notch or slur are most commonly seen in left lateral leads and less often in inferior leads.

The early repolarization pattern may mimic patterns seen in myocardial infarction, pericarditis, ventricular aneurysm, hyperkalemia,

and hypothermia,^{1,3} and misinterpreting the pattern can lead to unnecessary laboratory testing, imaging, medication use, and hospital admissions. On the other hand, misinterpreting it as benign in the presence of certain features of the history or clinical presentation can delay the diagnosis and treatment of a potentially critical condition.

PREVALENCE AND MECHANISMS

The prevalence of the early repolarization pattern in the general population ranges from 5% to 15%; the wide range reflects differences in the definition, as well as variability in the pattern of early repolarization over time.⁴

The early repolarization pattern is more commonly seen in African American men and in young, physically active individuals.³ In one study, it was observed in 15% of cases of idiopathic ventricular fibrillation and sudden cardiac death, especially in people ages 35 to 45.⁴ While there is evidence of a heritable basis in the general population, a family history of early repolarization is not known to increase the risk of sudden cardiac death.

A proposed mechanism for the early repolarization pattern is an imbalance in the ion channel system, resulting in variable refrac-

In patients without symptoms, early repolarization is nearly always a benign incidental finding

doi:10.3949/cjcm.86a.17032

TABLE 1

Early repolarization: High-risk features

J-point distribution	Lateral distribution is deemed low-risk compared with inferior and global distribution across all leads
J-wave amplitude	Amplitude ≥ 0.2 mV
ST-segment morphology	Horizontal or downsloping ST segment
QRS notching	Notching of the terminal portion of the QRS is more prevalent in patients with idiopathic ventricular fibrillation

toriness of multiple myocardial regions and varying excitability in the myocardium. This can produce a voltage gradient between myocardial regions, which is believed to cause the major hallmarks of the early repolarization pattern, ie, ST-segment elevation and QRS notching or slurring.³

Although the mechanistic basis of ventricular arrhythmia in patients with early repolarization is still incompletely understood, certain associations may help define the ECG

phenotype that suggests increased risk of sudden cardiac death (Table 1).

MANAGEMENT

The early repolarization pattern is nearly always a benign incidental finding on ECG, with no specific signs or symptoms attributed to it. High-risk features on ECG are associated with a modest increase in absolute risk of sudden cardiac death and warrant clinical correlation.

In the absence of syncope or family history of sudden cardiac death, early repolarization does not merit further workup.²

In patients with a history of unexplained syncope and a family history of sudden cardiac death, early repolarization should be considered in overall risk stratification.¹ Early repolarization in a patient with previous idiopathic ventricular fibrillation warrants referral for electrophysiologic study and, if indicated, insertion of an implantable cardiac defibrillator for secondary prevention.⁵

REFERENCES

1. Patton KK, Ellinor PT, Ezekowitz M, et al; American Heart Association Electrocardiography and Arrhythmias Committee of the Council on Clinical Cardiology and Council on Functional Genomics and Translational Biology. Electrocardiographic early repolarization: a scientific statement from the American Heart Association. *Circulation* 2016; 133(15):1520–1529. doi:10.1161/CIR.0000000000000388
2. Macfarlane PW, Antzelevitch C, Haissaguerre M, et al. The early repolarization pattern: a consensus paper. *J Am Coll Cardiol* 2015; 66(4):470–477. doi:10.1016/j.jacc.2015.05.033
3. Benito B, Guasch E, Rivard L, Nattel S. Clinical and mechanistic issues in early repolarization of normal variants and lethal arrhythmia syndromes. *J Am Coll Cardiol* 2010; 56(15):1177–1186. doi:10.1016/j.jacc.2010.05.037
4. Maury P, Rollin A. Prevalence of early repolarisation/J wave patterns in the normal population. *J Electrocardiol* 2013; 46(5):411–416. doi:10.1016/j.jelectrocard.2013.06.014
5. Mahida S, Sacher F, Berte B, et al. Evaluation of patients with early repolarization syndrome. *J Atr Fibrillation* 2014; 7(3):1083. doi:10.4022/jafib.1083

ADDRESS: M. Chadi Alraies, MD, Wayne State University, Detroit Medical Center, 311 Mack Avenue, Detroit, MI 48201; alraies@hotmail.com