

## How should diabetic dyslipidemia be treated?

SEPTEMBER 2019

**TO THE EDITOR:** The case presented by Hsueh and colleagues<sup>1</sup> is interesting and illustrative. The options for fibric acid derivatives are equally listed as gemfibrozil and fenofibrate. It should be noted, however, that current multisociety guidelines recommend statin treatment for most patients with diabetes,<sup>2</sup> and fenofibrate is the preferred fibric acid derivative to use in combination with a statin. Gemfibrozil has been associated with a higher risk of muscle-related toxicity when combined with statin therapy due to inhibitory effects on the statin metabolic pathway and subsequent increases in plasma statin concentrations.<sup>3</sup> US Food and Drug Administration labeling includes this precaution and states that the benefits of combination use of gemfibrozil and statins do not outweigh the risks.

TAHER MODARRESSI, MD  
Diabetes & Endocrine Associates of Hunterdon  
Hunterdon Medical Center  
Flemington, NJ

### ■ REFERENCES

1. Hsueh YC, Chou CL, Lee TI. Diabetic dyslipidemia with eruptive xanthoma. *Cleve Clin J Med* 2019; 86(9):575–576. doi:10.3949/ccjm.86a.18104
2. Grundy SM, Stone NJ, Bailey AL, et al. 2018 AHA/ACC/AACVPR/AAPA/ABC/ACPM/ADA/AGS/ APhA/ASPC/NLA/PCNA guideline on the management of blood cholesterol: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Circulation* 2019; 139(25):e1082–e1143. doi:10.1161/CIR.0000000000000625
3. Wiggins BS, Saseen JJ, Page RL, et al. Recommendations for management of clinically significant drug-drug interactions with statins and select agents used in patients with cardiovascular disease: a scientific statement from the American Heart Association. *Circulation* 2016; 134(21):e468–e495. PMID:27754879.

doi:10.3949/ccjm.87c.01003

**IN REPLY:** We agree that statin therapy is first-line treatment for primary prevention of atherosclerotic cardiovascular disease for patients with diabetes mellitus who are 40 to 75 years of age.<sup>1</sup> However, severe hypertriglyceridemia (fasting triglycerides  $\geq$  500 mg/dL and especially  $>$  1,000 mg/dL) in diabetic patients, such as our patient,<sup>2</sup> may warrant pharmacologic therapy with fibric acid derivatives, fish oil, or both to reduce the risk of acute pancreatitis.<sup>3</sup> Thus, lifestyle modifications, glycemic control with oral hypoglycemic agents, and fenofibrate therapy were initially prescribed to our patient.<sup>2</sup>

TING-I LEE, MD, PhD  
Head, Division of Endocrinology  
and Metabolism  
Taipei Medical University  
Director, Division of Endocrinology  
and Metabolism  
Wan Fang Hospital  
Taipei 116, Taiwan, ROC

### ■ REFERENCES

1. Grundy SM, Stone NJ, Bailey AL, et al. 2018 AHA/ACC/AACVPR/AAPA/ABC/ACPM/ADA/AGS/ APhA/ASPC/NLA/PCNA guideline on the management of blood cholesterol: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Circulation* 2019; 139(25):e1082–e1143. doi:10.1161/CIR.0000000000000625
2. Hsueh YC, Chou CL, Lee TI. Diabetic dyslipidemia with eruptive xanthoma. *Cleve Clin J Med* 2019; 86(9):575–576. doi:10.3949/ccjm.86a.18104
3. American Diabetes Association. Pharmacologic approaches to glycemic treatment: standards of medical care in diabetes—2019. *Diabetes Care* 2019; 42(suppl 1):S90–S102. doi:10.2337/dc19 S009

doi:10.3949/ccjm.87c.01004