

TABLE OF CONTENTS

FROM THE EDITOR

Celiac disease in the ‘nonclassic’ patient **327**

Adherence to a strict gluten-free diet is not easy, and even strict adherence does not guarantee relief of all symptoms. We need to be as certain as possible that trying the diet makes sense—and that is the real challenge.

Brian F. Mandell, MD, PhD

THE CLINICAL PICTURE

Eruption of erythema after dieting **335**

During the past 4 weeks, the patient had been increasing his exercise and was on a strict diet for bodybuilding.

Li-wen Zhang, MD; Wen-ju Wang, MD; Tao Chen, MD, PhD; Rong-hua Xu, MD

THE CLINICAL PICTURE

Finger nodules: Tip of the gouty iceberg **337**

Recurrent, dramatic flares are the cardinal symptom of gouty disease, but some patients exhibit tophi on physical examination or imaging—without flares or pain.

Brian F. Mandell, MD, PhD

SMART TESTING **CME MOC**

Serial serum lipase testing after the initial diagnostic workup for inpatients with acute pancreatitis: What is the evidence? **341**

Beyond the initial diagnostic workup, the results of serial testing may be misleading, with the potential for adverse effects on patient care and increases in healthcare spending.

Samantha Jill Magier, MD, MEng; Thiruvengadam Muniraj, MD, FRCP; Naseema Merchant, MBBS, FCCP, FACP, FHM

1-MINUTE CONSULT

Should urine antigen testing for *Legionella pneumophila* be ordered for all hospitalized patients with community-acquired pneumonia? **345**

Testing is recommended if the pneumonia is severe, if there has been recent travel, and if there is currently an outbreak of legionnaires disease.

Anna Cheek, MD; Ian Jackson, MD; Manasa Velagapudi, MBBS; Shraddha Narechania, MD, FCCP

CONTINUED ON PAGE 326

Upcoming Features

- Amiodarone and thyrotoxicosis
- Metformin for weight loss in adults with obesity
- Atypical hyperplasia of the breast: Management strategies



CLEVELAND
CLINIC
JOURNAL OF
MEDICINE

CONTINUED FROM PAGE 324

1-MINUTE CONSULT

Celiac disease: Who should I test, and how? 349

First-degree relatives of patients with celiac disease and patients with dermatitis herpetiformis, type 1 diabetes mellitus, and autoimmune thyroid disease are among those for whom testing is advisable.

Jason Nasser, MD; Claire Jansson-Knodell, MD; Alberto Rubio Tapia, MD

GUIDELINES TO PRACTICE

Guidelines for the management of trigeminal neuralgia 355

Care pathways for patients with trigeminal neuralgia vary widely. The most recent UK guidelines emphasize the need for evidence-based care plans for multidisciplinary management.

Mun Seng Chong, MBBS, MD, FRCP; Anish Bahra, MBChB, MD, FRCP; Joanna M. Zakrzewska, BDS, MB BChir, MD, FDSRCS, FFDRCSI, FFPM RCA, FHEA

REVIEW



Measuring exhaled nitric oxide when diagnosing and managing asthma 363

The authors examine the role of this biomarker of airway inflammation and provide guidance for its appropriate use.

Payal Sen, MD; Sumita B. Khatri, MD; Vickram Tejwani, MD

REVIEW

CME MOC

Deadly drug rashes: Early recognition and multidisciplinary care 373

An illustrated review of 4 severe drug rashes. Early recognition and prompt withdrawal of the causative agent are crucial.

Valerie Jaroenpuntaruk, MD; Adam Gray, MD

LETTERS TO THE EDITOR

Chondrodermatitis nodularis helioides 333

Li-wen Zhang, MD; Juan Wu, MD, PhD; Tao Chen, MD, PhD

In Reply: Chondrodermatitis nodularis helioides 333

Maggie So, MD, FACP; Randall S. Edson, MD, MACP, FIDSA

DEPARTMENTS

CME Calendar 330

CME/MOC Instructions 382