Table of Contents

FROM THE EDITOR

Diabetes: Putting off until tomorrow what could happen today can be good 715

Prevention of beta-cell loss and preservation of endogenous insulin production have been attractive treatment prerogatives in type 1 diabetes.

Brian F. Mandell, MD, PhD

THE CLINICAL PICTURE

Trident sign in osmotic demyelination syndrome 732

Brain magnetic resonance imaging showed a confluent trident-shaped lesion in the central pons 1 week after correction of the patient's chronic hyponatremia.

Erin Equinozzi, MSc; Clement D. Lee, MD, MSc; Pamela Schaefer, MD

THE CLINICAL PICTURE

Acute transient phlebitis after a morphine infusion 734

Prominent ascending erythema outlining the infused vein and its proximal branches developed quickly after infusion.

Alexandros Makris, MD, PhD, MSc; Alexandros Nikas, MD; Anastasios Bontozis, MD

THE CLINICAL PICTURE

Skin manifestations of mpox 736

The patient presented with a tender, ulcerated perianal lesion and scattered papules on the trunk and extremities.

James Kim, MD; Matt Biondi, MD; Helen Bartels, MB, BChir

1-MINUTE CONSULT

How should I incorporate emergency contraception counseling into my practice? 738

Emergency contraception counseling should be provided routinely to women at risk of unintended pregnancy as part of counseling on reproductive goals.

Alexa R. Wilden, MD; Heather M. Wainstein, MD

1-MINUTE CONSULT

Should I consider low-dose rivaroxaban (2.5 mg) for my patient with peripheral artery disease? 742

For patients with high ischemic risk, rivaroxaban 2.5 mg twice daily with aspirin 81 or 100 mg daily is a proven strategy to reduce major cardiovascular and limb events.

Pranay Marlecha, MD; Deborah Hornacek, MD; Pulkit Chaudhury, MD, MS

1-MINUTE CONSULT

How can I select the right thyroid hormone formulation for my patient with hypothyroidism? 745

Knowing the features of the available thyroid hormone formulations can provide a starting point for individualizing therapy for patients who have concerns about product formulations or remain symptomatic despite a normal thyroid-stimulating hormone level.

Lea El Hage, MD; Vinni Makin, MBBS, MD

CONTINUED ON PAGE 714

CONTINUED FROM PAGE 711

SYMPTOMS TO DIAGNOSIS



Finding the cause of severe hypokalemia: A 4-step approach 748

This case illustrates the approach to a group of conditions associated with hypokalemia that usually pose a diagnostic challenge.

Enmanuel A. Leiva-Murillo, MD; Erwin Bechtold-Javier, MD; Pol Candela-Parrilla, MD; Ana Matas-García, MD, PhD

REVIEW CME MOC

Latent autoimmune diabetes in adults: Not type 1, not type 2, a little of both 757

Latent autoimmune diabetes in adults shares features of type 1 and type 2 diabetes, often leading to misdiagnosis and a delay in starting needed insulin therapy.

Michelle D. Lundholm, MD; Keren Zhou, MD

EDITORIAL

LADA: A case for starting insulin therapy at diagnosis 764

The author discusses factors that support starting insulin therapy at diagnosis in all patients with latent autoimmune diabetes in adults.

Byron D. Hoogwerf, MD

LETTERS TO THE EDITOR

Barriers to cancer screening in transgender women 721

Maryam Sattari, MD, MS

All menopausal hormone therapy is not alike 721

Henry H. Lindner, MD

In Reply: All menopausal hormone therapy is not alike 722

Samantha F. Butts, MD, MSCE

Cystatin C for estimating glomerular filtration rate and guiding medication dosing 722

Shahzeb Saeed, MD

Corticosteroids in severe community-acquired pneumonia 723

Enmanuel A. Leiva-Murillo, MD; Rosanel Amaro-Rodríguez, MD, PhD

In Reply: Corticosteroids in severe community-acquired pneumonia 724

Alejandro Chapa-Rodriguez, MD; Tark Abou-Elmagd, MD; Cody O'Rear, DO; Shraddha Narechania, MD

DEPARTMENTS

CME Calendar 710

CME/MOC Instructions 767